

Coordinate Systems

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Basis of a Vector Space

Recall:

Let H be a subspace of a vector space V . A set \mathcal{B} of vectors in V is a **basis for H** if:

- ▶ \mathcal{B} is a **linearly independent set**, and
- ▶ the subspace spanned by \mathcal{B} equals H (ie, **$\text{Span } \mathcal{B} = H$**).

The Uniqueness Representation Theorem

One nice result of choosing a basis \mathcal{B} for a vector space V is to impose a **coordinate system** on V .

Example

- If $V = \mathbb{R}^4$ with standard basis $\mathcal{B} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4\}$, then

$$\begin{bmatrix} 2 \\ -5 \\ 17 \\ -8 \end{bmatrix} = 2\mathbf{e}_1 - 5\mathbf{e}_2 + 17\mathbf{e}_3 - 8\mathbf{e}_4.$$

- Consider the vector space $V = \text{Mat}_{2 \times 2}$ and the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$.

Then an element in $\text{Mat}_{2 \times 2}$ such as $\begin{bmatrix} 2 & -5 \\ 17 & -8 \end{bmatrix}$ can be written as the linear combination:

$$\begin{bmatrix} 2 & -5 \\ 17 & -8 \end{bmatrix} = 2\mathbf{v}_1 - 5\mathbf{v}_2 + 17\mathbf{v}_3 - 8\mathbf{v}_4.$$

The Uniqueness Representation Theorem

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then for each \mathbf{x} in V , there exists a **unique set of scalars** c_1, c_2, \dots, c_n (our coordinates) such that

$$\mathbf{x} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + \dots + c_n\mathbf{b}_n. \quad (1)$$

The Uniqueness Representation Theorem

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then for each \mathbf{x} in V , there exists a **unique set of scalars** c_1, c_2, \dots, c_n (our coordinates) such that

$$\mathbf{x} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + \dots + c_n\mathbf{b}_n. \quad (1)$$

Proof. Since \mathcal{B} spans V , we know there exists scalars such that equation (1) above holds. Suppose the representation is not unique, and there exist scalars d_1, d_2, \dots, d_n such that

$$\mathbf{x} = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n.$$

The Uniqueness Representation Theorem

Theorem

Let $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ be a basis for a vector space V . Then for each \mathbf{x} in V , there exists a **unique set of scalars** c_1, c_2, \dots, c_n (our coordinates) such that

$$\mathbf{x} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + \dots + c_n\mathbf{b}_n. \quad (1)$$

Proof. Since \mathcal{B} spans V , we know there exists scalars such that equation (1) above holds. Suppose the representation is not unique, and there exist scalars d_1, d_2, \dots, d_n such that

$$\mathbf{x} = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n.$$

Subtracting the two gives

$$\mathbf{0} = (c_1 - d_1)\mathbf{b}_1 + (c_2 - d_2)\mathbf{b}_2 + \dots + (c_n - d_n)\mathbf{b}_n. \quad (2)$$

Since we assumed \mathcal{B} is a basis for V , equation (2) has only the trivial solution, so $c_j = d_j$ for all $1 \leq j \leq n$. Thus we see the representation is **unique**. \square

Coordinates Relative to a Basis

Definition

Suppose $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a basis for a vector space V , and \mathbf{x} is in V . The **coordinates of \mathbf{x} relative to the basis \mathcal{B}** (or the **\mathcal{B} -coordinates of \mathbf{x}**) are the unique scalars c_1, c_2, \dots, c_n such that $\mathbf{x} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + \dots + c_n\mathbf{b}_n$.

Definition

If c_1, c_2, \dots, c_n are the \mathcal{B} -coordinates of \mathbf{x} , then the vector in \mathbb{R}^n

$$[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}$$

is the **\mathcal{B} -coordinate vector of \mathbf{x}** .

Mapping V to \mathbb{R}^n

Definition

The mapping $V \rightarrow \mathbb{R}^n : \mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}$ is the **coordinate mapping** (determined by \mathcal{B}).

$$\text{Mat}_{2 \times 2} \rightarrow \mathbb{R}^4 : \begin{bmatrix} 2 & -5 \\ 17 & -8 \end{bmatrix} \mapsto \begin{bmatrix} 2 \\ -5 \\ 17 \\ -8 \end{bmatrix}$$

Example

Let $V = \mathbb{R}^2$ and consider
the vector $\mathbf{x} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$.

Consider two different bases for V :

$$\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{B}_2 = \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

Example

Let $V = \mathbb{R}^2$ and consider the vector $\mathbf{x} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$.

We have $\mathbf{x} = 3\mathbf{e}_1 - 5\mathbf{e}_2$, so

$$[\mathbf{x}]_{\mathcal{B}_1} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}.$$

Consider two different bases for V :

$$\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{B}_2 = \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

We solve the vector equation $c_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$.

This system of linear equations has augmented matrix


$$\begin{bmatrix} 1 & -1 & 3 \\ -1 & 1 & -5 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 4 \\ 0 & 1 & -1 \end{bmatrix}$$

Thus we have $[\mathbf{x}]_{\mathcal{B}_2} = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$ since $\mathbf{x} = \begin{bmatrix} 3 \\ -5 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ -1 \end{bmatrix} - 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Geometric Interpretation

Let $V = \mathbb{R}^2$ and consider the vector $\mathbf{x} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$.

Consider two different bases for V : $\mathcal{B}_1 = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ and $\mathcal{B}_2 = \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$.



We have $\begin{bmatrix} 3 \\ -5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} - 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$$[\mathbf{x}]_{\mathcal{B}_1} = \begin{bmatrix} 3 \\ -5 \end{bmatrix}$$

We have $\begin{bmatrix} 3 \\ -5 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ -1 \end{bmatrix} - 1 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

$$[\mathbf{x}]_{\mathcal{B}_2} = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$$

Change of Coordinates Matrix for Basis in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Suppose vector \mathbf{x} in \mathbb{R}^2 has \mathcal{B} -coordinates $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ -6 \end{bmatrix}$.

How can we determine what the coordinates of this vector are in the standard basis?

Change of Coordinates Matrix for Basis in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Suppose vector \mathbf{x} in \mathbb{R}^2 has \mathcal{B} -coordinates $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ -6 \end{bmatrix}$.

How can we determine what the coordinates of this vector are in the standard basis?

$$2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} - 6 \begin{bmatrix} -3 \\ 4 \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -6 \end{bmatrix} = \begin{bmatrix} 20 \\ -28 \end{bmatrix}$$

Change of Coordinates Matrix for Basis in \mathbb{R}^n

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Suppose vector \mathbf{x} in \mathbb{R}^2 has \mathcal{B} -coordinates $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ -6 \end{bmatrix}$.

How can we determine what the coordinates of this vector are in the standard basis?

$$2 \begin{bmatrix} 1 \\ -2 \end{bmatrix} - 6 \begin{bmatrix} -3 \\ 4 \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -6 \end{bmatrix} = \begin{bmatrix} 20 \\ -28 \end{bmatrix}$$

The matrix $P = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} = [\mathbf{b}_1 \quad \mathbf{b}_2]$ is called the **change of coordinates matrix** from \mathcal{B} to the standard basis in \mathbb{R}^2 .

In general, if $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a basis for \mathbb{R}^n , then the **change of coordinates matrix** from \mathcal{B} to the standard basis in \mathbb{R}^n is given by the matrix $P_{\mathcal{B}} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \dots \quad \mathbf{b}_n]$.

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Let $\mathbf{x} = \begin{bmatrix} 8 \\ -11 \end{bmatrix}$. Find the \mathcal{B} -coordinates of \mathbf{x} .

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Let $\mathbf{x} = \begin{bmatrix} 8 \\ -11 \end{bmatrix}$. Find the \mathcal{B} -coordinates of \mathbf{x} .

- ▶ We have the change of coordinate matrix $P = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix}$ which maps $[\mathbf{x}]_{\mathcal{B}}$ to the standard basis.
- ▶ We want to do the inverse of this map (map standard basis to $[\mathbf{x}]_{\mathcal{B}}$).
- ▶ The inverse matrix P^{-1} of P will do the job.

Let $V = \mathbb{R}^2$ and consider the basis $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \begin{bmatrix} -3 \\ 4 \end{bmatrix} \right\}$.

Let $\mathbf{x} = \begin{bmatrix} 8 \\ -11 \end{bmatrix}$. Find the \mathcal{B} -coordinates of \mathbf{x} .

- ▶ We have the change of coordinate matrix $P = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix}$ which maps $[\mathbf{x}]_{\mathcal{B}}$ to the standard basis.
- ▶ We want to do the inverse of this map (map standard basis to $[\mathbf{x}]_{\mathcal{B}}$).
- ▶ The inverse matrix P^{-1} of P will do the job.

In this example, we have

$$P = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix} \quad \text{and} \quad P^{-1} = \begin{bmatrix} -2 & -1.5 \\ -1 & -0.5 \end{bmatrix}.$$

This gives

$$[\mathbf{x}]_{\mathcal{B}} = P^{-1}\mathbf{x} = \begin{bmatrix} -2 & -1.5 \\ -1 & -0.5 \end{bmatrix} \begin{bmatrix} 8 \\ -11 \end{bmatrix} = \begin{bmatrix} 0.5 \\ -2.5 \end{bmatrix}.$$

Coordinate Mappings from V into \mathbb{R}^n

Given a vector space V with basis $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$, we can define the coordinate map

$$T : V \rightarrow \mathbb{R}^n : \mathbf{x} \mapsto [\mathbf{x}]_{\mathcal{B}}.$$

The coordinate mapping above is a **one-to-one** linear transformation from V **onto** \mathbb{R}^n .
A linear transformation that is **both** one-to-one and onto is called an **isomorphism**.

Thus, V and \mathbb{R}^n have the “**same shape**” with respect to vector space properties (ie, vector addition and scalar multiplication).

Example

Let $V = \mathbb{P}_3$ with basis $\mathcal{B} = \{1, t, t^2, t^3\}$. Then the coordinate mapping onto \mathbb{R}^4 is given by

$$T(1) = \mathbf{e}_1, T(t) = \mathbf{e}_2, T(t^2) = \mathbf{e}_3, \text{ and } T(t^3) = \mathbf{e}_4.$$

Find the \mathcal{B} -coordinates of the vector (polynomial) $\mathbf{x} = 2 + 2t^2 - 5t^3$.

Example

Let $V = \mathbb{P}_3$ with basis $\mathcal{B} = \{1, t, t^2, t^3\}$. Then the coordinate mapping onto \mathbb{R}^4 is given by

$$T(1) = \mathbf{e}_1, T(t) = \mathbf{e}_2, T(t^2) = \mathbf{e}_3, \text{ and } T(t^3) = \mathbf{e}_4.$$

Find the \mathcal{B} -coordinates of the vector (polynomial) $\mathbf{x} = 2 + 2t^2 - 5t^3$.

We have $T(\mathbf{x}) = T(2 + 0t + 2t^2 - 5t^3) = 2\mathbf{e}_1 + 0\mathbf{e}_2 + 2\mathbf{e}_3 - 5\mathbf{e}_4$, which gives $[\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 0 \\ 2 \\ -5 \end{bmatrix}$.

Let $V = \mathbb{P}_3$ with basis $\mathcal{B} = \{1, t, t^2, t^3\}$. Determine whether polynomials $\mathbf{v}_1 = 2 + 2t^2 - 5t^3$, $\mathbf{v}_2 = 5 + t^3$, $\mathbf{v}_3 = 7t - 3t^2$, and $\mathbf{v}_4 = -1 + 7t + t^2 - 11t^3$ form a linearly independent set.

Let $V = \mathbb{P}_3$ with basis $\mathcal{B} = \{1, t, t^2, t^3\}$. Determine whether polynomials $\mathbf{v}_1 = 2 + 2t^2 - 5t^3$, $\mathbf{v}_2 = 5 + t^3$, $\mathbf{v}_3 = 7t - 3t^2$, and $\mathbf{v}_4 = -1 + 7t + t^2 - 11t^3$ form a linearly independent set.

$$\text{We have } [\mathbf{v}_1]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 0 \\ 2 \\ -5 \end{bmatrix}, [\mathbf{v}_2]_{\mathcal{B}} = \begin{bmatrix} 5 \\ 0 \\ 0 \\ 1 \end{bmatrix}, [\mathbf{v}_3]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 7 \\ -3 \\ 0 \end{bmatrix}, [\mathbf{v}_4]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 7 \\ 1 \\ -11 \end{bmatrix}.$$

Since \mathbb{P}_3 and \mathbb{R}^4 are **isomorphic**,

$\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ is lin indep in \mathbb{P}_3 **if and only if** $\{[\mathbf{v}_1]_{\mathcal{B}}, [\mathbf{v}_2]_{\mathcal{B}}, [\mathbf{v}_3]_{\mathcal{B}}, [\mathbf{v}_4]_{\mathcal{B}}\}$ is lin indep in \mathbb{R}^4 .

Let $V = \mathbb{P}_3$ with basis $\mathcal{B} = \{1, t, t^2, t^3\}$. Determine whether polynomials $\mathbf{v}_1 = 2 + 2t^2 - 5t^3$, $\mathbf{v}_2 = 5 + t^3$, $\mathbf{v}_3 = 7t - 3t^2$, and $\mathbf{v}_4 = -1 + 7t + t^2 - 11t^3$ form a linearly independent set.

$$\text{We have } [\mathbf{v}_1]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 0 \\ 2 \\ -5 \end{bmatrix}, [\mathbf{v}_2]_{\mathcal{B}} = \begin{bmatrix} 5 \\ 0 \\ 0 \\ 1 \end{bmatrix}, [\mathbf{v}_3]_{\mathcal{B}} = \begin{bmatrix} 0 \\ 7 \\ -3 \\ 0 \end{bmatrix}, [\mathbf{v}_4]_{\mathcal{B}} = \begin{bmatrix} -1 \\ 7 \\ 1 \\ -11 \end{bmatrix}.$$

Since \mathbb{P}_3 and \mathbb{R}^4 are isomorphic,

$\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ is lin indep in \mathbb{P}_3 if and only if $\{[\mathbf{v}_1]_{\mathcal{B}}, [\mathbf{v}_2]_{\mathcal{B}}, [\mathbf{v}_3]_{\mathcal{B}}, [\mathbf{v}_4]_{\mathcal{B}}\}$ is lin indep in \mathbb{R}^4 .

$$\begin{bmatrix} 2 & 5 & 0 & -1 \\ 0 & 0 & 7 & 7 \\ 2 & 0 & -3 & 1 \\ -5 & 1 & 0 & -11 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Setting the free variable $x_4 = 1$, we obtain:

$$-2[\mathbf{v}_1]_{\mathcal{B}} + 1[\mathbf{v}_2]_{\mathcal{B}} - 1[\mathbf{v}_3]_{\mathcal{B}} + 1[\mathbf{v}_4]_{\mathcal{B}} = \mathbf{0}.$$

So we have $-2(2 + 2t^2 - 5t^3) + 1(5 + t^3) - 1(7t - 3t^2) + 1(-1 + 7t + t^2 - 11t^3) = 0$.