

The Inverse of a Matrix

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$2x = 7$$

$$x = \frac{7}{2}$$

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$2x = 7$$

$$\frac{1}{2}2x = \frac{1}{2}7$$

$$x = \frac{7}{2}$$

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$\begin{aligned}2x &= 7 \\ \frac{1}{2}2x &= \frac{1}{2}7 \\ x &= \frac{7}{2}\end{aligned}$$

Multiplying by $\frac{1}{2}$ is the **inverse** of the operation of multiplying by 2.

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$\begin{aligned}2x &= 7 \\ \frac{1}{2}2x &= \frac{1}{2}7 \\ x &= \frac{7}{2}\end{aligned}$$

Multiplying by $\frac{1}{2}$ is the **inverse** of the operation of multiplying by 2.

We can solve $ax = b$ (a, b real) whenever a has a **multiplicative inverse**,

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$\begin{aligned}2x &= 7 \\ \frac{1}{2}2x &= \frac{1}{2}7 \\ x &= \frac{7}{2}\end{aligned}$$

Multiplying by $\frac{1}{2}$ is the **inverse** of the operation of multiplying by 2.

We can solve $ax = b$ (a, b real) whenever a has a **multiplicative inverse**, which is when $a \neq 0$.

The Inverse of an Operation

Consider solving the following linear equation with one variable:

$$\begin{aligned}2x &= 7 \\ \frac{1}{2}2x &= \frac{1}{2}7 \\ x &= \frac{7}{2}\end{aligned}$$

Multiplying by $\frac{1}{2}$ is the **inverse** of the operation of multiplying by 2.

We can solve $ax = b$ (a, b real) whenever a has a **multiplicative inverse**, which is when $a \neq 0$.

Can we solve the **matrix equation** $A\mathbf{x} = \mathbf{b}$ in a similar way?

The Inverse of a Matrix

Let A denote a **square** $n \times n$ matrix. The **inverse** of A (if it exists) is denoted A^{-1} and it is the unique matrix such that

- ▶ $AA^{-1} = I_n$ and
- ▶ $A^{-1}A = I_n$.

For example if $A = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix}$, then $A^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{5} \end{bmatrix}$.

Check:

$$AA^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{5} \end{bmatrix} =$$

Solving the Matrix Equation $A\mathbf{x} = \mathbf{b}$

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the matrix equation $A\mathbf{x} = \mathbf{b}$ has the **unique solution** $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

Solving the Matrix Equation $A\mathbf{x} = \mathbf{b}$

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the matrix equation $A\mathbf{x} = \mathbf{b}$ has the **unique solution** $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

First, we show that $A^{-1}\mathbf{b}$ is a solution:

$$A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}.$$

Solving the Matrix Equation $A\mathbf{x} = \mathbf{b}$

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the matrix equation $A\mathbf{x} = \mathbf{b}$ has the **unique solution** $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

First, we show that $A^{-1}\mathbf{b}$ is a solution:

$$A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}.$$

Second, we show the solution is unique. Suppose \mathbf{y} is a solution. Then $A\mathbf{y} = \mathbf{b}$. Multiplying both sides on the left by A^{-1} and simplifying, we obtain:

$$A^{-1}(A\mathbf{y}) = A^{-1}\mathbf{b} \Rightarrow (A^{-1}A)\mathbf{y} = A^{-1}\mathbf{b} \Rightarrow I_n\mathbf{y} = A^{-1}\mathbf{b} \Rightarrow \mathbf{y} = A^{-1}\mathbf{b}.$$



Properties of Inverses

If A is an invertible matrix, then A^{-1} is invertible and its inverse is

$$(A^{-1})^{-1} = A.$$

Why?

- ▶ A matrix that is **not invertible** is called a **singular matrix**.
- ▶ A matrix that **is invertible** is called a **nonsingular matrix**.

Properties of Inverses

If A and B are $n \times n$ invertible matrices, then AB is an $n \times n$ invertible matrix. The inverse $(AB)^{-1}$ is obtained by taking the **product of inverses in the reverse order**:

$$(AB)^{-1} = B^{-1}A^{-1}.$$

Why?

Properties of Inverses

If A is an invertible matrix, then so is A^T , and the inverse is

$$(A^T)^{-1} = (A^{-1})^T.$$

Why?

Elementary Matrices

An **elementary matrix** is one that is obtained by performing an elementary row operation on an identity matrix.

$$E_1 = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \quad E_3 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

Elementary Row Operations via Elementary Matrices

Theorem

If an elementary row operation is performed on an $m \times n$ matrix A , the resulting matrix can be written as EA , where the $m \times m$ matrix E is an elementary matrix created by performing the same row operation on I_m .

$$E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3} \quad A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{bmatrix}_{3 \times 4} \quad E_2 A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -13 & -14 & -15 & -16 \\ 9 & 10 & 11 & 12 \end{bmatrix}_{3 \times 4}$$

Elementary Matrices are Invertible

Every elementary matrix E is invertible, and the inverse E^{-1} is formed by applying the inverse elementary row operation to I_m .

$$E_1 = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} \quad E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \quad E_3 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

Write out matrices to represent E_1^{-1} , E_2^{-1} , and E_3^{-1} .

RREF of an Invertible Matrix

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n .

Proof.

RREF of an Invertible Matrix

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n .

Proof.

(\Rightarrow) Suppose that A is invertible. Then $A\mathbf{x} = \mathbf{b}$ is solvable for every \mathbf{b} in \mathbb{R}^n . Thus, there must be a pivot in every row of A . Since A is square, the RREF of A is I_n .

RREF of an Invertible Matrix

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n .

Proof.

(\Rightarrow) Suppose that A is invertible. Then $A\mathbf{x} = \mathbf{b}$ is solvable for every \mathbf{b} in \mathbb{R}^n . Thus, there must be a pivot in every row of A . Since A is square, the RREF of A is I_n .

(\Leftarrow) Suppose that the RREF of A is I_n . Then there is a sequence R_1, R_2, \dots, R_p of elementary row operations that transforms A into I_n . Let E_i be the elementary matrix corresponding to R_i . Then

$$E_p E_{p-1} \dots E_2 E_1 A = I_n$$



RREF of an Invertible Matrix

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n .

Proof.

(\Rightarrow) Suppose that A is invertible. Then $A\mathbf{x} = \mathbf{b}$ is solvable for every \mathbf{b} in \mathbb{R}^n . Thus, there must be a pivot in every row of A . Since A is square, the RREF of A is I_n .

(\Leftarrow) Suppose that the RREF of A is I_n . Then there is a sequence R_1, R_2, \dots, R_p of elementary row operations that transforms A into I_n . Let E_i be the elementary matrix corresponding to R_i . Then

$$\underbrace{E_p E_{p-1} \dots E_2 E_1}_{A^{-1}} A = I_n$$



An Algorithm for Finding A^{-1}

Row reduce the augmented matrix $[A \mid I_n]$.

- ▶ If RREF of A is I_n , then RREF of $[A \mid I_n]$ is $[I_n \mid A^{-1}]$.
- ▶ Otherwise, A does not have an inverse.

$$\begin{aligned} \left[\begin{array}{ccc|ccc} 0 & 1 & -2 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 5 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 0 & 1 & -2 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & \frac{1}{5} \end{array} \right] \rightarrow \\ \left[\begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & 0 & \frac{2}{5} \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & \frac{1}{5} \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & \frac{2}{5} \\ 0 & 0 & 1 & 0 & 0 & \frac{1}{5} \end{array} \right] \end{aligned}$$

Practice

If possible, find the inverse of the given matrix.

a) $\begin{bmatrix} 3 & 6 \\ 1 & 2 \end{bmatrix}$

b) $\begin{bmatrix} -2 & 4 \\ -3 & 1 \end{bmatrix}$

Inverse of a 2×2 Matrix

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. How does A fail to be invertible?

Inverse of a 2×2 Matrix

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. How does A fail to be invertible? If one row is a multiple of the other!

$$\begin{bmatrix} a \\ b \end{bmatrix} = k \begin{bmatrix} c \\ d \end{bmatrix} \Leftrightarrow \begin{matrix} a = kc \\ b = kd \end{matrix} \Rightarrow \frac{a}{c} = k = \frac{b}{d} \Rightarrow ad = bc \Leftrightarrow ad - bc = 0$$

Inverse of a 2×2 Matrix

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. How does A fail to be invertible? If one row is a multiple of the other!

$$\begin{bmatrix} a \\ b \end{bmatrix} = k \begin{bmatrix} c \\ d \end{bmatrix} \Leftrightarrow \begin{matrix} a = kc \\ b = kd \end{matrix} \Rightarrow \frac{a}{c} = k = \frac{b}{d} \Rightarrow ad = bc \Leftrightarrow ad - bc = 0$$

(Need to consider some special cases: c or d is 0, and $(c, d) = 0(a, b)$.)

Inverse of a 2×2 Matrix

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. How does A **fail** to be invertible? If one row is a multiple of the other!

$$\begin{bmatrix} a \\ b \end{bmatrix} = k \begin{bmatrix} c \\ d \end{bmatrix} \Leftrightarrow \begin{matrix} a = kc \\ b = kd \end{matrix} \Rightarrow \frac{a}{c} = k = \frac{b}{d} \Rightarrow ad = bc \Leftrightarrow ad - bc = 0$$

(Need to consider some special cases: c or d is 0, and $(c, d) = 0(a, b)$.)

Theorem

A is **not invertible** if and only if $ad - bc = 0$.

Inverse of a 2×2 Matrix

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. How does A **fail** to be invertible? If one row is a multiple of the other!

$$\begin{bmatrix} a \\ b \end{bmatrix} = k \begin{bmatrix} c \\ d \end{bmatrix} \Leftrightarrow \begin{matrix} a = kc \\ b = kd \end{matrix} \Rightarrow \frac{a}{c} = k = \frac{b}{d} \Rightarrow ad = bc \Leftrightarrow ad - bc = 0$$

(Need to consider some special cases: c or d is 0, and $(c, d) = 0(a, b)$.)

Theorem

A is **not invertible** if and only if $ad - bc = 0$.

If A is **invertible**, then

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Practice

If possible, find the inverse of the given matrix.

a)
$$\begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$$

b)
$$\begin{bmatrix} 2 & 0 & 1 \\ 0 & 1 & 3 \\ -4 & 0 & -2 \end{bmatrix}$$

Solving Systems of Linear Equations

Solve the system of equations

$$\begin{aligned}x_1 + 3x_3 &= 2 \\ -2x_2 - 2x_3 &= 1 \\ x_1 - 3x_2 + x_3 &= 0\end{aligned}$$

The equation above has matrix equation $A\mathbf{x} = \mathbf{b}$ given by

$$\begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

Solving Systems of Linear Equations

Solve the system of equations

$$\begin{aligned}x_1 + 3x_3 &= 2 \\ -2x_2 - 2x_3 &= 1 \\ x_1 - 3x_2 + x_3 &= 0\end{aligned}$$

The equation above has matrix equation $A\mathbf{x} = \mathbf{b}$ given by

$$\begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

We compute the solution by $\mathbf{x} = A^{-1}\mathbf{b}$:

$$\mathbf{x} = \begin{bmatrix} 4 & 4.5 & -3 \\ 1 & 1 & -1 \\ -1 & -1.5 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 12.5 \\ 3 \\ -3.5 \end{bmatrix}$$

Solving Systems with Inverse Matrices

1. Write the system of linear equations as a matrix equation $A\mathbf{x} = \mathbf{b}$.
2. Find A^{-1} (if possible).
3. Multiply both sides of the equation in step (1) on the left by A^{-1} .
4. The solution is $\mathbf{x} = A^{-1}\mathbf{b}$.

Note. If the number of equations and variables are not equal, then A will not be a square matrix, and it will not be possible to find A^{-1} . Use row reduction on the augmented matrix instead.

Solving Systems with Inverse Matrices

1. Write the system of linear equations as a matrix equation $A\mathbf{x} = \mathbf{b}$.
2. Find A^{-1} (if possible).
3. Multiply both sides of the equation in step (1) on the left by A^{-1} .
4. The solution is $\mathbf{x} = A^{-1}\mathbf{b}$.

Note. If the number of equations and variables are not equal, then A will not be a square matrix, and it will not be possible to find A^{-1} . Use row reduction on the augmented matrix instead.

How does solving $A\mathbf{x} = \mathbf{b}$ with inverse matrices **compare** to finding the RREF of the augmented matrix?