

Linear Transformations

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Interpretations of the Matrix Equation $A\mathbf{x} = \mathbf{b}$

- ▶ Our focus has been on using matrices and vectors to solve systems of linear equations.
- ▶ We have applied these techniques to determine whether a vector is in the span of a set of vectors.
- ▶ Now we will interpret the matrix-vector product $A\mathbf{x}$ as a **function mapping from one space into another.**

Interpretations of the Matrix Equation $A\mathbf{x} = \mathbf{b}$

- ▶ Our focus has been on using matrices and vectors to solve systems of linear equations.
- ▶ We have applied these techniques to determine whether a vector is in the span of a set of vectors.
- ▶ Now we will interpret the matrix-vector product $A\mathbf{x}$ as a **function mapping from one space into another**.

Example

- ▶ $f : x \mapsto x^2$ maps from $\mathbb{R} \rightarrow \mathbb{R}$.
- ▶ $f : x \mapsto 7x$ maps from $\mathbb{R} \rightarrow \mathbb{R}$.
- ▶ $f : \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} 7x_1 \\ 7x_2 \end{bmatrix}$ maps from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$.
- ▶ $f : \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} x_2 \\ x_1 \\ x_1 + x_2 \end{bmatrix}$ maps from $\mathbb{R}^2 \rightarrow \mathbb{R}^3$.

Transformations from \mathbb{R}^n to \mathbb{R}^m

A **transformation** (also called a function or mapping) T from \mathbb{R}^n to \mathbb{R}^m is a rule that assigns each vector \mathbf{x} in \mathbb{R}^n to a vector $T(\mathbf{x})$ in \mathbb{R}^m .

- ▶ The set of inputs, \mathbb{R}^n , is called the **domain** of T .
- ▶ The space where the inputs are mapped to, \mathbb{R}^m , is called the **codomain** of T .
- ▶ For each \mathbf{x} in the domain, the vector $T(\mathbf{x})$ in \mathbb{R}^m is called the **image** of \mathbf{x} .
- ▶ The set of all images (outputs) is called the **range** of T .

Transformations from \mathbb{R}^n to \mathbb{R}^m

A **transformation** (also called a function or mapping) T from \mathbb{R}^n to \mathbb{R}^m is a rule that assigns each vector \mathbf{x} in \mathbb{R}^n to a vector $T(\mathbf{x})$ in \mathbb{R}^m .

- ▶ The set of inputs, \mathbb{R}^n , is called the **domain** of T .
- ▶ The space where the inputs are mapped to, \mathbb{R}^m , is called the **codomain** of T .
- ▶ For each \mathbf{x} in the domain, the vector $T(\mathbf{x})$ in \mathbb{R}^m is called the **image** of \mathbf{x} .
- ▶ The set of all images (outputs) is called the **range** of T .

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 + x_2 \\ x_2 \\ 0 \end{bmatrix}$$

Matrix Transformations

Let A be an $m \times n$ matrix. Then we can define a transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ where

$$T: \mathbf{x} \mapsto A\mathbf{x}.$$

Matrix Transformations

Let A be an $m \times n$ matrix. Then we can define a transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ where

$$T: \mathbf{x} \mapsto A\mathbf{x}.$$

1. $A = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$. $T(\mathbf{x}) = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 7x_1 \\ 7x_2 \end{bmatrix}$ maps from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Matrix Transformations

Let A be an $m \times n$ matrix. Then we can define a transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ where

$$T: \mathbf{x} \mapsto A\mathbf{x}.$$

1. $A = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$. $T(\mathbf{x}) = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 7x_1 \\ 7x_2 \end{bmatrix}$ maps from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$.

2. $T: \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \mapsto \begin{bmatrix} x_2 \\ x_1 \\ x_1 + x_2 \end{bmatrix}$ maps from $\mathbb{R}^2 \rightarrow \mathbb{R}^3$.

$$T(\mathbf{x}) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_2 \\ x_1 \\ x_1 + x_2 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 1 & -3 & 2 \\ -4 & 1 & 0 \\ 5 & 3 & -9 \end{bmatrix}$,

and let T be the matrix transformation $T: \mathbf{x} \mapsto A\mathbf{x}$.

1. Find the image of the vectors $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, and $\mathbf{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

Example

Let $A = \begin{bmatrix} 1 & -3 & 2 \\ -4 & 1 & 0 \\ 5 & 3 & -9 \end{bmatrix}$,

and let T be the matrix transformation $T: \mathbf{x} \mapsto A\mathbf{x}$.

1. Find the image of the vectors $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, and $\mathbf{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

2. Is $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ in the range of T ? If so, what is the preimage?

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies:

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ **for all** vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .
2. $T(c\mathbf{v}) = c(T(\mathbf{v}))$ **for all** scalars c and vectors \mathbf{v} in \mathbb{R}^n .

Linear Transformations

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called a **linear transformation** if it satisfies:

1. $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$ **for all** vectors \mathbf{u} and \mathbf{v} in \mathbb{R}^n .
2. $T(c\mathbf{v}) = c(T(\mathbf{v}))$ **for all** scalars c and vectors \mathbf{v} in \mathbb{R}^n .

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 + x_2 \\ x_2 \\ 0 \end{bmatrix} \text{ is a linear transformation.}$$

Matrix Transformations are Linear Transformations

Theorem

Every **matrix transformation** is a **linear transformation**. Namely, if $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ can be expressed as $T(\mathbf{x}) = A\mathbf{x}$ for some matrix A , then T is a linear transformation.

Proof.

Image of the Zero Vector

Theorem

If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear map, then $T(\mathbf{0}) = \mathbf{0}$.

Proof.

Example

Consider the linear transformation $T : \mathbb{R}^4 \rightarrow \mathbb{R}^4 : \mathbf{x} \mapsto A\mathbf{x}$ where $A = \begin{bmatrix} 4 & -2 & 5 & -5 \\ -9 & 7 & -8 & 0 \\ -6 & 4 & 5 & 3 \\ 5 & -3 & 8 & -4 \end{bmatrix}$.

1. Find all \mathbf{x} such that $T(\mathbf{x}) = \mathbf{0}$.

2. Find all \mathbf{x} (if any) such that $T(\mathbf{x}) = \begin{bmatrix} 7 \\ 5 \\ 9 \\ 7 \end{bmatrix}$.