

Matrix Arithmetic

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Representations of Matrices

- ▶ If A is an $m \times n$ matrix, it has m rows and n columns.
- ▶ The j^{th} column vector is denoted \mathbf{a}_j .
- ▶ We have n column vectors, and each \mathbf{a}_j is in \mathbb{R}^m .
- ▶ The entry in the i^{th} row and j^{th} column is denoted a_{ij} .
- ▶ **Rows First!** First we give the row index, then the column index.
- ▶ **Recall that in Python, indexing starts at 0.**

$$A = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \dots \quad \mathbf{a}_j \quad \dots \quad \mathbf{a}_n] = \begin{bmatrix} a_{11} & a_{12} & \dots & \mathbf{a}_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & \mathbf{a}_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \dots & \mathbf{a}_{ij} & \dots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & \mathbf{a}_{mj} & \dots & a_{mn} \end{bmatrix}_{m \times n}$$

Types of Matrices

- ▶ An $m \times n$ whose entries are all 0 is called a **zero matrix**, $M = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$.
- ▶ A matrix that has the same number of rows and columns is called a **square matrix**.
- ▶ We call the entries along the diagonal of a square matrix the **diagonal entries**.
- ▶ A square matrix whose **off-diagonal** entries are all 0 is called a **diagonal matrix**.
- ▶ An $n \times n$ matrix with each $a_{ij} = 1$ if $i = j$ and $a_{ij} = 0$ if $i \neq j$ is called the **identity matrix** and is denoted I_n .
- ▶ An $n \times n$ matrix for which $a_{ij} = a_{ji}$ for all $1 \leq i, j \leq n$ is called a **symmetric matrix**.

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -8 & 0 & 0 \\ 0 & 0 & 6 & 0 \\ 0 & 0 & 0 & -7 \end{bmatrix}, I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 6 & 0 & 8 \\ 3 & 0 & 11 & 12 \\ 4 & 8 & 12 & 16 \end{bmatrix}$$

Arithmetic with Matrices

- ▶ Let A and B be $m \times n$ matrices of the same size. The **sum** $C = A + B$ is the $m \times n$ matrix whose entries are the sums of the corresponding entries of A and B . In other words, $c_{ij} = a_{ij} + b_{ij}$.
- ▶ Let A denote an $m \times n$ matrix and r a scalar. To compute **scalar multiple** rA we multiply every entry in A by the scalar r . Thus, if $C = rA$, then $c_{ij} = ra_{ij}$.
- ▶ Two matrices A and B are equal if they have the same size and $a_{ij} = b_{ij}$ for all entries.

Arithmetic with Matrices

- ▶ Let A and B be $m \times n$ matrices of the same size. The **sum** $C = A + B$ is the $m \times n$ matrix whose entries are the sums of the corresponding entries of A and B . In other words, $c_{ij} = a_{ij} + b_{ij}$.
- ▶ Let A denote an $m \times n$ matrix and r a scalar. To compute **scalar multiple** rA we multiply every entry in A by the scalar r . Thus, if $C = rA$, then $c_{ij} = ra_{ij}$.
- ▶ Two matrices A and B are equal if they have the same size and $a_{ij} = b_{ij}$ for all entries.

Example

Let $A = \begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 6 \\ -1 & 0 & 5 \end{bmatrix}$. Compute $2A - B$.

Properties of Matrix Addition and Scalar Multiplication

Let A , B , and C be $m \times n$ matrices of the same size, and let r and s denote scalars.

a. $A + B = B + A$

b. $(A + B) + C = A + (B + C)$

c. $A + 0 = A$

d. $r(A + B) = rA + rB$

e. $(r + s)A = rA + sA$

f. $r(sA) = (rs)A$

Matrix Multiplication

Let A be an $m \times n$ matrix and B be an $n \times p$ matrix. Then

$$AB = A [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \dots \quad \mathbf{b}_p] = [A\mathbf{b}_1 \quad A\mathbf{b}_2 \quad \dots \quad A\mathbf{b}_p].$$

Let $A = \begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 & 0 & 5 \\ 6 & 0 & -1 & 2 \\ 1 & 2 & 1 & 1 \end{bmatrix}$. We define the product AB as follows:

$$\begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 6 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -7 \end{bmatrix} \quad \begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \end{bmatrix} \quad \begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 15 \end{bmatrix}$$

$$AB = \begin{bmatrix} 1 & 5 & 3 & -2 \\ -7 & 4 & 3 & 15 \end{bmatrix}$$

Checking Matrix Sizes

Let A be an $m \times n$ matrix and B be an $n \times p$ matrix. The matrix product AB :

- ▶ Is only defined when A has the same number of columns as B has rows.
- ▶ Is undefined when A has a different number of columns as B has rows.
- ▶ Results in an $m \times p$ matrix when it is defined.

inner sizes match, outer sizes give result: $A_{m \times n} B_{n \times p} = (AB)_{m \times p}$.

$$\begin{aligned} AB &= \begin{bmatrix} -1 & 0 & 3 \\ 2 & -2 & 1 \end{bmatrix}_{2 \times 3} \begin{bmatrix} 2 & 1 & 0 & 5 \\ 6 & 0 & -1 & 2 \\ 1 & 2 & 1 & 1 \end{bmatrix}_{3 \times 4} \\ &= \begin{bmatrix} 1 & 5 & 3 & -2 \\ -7 & 4 & 3 & 15 \end{bmatrix}_{2 \times 4} \end{aligned}$$

Is the Operation Defined?

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E = \begin{bmatrix} 4 & 6 \\ 0 & 3 \\ 1 & 0 \end{bmatrix}, \quad F = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

1. For which matrices is **addition** with A defined? [In what order?]
2. For which matrices is **addition** with B defined?
3. For which matrices is **addition** with C defined?
4. For which matrices is **multiplication** with A defined? [In what order?]
5. For which matrices is **multiplication** with B defined?
6. For which matrices is **multiplication** with C defined?

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E = \begin{bmatrix} 4 & 6 \\ 0 & 3 \\ 1 & 0 \end{bmatrix}, \quad F = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

Evaluate each of the products below:

a) AD

b) DA

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E = \begin{bmatrix} 4 & 6 \\ 0 & 3 \\ 1 & 0 \end{bmatrix}, \quad F = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

Evaluate each of the products below:

a) BF

b) FB

c) B^2

$$A = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E = \begin{bmatrix} 4 & 6 \\ 0 & 3 \\ 1 & 0 \end{bmatrix}, \quad F = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$$

Evaluate each of the products below:

a) $A(BC)$

b) $(AB)C$

Finding the ij^{th} Entry

$$AB = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix} = \left[\begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix} \quad \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right] = \begin{bmatrix} 12 & -3 \\ -4 & 5 \\ 4 & 1 \end{bmatrix}$$

$$AB = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1j} & \dots & b_{1p} \\ b_{21} & b_{22} & \dots & b_{2j} & \dots & b_{2p} \\ \vdots & \vdots & & \vdots & & \vdots \\ b_{i1} & b_{i2} & \dots & b_{ij} & \dots & b_{ip} \\ \vdots & \vdots & & \vdots & & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nj} & \dots & b_{np} \end{bmatrix}$$

$$(AB)_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj} = \sum_{k=1}^n a_{ik}b_{kj}.$$

Properties of Matrix Multiplication

Let A , B , and C be matrices for which the indicated sums and products are defined. Let r denote a scalar.

- a. $A(BC) = (AB)C$ (associative law)
- b. $A(B + C) = AB + AC$ (left distributive law)
- c. $(B + C)A = BA + CA$ (right distributive law)
- d. $r(AB) = (rA)B = A(rB)$ (scalar multiplication)
- e. $I_m A = A = A I_m$ (identity for a square $m \times m$ matrix A)

Properties of Matrix Multiplication

Let A , B , and C be matrices for which the indicated sums and products are defined. Let r denote a scalar.

- a. $A(BC) = (AB)C$ (associative law)
- b. $A(B + C) = AB + AC$ (left distributive law)
- c. $(B + C)A = BA + CA$ (right distributive law)
- d. $r(AB) = (rA)B = A(rB)$ (scalar multiplication)
- e. $I_m A = A = A I_m$ (identity for a square $m \times m$ matrix A)

Partial proof for identity.

Note that $I_m = [\mathbf{e}_1 \ \mathbf{e}_2 \ \dots \ \mathbf{e}_m]$. Then

$$A I_m = A[\mathbf{e}_1 \ \mathbf{e}_2 \ \dots \ \mathbf{e}_m] = [A\mathbf{e}_1 \ A\mathbf{e}_2 \ \dots \ A\mathbf{e}_m] = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_m] = A.$$



The Transpose of a Matrix

Let A denote an $m \times n$ matrix. The **transpose** of A is the $n \times m$ matrix, denoted by A^T , whose columns are formed from the corresponding rows of A .

Example

Give the transpose of

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ -5 & -6 & -7 & -8 \\ 9 & 10 & 11 & 12 \end{bmatrix}$$

Properties of Transpose

Let A and B be matrices for which the indicated sums and products are defined. Let r denote a scalar.

a. $(A^T)^T = A$

b. $(A + B)^T = A^T + B^T$

c. $(rA)^T = rA^T$

d. $(AB)^T = B^T A^T$

e. $A^T = A$ if A is $m \times m$ symmetric

Properties of Transpose

Let A and B be matrices for which the indicated sums and products are defined. Let r denote a scalar.

a. $(A^T)^T = A$

b. $(A + B)^T = A^T + B^T$

c. $(rA)^T = rA^T$

d. $(AB)^T = B^T A^T$

e. $A^T = A$ if A is $m \times m$ symmetric

Other proof for identity.

We previously showed that $A_{m \times m} I_m = A$. We wish to show that $I_m A = A$.

$$I_m A = \left((I_m A)^T \right)^T = \left(A^T I_m^T \right)^T = \left(A^T I_m \right)^T = \left(A^T \right)^T = A.$$

