

Diagonalization

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Diagonal Matrices

Diagonal matrices are particularly nice to work with in many contexts, and can simplify calculations considerably.

Consider the matrix $D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{bmatrix}$.

- ▶ We know $\det D = (2)(3)(5) = 30$, and thus D is invertible.
- ▶ We know that D has eigenvalues $\lambda = 2, 3, 5$.
- ▶ We can easily raise D to powers:

$$D^k = \underbrace{DD \dots D}_{k \text{ times}} = \begin{bmatrix} 2^k & 0 & 0 \\ 0 & 3^k & 0 \\ 0 & 0 & 5^k \end{bmatrix}.$$

Example

Recall this example where A is similar to the diagonal matrix B .

$$\underbrace{\begin{bmatrix} -6 & -4 & -1 \\ -3 & -2 & -1 \\ 5 & 3 & 1 \end{bmatrix}}_{P^{-1}} \underbrace{\begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 1 & 1 & 2 \\ -2 & -1 & -3 \\ 1 & -2 & 0 \end{bmatrix}}_P = \underbrace{\begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix}}_B.$$

- A has eigenvalues $\lambda = -1, 3$ (with $\lambda = -1$ having multiplicity 2).

Example

Recall this example where A is similar to the diagonal matrix B .

$$\underbrace{\begin{bmatrix} -6 & -4 & -1 \\ -3 & -2 & -1 \\ 5 & 3 & 1 \end{bmatrix}}_{P^{-1}} \underbrace{\begin{bmatrix} -13 & -8 & -4 \\ 12 & 7 & 4 \\ 24 & 16 & 7 \end{bmatrix}}_A \underbrace{\begin{bmatrix} 1 & 1 & 2 \\ -2 & -1 & -3 \\ 1 & -2 & 0 \end{bmatrix}}_P = \underbrace{\begin{bmatrix} -1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -1 \end{bmatrix}}_B.$$

- ▶ A has eigenvalues $\lambda = -1, 3$ (with $\lambda = -1$ having multiplicity 2).

Definition

An $n \times n$ matrix A is said to be diagonalizable if it is similar to a diagonal matrix D .

- ▶ Not all matrices are diagonalizable.
- ▶ How do we determine if a matrix is diagonalizable?
If so, how can we find an invertible matrix P such that $P^{-1}AP = D$?

Diagonalizable Matrices

Suppose an $n \times n$ matrix A is **diagonalizable**. Then A is similar to a diagonal matrix D , and there is an **invertible matrix** P such that $P^{-1}AP = D$. Equivalently, $AP = PD$.

$$\text{Let } P = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \dots \quad \mathbf{v}_n] = \begin{bmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{bmatrix} \text{ and } D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{bmatrix}.$$

Diagonalizable Matrices

Suppose an $n \times n$ matrix A is **diagonalizable**. Then A is similar to a diagonal matrix D , and there is an **invertible matrix** P such that $P^{-1}AP = D$. Equivalently, $AP = PD$.

$$\text{Let } P = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \dots \quad \mathbf{v}_n] = \begin{bmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{bmatrix} \text{ and } D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{bmatrix}.$$

We compute

$$AP = [A\mathbf{v}_1 \quad A\mathbf{v}_2 \quad \dots \quad A\mathbf{v}_n]$$
$$PD =$$

Diagonalizable Matrices

Suppose an $n \times n$ matrix A is **diagonalizable**. Then A is similar to a diagonal matrix D , and there is an **invertible matrix** P such that $P^{-1}AP = D$. Equivalently, $AP = PD$.

$$\text{Let } P = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \dots \quad \mathbf{v}_n] = \begin{bmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{bmatrix} \text{ and } D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{bmatrix}.$$

We compute

$$\begin{aligned} AP &= [A\mathbf{v}_1 \quad A\mathbf{v}_2 \quad \dots \quad A\mathbf{v}_n] \\ PD &= [d_1\mathbf{v}_1 \quad d_2\mathbf{v}_2 \quad \dots \quad d_n\mathbf{v}_n] \end{aligned}$$

Diagonalizable Matrices

Suppose an $n \times n$ matrix A is **diagonalizable**. Then A is similar to a diagonal matrix D , and there is an **invertible matrix** P such that $P^{-1}AP = D$. Equivalently, $AP = PD$.

$$\text{Let } P = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \dots \quad \mathbf{v}_n] = \begin{bmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{bmatrix} \text{ and } D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{bmatrix}.$$

We compute

$$\begin{aligned} AP &= [A\mathbf{v}_1 \quad A\mathbf{v}_2 \quad \dots \quad A\mathbf{v}_n] \\ PD &= [d_1\mathbf{v}_1 \quad d_2\mathbf{v}_2 \quad \dots \quad d_n\mathbf{v}_n] \end{aligned}$$

Since $A\mathbf{v}_i = d_i\mathbf{v}_i$, d_i is an **eigenvalue** of A with corresponding **eigenvector** \mathbf{v}_i .

Diagonalizable Matrices

Suppose an $n \times n$ matrix A is **diagonalizable**. Then A is similar to a diagonal matrix D , and there is an **invertible matrix** P such that $P^{-1}AP = D$. Equivalently, $AP = PD$.

$$\text{Let } P = [\mathbf{v}_1 \quad \mathbf{v}_2 \quad \dots \quad \mathbf{v}_n] = \begin{bmatrix} v_{11} & v_{12} & \dots & v_{1n} \\ v_{21} & v_{22} & \dots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \dots & v_{nn} \end{bmatrix} \text{ and } D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & d_n \end{bmatrix}.$$

We compute

$$\begin{aligned} AP &= [A\mathbf{v}_1 \quad A\mathbf{v}_2 \quad \dots \quad A\mathbf{v}_n] \\ PD &= [d_1\mathbf{v}_1 \quad d_2\mathbf{v}_2 \quad \dots \quad d_n\mathbf{v}_n] \end{aligned}$$

Since $A\mathbf{v}_i = d_i\mathbf{v}_i$, d_i is an **eigenvalue** of A with corresponding **eigenvector** \mathbf{v}_i .

Since P is **invertible**, the columns of P are **linearly independent**.

Thus, the \mathbf{v}_i s are n linearly independent **eigenvectors** of A .

Diagonalization Theorem

Theorem

An $n \times n$ matrix A is **diagonalizable** if and only if A has n linearly independent eigenvectors.

If A is diagonalizable, then $P^{-1}AP = D$ where:

- ▶ P is an $n \times n$ matrix whose columns are the n linearly independent eigenvectors.
- ▶ D is an $n \times n$ diagonal matrix whose diagonal entries are the eigenvalues of A that correspond, respectively, to the eigenvectors in P .

$$P^{-1}A \underbrace{\begin{bmatrix} | & | & & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_n \\ | & | & & | \end{bmatrix}}_P = \underbrace{\begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}}_D$$

A 2×2 Example

Diagonalize if possible the matrix $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$.

A 2×2 Example

Diagonalize if possible the matrix $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$.

We have

$$\begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 5 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

Thus $\lambda_1 = 5$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\lambda_2 = 4$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

A 2×2 Example

Diagonalize if possible the matrix $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$.

We have

$$\begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 5 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = 4 \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

Thus $\lambda_1 = 5$ with eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\lambda_2 = 4$ with eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

Since the eigenvectors \mathbf{v}_1 and \mathbf{v}_2 are linearly independent, we have

$$P^{-1}AP = \left(\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \right)^{-1} \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & 4 \end{bmatrix} = D.$$

Linear Independence of Eigenvectors

Theorem

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \dots, \lambda_p$ of an $n \times n$ matrix A , then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is **linearly independent**.

Linear Independence of Eigenvectors

Theorem

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \dots, \lambda_p$ of an $n \times n$ matrix A , then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is **linearly independent**.

Proof. Suppose $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is linearly dependent. Thus there are scalars c_i , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such c_i s that are not all zero, choose the scalars such that the **fewest** c_i s are nonzero. Since the \mathbf{v}_i s are eigenvectors and not the zero vector, at least two of the c_i s must be nonzero. Assume that c_j and c_k are nonzero.

Linear Independence of Eigenvectors

Theorem

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \dots, \lambda_p$ of an $n \times n$ matrix A , then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is **linearly independent**.

Proof. Suppose $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is linearly dependent. Thus there are scalars c_i , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such c_i s that are not all zero, choose the scalars such that the **fewest** c_i s are nonzero. Since the \mathbf{v}_i s are eigenvectors and not the zero vector, at least two of the c_i s must be nonzero. Assume that c_j and c_k are nonzero.

Multiplying both sides of equation (1) by A gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Linear Independence of Eigenvectors

Theorem

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \dots, \lambda_p$ of an $n \times n$ matrix A , then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is **linearly independent**.

Proof. Suppose $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is linearly dependent. Thus there are scalars c_i , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such c_i s that are not all zero, choose the scalars such that the **fewest** c_i s are nonzero. Since the \mathbf{v}_i s are eigenvectors and not the zero vector, at least two of the c_i s must be nonzero. Assume that c_j and c_k are nonzero.

Multiplying both sides of equation (1) by A gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Multiplying both sides of equation (1) by λ_j and subtracting from equation (2) gives

$$c_1(\lambda_1 - \lambda_j)\mathbf{v}_1 + c_2(\lambda_2 - \lambda_j)\mathbf{v}_2 + \dots + 0\mathbf{v}_j + \dots + c_p(\lambda_p - \lambda_j)\mathbf{v}_p = \mathbf{0}. \quad (3)$$

Linear Independence of Eigenvectors

Theorem

If $\mathbf{v}_1, \dots, \mathbf{v}_p$ are eigenvectors corresponding to **distinct** eigenvalues $\lambda_1, \dots, \lambda_p$ of an $n \times n$ matrix A , then the set $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is **linearly independent**.

Proof. Suppose $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is linearly dependent. Thus there are scalars c_i , not all zero, such that

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{0}. \quad (1)$$

Among all such c_i s that are not all zero, choose the scalars such that the **fewest** c_i s are nonzero. Since the \mathbf{v}_i s are eigenvectors and not the zero vector, at least two of the c_i s must be nonzero. Assume that c_j and c_k are nonzero.

Multiplying both sides of equation (1) by A gives

$$A(c_1\mathbf{v}_1 + \dots + c_j\mathbf{v}_j + \dots + c_p\mathbf{v}_p) = c_1\lambda_1\mathbf{v}_1 + \dots + c_j\lambda_j\mathbf{v}_j + \dots + c_p\lambda_p\mathbf{v}_p = \mathbf{0}. \quad (2)$$

Multiplying both sides of equation (1) by λ_j and subtracting from equation (2) gives

$$c_1(\lambda_1 - \lambda_j)\mathbf{v}_1 + c_2(\lambda_2 - \lambda_j)\mathbf{v}_2 + \dots + 0\mathbf{v}_j + \dots + c_p(\lambda_p - \lambda_j)\mathbf{v}_p = \mathbf{0}. \quad (3)$$

Note that $c_k(\lambda_k - \lambda_j) \neq 0$ since $\lambda_k \neq \lambda_j$. Thus, we have expressed $\mathbf{0}$ as a linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_p$ with not all scalars 0 that has **fewer** nonzero scalars than what we assumed was the fewest.

Contradiction!



3×3 Example 1

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$.

3×3 Example 1

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$.

1. Find the eigenvalues by solving the characteristic equation $\det(A - \lambda I) = 0$.

$$\det(A - \lambda I) = \det \begin{bmatrix} 2 - \lambda & 0 & 0 \\ 1 & 2 - \lambda & 1 \\ -1 & 0 & 1 - \lambda \end{bmatrix} = (2 - \lambda)^2(1 - \lambda) = 0$$

A has eigenvalues $\lambda = 2$ (with multiplicity 2) and $\lambda = 1$.

3×3 Example 1

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$.

1. Find the eigenvalues by solving the characteristic equation $\det(A - \lambda I) = 0$.

$$\det(A - \lambda I) = \det \begin{bmatrix} 2 - \lambda & 0 & 0 \\ 1 & 2 - \lambda & 1 \\ -1 & 0 & 1 - \lambda \end{bmatrix} = (2 - \lambda)^2(1 - \lambda) = 0$$

A has eigenvalues $\lambda = 2$ (with multiplicity 2) and $\lambda = 1$.

2. If possible, find three linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

2. If possible, find three linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

For $\lambda = 1$ we have:

$$\begin{bmatrix} 2 - 1 & 0 & 0 \\ 1 & 2 - 1 & 1 \\ -1 & 0 & 1 - 1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ -1 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

if $\mathbf{x} = x_3 \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$. Thus the set $\left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$ is a basis
for the eigenspace corresponding to $\lambda = 1$.

2. If possible, find three linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

For $\lambda = 1$ we have:

$$\begin{bmatrix} 2-1 & 0 & 0 \\ 1 & 2-1 & 1 \\ -1 & 0 & 1-1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ -1 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

if $\mathbf{x} = x_3 \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$. Thus the set $\left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$ is a basis
for the eigenspace corresponding to $\lambda = 1$.

For $\lambda = 2$ we have:

$$\begin{bmatrix} 2-2 & 0 & 0 \\ 1 & 2-2 & 1 \\ -1 & 0 & 1-2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

if $\mathbf{x} = x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$. Thus $\left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$
forms a basis for the eigenspace of $\lambda = 2$.

2. If possible, find three linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.

For $\lambda = 1$ we have:

$$\begin{bmatrix} 2-1 & 0 & 0 \\ 1 & 2-1 & 1 \\ -1 & 0 & 1-1 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ -1 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

if $\mathbf{x} = x_3 \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$. Thus the set $\left\{ \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}$ is a basis
for the eigenspace corresponding to $\lambda = 1$.

For $\lambda = 2$ we have:

$$\begin{bmatrix} 2-2 & 0 & 0 \\ 1 & 2-2 & 1 \\ -1 & 0 & 1-2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

if $\mathbf{x} = x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$. Thus $\left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$
forms a basis for the eigenspace of $\lambda = 2$.

3. Construct matrix P from the vectors above.

$$P = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \quad \text{and} \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

4. Construct matrix D from the corresponding eigenvalues.

Checking our Work

1. Find the eigenvalues by solving the characteristic equation $\det(A - \lambda I) = 0$.
2. If possible, find n linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.
3. Construct matrix P from the vectors above.
4. Construct matrix D from the corresponding eigenvalues.
5. Verify your results.

Checking our Work

1. Find the eigenvalues by solving the characteristic equation $\det(A - \lambda I) = \mathbf{0}$.
2. If possible, find n linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = \mathbf{0}$.
3. Construct matrix P from the vectors above.
4. Construct matrix D from the corresponding eigenvalues.
5. Verify your results.

$$P^{-1}AP = \left(\begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \right)^{-1} \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = D.$$

Checking our Work

1. Find the eigenvalues by solving the characteristic equation $\det(A - \lambda I) = 0$.
2. If possible, find n linearly independent eigenvectors by solving $(A - \lambda I)\mathbf{x} = 0$.
3. Construct matrix P from the vectors above.
4. Construct matrix D from the corresponding eigenvalues.
5. Verify your results.

$$P^{-1}AP = \left(\begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \right)^{-1} \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = D.$$

Or you can check that

$$AP = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & -2 \\ -1 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix} \quad PD = \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & -2 \\ -1 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}$$

Computing Powers of a Matrix

Compute A^7 for $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$.

Computing Powers of a Matrix

Compute A^7 for $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$. We previously showed that A is diagonalizable.

Thus, there is an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$.

Computing Powers of a Matrix

Compute A^7 for $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$. We previously showed that A is diagonalizable.

Thus, there is an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$.

Solving for A , we have that $A = PDP^{-1}$. Thus,

$$A^7 = \underbrace{(PDP^{-1})(PDP^{-1}) \dots (PDP^{-1})}_{7 \text{ times}} = \underbrace{PD(P^{-1}P)D(P^{-1}P) \dots (P^{-1}P)DP^{-1}}_{7 \text{ times}} = PD^7P^{-1}$$

Computing Powers of a Matrix

Compute A^7 for $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$. We previously showed that A is diagonalizable.

Thus, there is an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$.

Solving for A , we have that $A = PDP^{-1}$. Thus,

$$\begin{aligned} A^7 &= \underbrace{(PDP^{-1})(PDP^{-1}) \dots (PDP^{-1})}_{7 \text{ times}} = \underbrace{PD(P^{-1}P)D(P^{-1}P) \dots (P^{-1}P)DP^{-1}}_{7 \text{ times}} = PD^7P^{-1} \\ &= \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1^7 & 0 & 0 \\ 0 & 2^7 & 0 \\ 0 & 0 & 2^7 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 128 & 0 & 0 \\ 127 & 128 & 127 \\ -127 & 0 & 1 \end{bmatrix} \end{aligned}$$

Computing Powers of a Matrix

Compute A^7 for $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$. We previously showed that A is diagonalizable.

Thus, there is an invertible matrix P and a diagonal matrix D such that $P^{-1}AP = D$.

Solving for A , we have that $A = PDP^{-1}$. Thus,

$$\begin{aligned} A^7 &= \underbrace{(PDP^{-1})(PDP^{-1}) \dots (PDP^{-1})}_{7 \text{ times}} = \underbrace{PD(\overset{\text{red}}{P^{-1}P})D(\overset{\text{red}}{P^{-1}P}) \dots (\overset{\text{red}}{P^{-1}P})DP^{-1}}_{7 \text{ times}} = PD^7P^{-1} \\ &= \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1^7 & 0 & 0 \\ 0 & 2^7 & 0 \\ 0 & 0 & 2^7 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ -1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 128 & 0 & 0 \\ 127 & 128 & 127 \\ -127 & 0 & 1 \end{bmatrix} \end{aligned}$$

In general, $\overset{\text{red}}{A^k} = (PDP^{-1})^k = P\overset{\text{red}}{D^k}P^{-1}$ for a positive integer k .

3×3 Example 2

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$.

3×3 Example 2

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$. Since A is **triangular**, its eigenvals are 2, 4.

3×3 Example 2

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$. Since A is **triangular**, its eigenvals are 2, 4.

For $\lambda = 4$ we have

$$(A - 4I)\mathbf{x} = \begin{bmatrix} -2 & 4 & 6 \\ 0 & -2 & 2 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \text{ when } \mathbf{x} = x_3 \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}, \text{ thus a basis is } \left\{ \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

3×3 Example 2

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$. Since A is **triangular**, its eigenvals are 2, 4.

For $\lambda = 4$ we have

$$(A - 4I)\mathbf{x} = \begin{bmatrix} -2 & 4 & 6 \\ 0 & -2 & 2 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \text{ when } \mathbf{x} = x_3 \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}, \text{ thus a basis is } \left\{ \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

For $\lambda = 2$ we have

$$(A - 2I)\mathbf{x} = \begin{bmatrix} 0 & 4 & 6 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{bmatrix} \mathbf{x} = \mathbf{0} \text{ when } \mathbf{x} = x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \text{ thus a basis is } \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}.$$

3×3 Example 2

If possible, diagonalize the matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$. Since A is **triangular**, its eigenvals are 2, 4.

For $\lambda = 4$ we have

$$(A - 4I)\mathbf{x} = \begin{bmatrix} -2 & 4 & 6 \\ 0 & -2 & 2 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \text{ when } \mathbf{x} = x_3 \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}, \text{ thus a basis is } \left\{ \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix} \right\}.$$

For $\lambda = 2$ we have

$$(A - 2I)\mathbf{x} = \begin{bmatrix} 0 & 4 & 6 \\ 0 & 0 & 2 \\ 0 & 0 & 2 \end{bmatrix} \mathbf{x} = \mathbf{0} \text{ when } \mathbf{x} = x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \text{ thus a basis is } \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}.$$

Since we do **NOT** have three linearly independent eigenvectors, A is **NOT diagonalizable**.

3×3 Example 3

Determine whether or not $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$ is diagonalizable.

3×3 Example 3

Determine whether or not $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$ is diagonalizable.

- Notice matrix A is **upper triangular**, so A has eigenvalues $\lambda = 3, 6, 2$.

3×3 Example 3

Determine whether or not $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$ is diagonalizable.

- ▶ Notice matrix A is **upper triangular**, so A has eigenvalues $\lambda = 3, 6, 2$.
- ▶ If \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are eigenvectors corresponding to **distinct** eigenvalues λ_1 , λ_2 , and λ_3 , then we proved that the set of eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is **linearly independent**.

3×3 Example 3

Determine whether or not $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$ is diagonalizable.

- ▶ Notice matrix A is **upper triangular**, so A has eigenvalues $\lambda = 3, 6, 2$.
- ▶ If \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are eigenvectors corresponding to **distinct** eigenvalues λ_1 , λ_2 , and λ_3 , then we proved that the set of eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is **linearly independent**.
- ▶ Since A has **three linearly independent eigenvectors**, it must be diagonalizable.

3×3 Example 3

Determine whether or not $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$ is diagonalizable.

- ▶ Notice matrix A is **upper triangular**, so A has eigenvalues $\lambda = 3, 6, 2$.
- ▶ If \mathbf{v}_1 , \mathbf{v}_2 , and \mathbf{v}_3 are eigenvectors corresponding to **distinct** eigenvalues λ_1 , λ_2 , and λ_3 , then we proved that the set of eigenvectors $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is **linearly independent**.
- ▶ Since A has **three linearly independent eigenvectors**, it must be diagonalizable.

Theorem

An $n \times n$ matrix with **n distinct eigenvalues** is diagonalizable.

Matrices with Eigenvalues of Higher Multiplicity

Theorem

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \dots, \lambda_p$.

- (a) For $1 \leq k \leq p$, the **dimension of the eigenspace** for λ_k (called the *geometric multiplicity*) is at most the algebraic multiplicity of the eigenvalue λ_k .

Matrices with Eigenvalues of Higher Multiplicity

Theorem

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \dots, \lambda_p$.

- (a) For $1 \leq k \leq p$, the **dimension of the eigenspace** for λ_k (called the *geometric multiplicity*) is at most the algebraic multiplicity of the eigenvalue λ_k .
- (b) The matrix A is diagonalizable if and only if the sum of the dimensions of the eigenspaces equals n . This happens if and only if:
 - (i) The characteristic polynomial factors completely into linear factors, and
 - (ii) The dimension of the eigenspace for each λ_k **equals** the multiplicity of λ_k .

Matrices with Eigenvalues of Higher Multiplicity

Theorem

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \dots, \lambda_p$.

- (a) For $1 \leq k \leq p$, the **dimension of the eigenspace** for λ_k (called the *geometric multiplicity*) is at most the algebraic multiplicity of the eigenvalue λ_k .
- (b) The matrix A is diagonalizable if and only if the sum of the dimensions of the eigenspaces equals n . This happens if and only if:
 - (i) The characteristic polynomial factors completely into linear factors, and
 - (ii) The dimension of the eigenspace for each λ_k **equals** the multiplicity of λ_k .
- (c) If A is diagonalizable and \mathcal{B}_k is a basis for the eigenspace corresponding to eigenvalue λ_k , then all the vectors in the sets $\mathcal{B}_1, \dots, \mathcal{B}_p$ form an **eigenvector basis** for \mathbb{R}^n .

Reviewing Previous Examples

The matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 2$ with eigenspace dimension equal to 2, and
- ▶ $\lambda = 1$ with eigenspace dimension equal to 1.
- ▶ $2 + 1 = 3$, so A is diagonalizable.

Reviewing Previous Examples

The matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 2$ with eigenspace dimension equal to 2, and
- ▶ $\lambda = 1$ with eigenspace dimension equal to 1.
- ▶ $2 + 1 = 3$, so A is diagonalizable.

The matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 2$ with eigenspace dimension equal to 1, and
- ▶ $\lambda = 4$ with eigenspace dimension equal to 1.
- ▶ $1 + 1 = 2$, so A is NOT diagonalizable.

Reviewing Previous Examples

The matrix $A = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 2$ with eigenspace dimension equal to 2, and
- ▶ $\lambda = 1$ with eigenspace dimension equal to 1.
- ▶ $2 + 1 = 3$, so A is diagonalizable.

The matrix $A = \begin{bmatrix} 2 & 4 & 6 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 2$ with eigenspace dimension equal to 1, and
- ▶ $\lambda = 4$ with eigenspace dimension equal to 1.
- ▶ $1 + 1 = 2$, so A is NOT diagonalizable.

The matrix $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 6 & 2 \\ 0 & 0 & 2 \end{bmatrix}$

has eigenvalues:

- ▶ $\lambda = 3, 6, 2$.
- ▶ Since A has three distinct eigenvalues, it is diagonalizable.

Interpretation of Similarity

Suppose that A is an $n \times n$ diagonalizable matrix: $A = PDP^{-1}$.

The columns of P are bases for the eigenspaces of A , and together form a basis \mathcal{B} for \mathbb{R}^n .

Consider the linear transform $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $T(\mathbf{x}) = A\mathbf{x} = PDP^{-1}\mathbf{x}$.

Interpretation of Similarity

Suppose that A is an $n \times n$ diagonalizable matrix: $A = PDP^{-1}$.

The columns of P are bases for the eigenspaces of A , and together form a basis \mathcal{B} for \mathbb{R}^n .

Consider the linear transform $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $T(\mathbf{x}) = A\mathbf{x} = PDP^{-1}\mathbf{x}$.

P is the change of basis matrix from \mathcal{B} to the standard basis.

P^{-1} is the change of basis matrix from the standard basis to \mathcal{B} .

So to perform T on a vector \mathbf{x} with respect to the standard basis, we can

1. change to the basis \mathcal{B} by multiplying by P^{-1} ,
2. then scale each coordinate by multiplying by D ,
3. then change back to the standard basis by multiplying by P .

Example

Consider the linear transform $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1.5 & 0.5 \\ 1 & 1 \end{bmatrix} \mathbf{x}$.

$$\begin{aligned} A &= PDP^{-1} \\ &= \begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 0.5 \end{bmatrix} \left(\begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \right)^{-1}. \end{aligned}$$

Desmos:

<https://www.desmos.com/calculator/n7l5cf9xjo>

Example

Consider the linear transform $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1.5 & 0.5 \\ 1 & 1 \end{bmatrix} \mathbf{x}$.

$$\begin{aligned} A &= PDP^{-1} \\ &= \begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 0.5 \end{bmatrix} \left(\begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \right)^{-1}. \end{aligned}$$

Desmos:

<https://www.desmos.com/calculator/n7l5cf9xjo>

