

Vector Equations

Linear Algebra


These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Vectors


A **vector** is a quantity that has both a length (also called magnitude) and a direction.

A car is traveling 60 miles per hour directly East.




$$\mathbf{v} = \vec{v} = \begin{bmatrix} 60 \\ 0 \end{bmatrix}$$

A car is traveling 40 miles per hour directly South.



$$\mathbf{w} = \vec{w} = \begin{bmatrix} 0 \\ -40 \end{bmatrix}$$

A car is traveling 64 miles per hour directly Northwest.



$$\mathbf{u} = \vec{u} = \begin{bmatrix} -\frac{64}{\sqrt{2}} \\ \frac{64}{\sqrt{2}} \end{bmatrix}$$

A **vector** (or a **column vector**) is a matrix with only one column.

- ▶ We denote vectors in bold such as \mathbf{v} , and/or
- ▶ Using an arrow superscript such as \vec{v} .

A **vector** (or a **column vector**) is a matrix with only one column.

- ▶ We denote vectors in bold such as \mathbf{v} , and/or
 - ▶ Using an arrow superscript such as \vec{v} .
-
- ▶ The vector $\vec{v} = \begin{bmatrix} -12 \\ 8 \end{bmatrix}$ consists of an ordered pair of two real numbers. We say \vec{v} is in \mathbb{R}^2 (read as “r two” or “r squared”).
 - ▶ If the vector \vec{v} consists of n different real numbers, then we have

$$\vec{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \in \mathbb{R}^n.$$

Equality of Vectors

We say that two vectors $\vec{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ and $\vec{w} = \begin{bmatrix} w_1 \\ \vdots \\ w_n \end{bmatrix}$ are **equal** if:

- ▶ They have both the **same direction and the same length**, or equivalently,
- ▶ All corresponding entries of the two vectors are equal:

$$v_i = w_i \text{ for all } 1 \leq i \leq n.$$

Vector Addition

Given two vectors $\mathbf{u} = \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ in \mathbb{R}^n , we define their sum as the vector $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ \vdots \\ u_n + v_n \end{bmatrix}$.

Vector Addition

Given two vectors $\mathbf{u} = \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ in \mathbb{R}^n , we define their sum as the vector $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ \vdots \\ u_n + v_n \end{bmatrix}$.

Example

Add the vectors.

a) $\mathbf{u} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -6 \\ 2 \end{bmatrix}$.

b) $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \\ 8 \\ 0 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 8 \\ 2 \\ -1 \\ 3 \end{bmatrix}$.

Vector Addition

Given two vectors $\mathbf{u} = \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ in \mathbb{R}^n , we define their sum as the vector $\mathbf{u} + \mathbf{v} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \\ \vdots \\ u_n + v_n \end{bmatrix}$.

Example

Add the vectors.

a) $\mathbf{u} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} -6 \\ 2 \end{bmatrix}$.

b) $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \\ 8 \\ 0 \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} 8 \\ 2 \\ -1 \\ 3 \end{bmatrix}$.

- ▶ We can only add two vectors if they have the same **dimensions**.
- ▶ When we add two vectors, the result is a vector with the same number of dimensions.

Scalar Multiplication

A quantity that has only a magnitude (and does not have a direction) is called a **scalar**. We usually refer to scalars as numbers. For example, -4 , $\sqrt{5}$, 0 , and e^2 are all scalars.

Given a vector $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ and scalar c , we define the **scalar product** of c and \mathbf{v} as $c\mathbf{v} = \begin{bmatrix} c \cdot v_1 \\ c \cdot v_2 \\ \vdots \\ c \cdot v_n \end{bmatrix}$.

Scalar Multiplication

A quantity that has only a magnitude (and does not have a direction) is called a **scalar**. We usually refer to scalars as numbers. For example, -4 , $\sqrt{5}$, 0 , and e^2 are all scalars.

Given a vector $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ and scalar c , we define the **scalar product** of c and \mathbf{v} as $c\mathbf{v} = \begin{bmatrix} c \cdot v_1 \\ c \cdot v_2 \\ \vdots \\ c \cdot v_n \end{bmatrix}$.

Example

Compute the scalar product

a) $c = -3$ and $\mathbf{v} = \begin{bmatrix} 2 \\ -5 \end{bmatrix}$.

b) $c = 2$ and $\mathbf{v} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$.

Scalar Multiplication

A quantity that has only a magnitude (and does not have a direction) is called a **scalar**. We usually refer to scalars as numbers. For example, -4 , $\sqrt{5}$, 0 , and e^2 are all scalars.

Given a vector $\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$ and scalar c , we define the **scalar product** of c and \mathbf{v} as $c\mathbf{v} = \begin{bmatrix} c \cdot v_1 \\ c \cdot v_2 \\ \vdots \\ c \cdot v_n \end{bmatrix}$.

Example

Compute the scalar product

a) $c = -3$ and $\mathbf{v} = \begin{bmatrix} 2 \\ -5 \end{bmatrix}$.

b) $c = 2$ and $\mathbf{v} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$.

- ▶ We can multiply any scalar and any vector.
- ▶ The result is a vector with the same dimension as the original vector.

Geometric Descriptions of Vectors in \mathbb{R}^2

We can think of the vector $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \in \mathbb{R}^2$ as a directed line segment (an arrow).

Consider two vectors \mathbf{u} and $\mathbf{v} \in \mathbb{R}^2$. The sum $\mathbf{u} + \mathbf{v}$ is the result of first moving \mathbf{u} and then moving \mathbf{v} .

Parallelogram Rule for Addition: If \mathbf{u} and \mathbf{v} are in \mathbb{R}^2 , then the sum $\mathbf{u} + \mathbf{v}$ is the diagonal of the parallelogram spanned by \mathbf{u} and \mathbf{v} .

Geometric Descriptions of Vectors in \mathbb{R}^2

We can think of the vector $\mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} \in \mathbb{R}^2$ as a directed line segment (an arrow).

Consider two vectors \mathbf{u} and $\mathbf{v} \in \mathbb{R}^2$. The sum $\mathbf{u} + \mathbf{v}$ is the result of first moving \mathbf{u} and then moving \mathbf{v} .

Consider the vector $\mathbf{u} \in \mathbb{R}^2$ and nonzero scalar c . The scalar product $c\mathbf{u}$ gives a vector parallel to \mathbf{u} whose magnitude has been scaled by a factor of c .

- ▶ If $c > 0$, then $c\mathbf{u}$ has the same direction as \mathbf{u} .
- ▶ If $c < 0$, then $c\mathbf{u}$ has the opposite direction as \mathbf{u} .
- ▶ If $|c| < 1$, then $c\mathbf{u}$ is a compression of \mathbf{u} .
- ▶ If $|c| > 1$, then $c\mathbf{u}$ is a stretching of \mathbf{u} .

Parallelogram Rule for Addition: If \mathbf{u} and \mathbf{v} are in \mathbb{R}^2 , then the sum $\mathbf{u} + \mathbf{v}$ is the diagonal of the parallelogram spanned by \mathbf{u} and \mathbf{v} .

Properties of Vector Arithmetic

The **zero vector** is defined as the vector in \mathbb{R}^n with no magnitude, and it is denoted by

0 or $\vec{0}$. As a result of this definition, it follows that $\mathbf{0} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$.

Properties of Vector Arithmetic

The **zero vector** is defined as the vector in \mathbb{R}^n with no magnitude, and it is denoted by

$\mathbf{0}$ or $\vec{0}$. As a result of this definition, it follows that $\mathbf{0} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$.

For all $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in \mathbb{R}^n and all scalars c and d :

(i) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

(v) $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$

(ii) $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$

(vi) $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$

(iii) $\mathbf{u} + \mathbf{0} = \mathbf{0} + \mathbf{u} = \mathbf{u}$

(vii) $c(d\mathbf{u}) = (cd)\mathbf{u}$

(iv) $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$


(viii) $1\mathbf{u} = \mathbf{u}$

Giving Directions in Box-Grid City

Example

Find constants x_1 and x_2 such that

$$\begin{bmatrix} -4 \\ -5 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$




Giving Directions in Box-Grid City

Example

Find constants x_1 and x_2 such that

$$\begin{bmatrix} -4 \\ -5 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$




The vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are called **standard column vectors**.

Giving Directions in Diagonaland

Example

Find constants x_1 and x_2 such that

$$\begin{bmatrix} -4 \\ -5 \end{bmatrix} = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$




Setting up a System of Equations

Example

Find constants x_1 and x_2 such that

$$\begin{bmatrix} -4 \\ -5 \end{bmatrix} = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$



$$2x_1 - x_2 = -4$$

$$x_1 + x_2 = -5$$

$$\begin{bmatrix} 2 & -1 & -4 \\ 1 & 1 & -5 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & -2 \end{bmatrix}$$

$$\text{Thus, } \begin{cases} x_1 = -3 \\ x_2 = -2 \end{cases}$$

Given any vector \mathbf{w} in \mathbb{R}^2 , we can find constants x_1 and x_2 such that

$$\mathbf{w} = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

Linear Combinations

Given vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ in \mathbb{R}^n and scalars c_1, c_2, \dots, c_p , the vector \mathbf{y} defined by


$$\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \sum_{i=1}^p c_i\mathbf{v}_i$$

is called a **linear combination** of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ with weights c_1, c_2, \dots, c_p .

Can you think of two different vectors in \mathbb{R}^2 such that $\mathbf{w} = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$ **cannot** be written as a linear combination of the two vectors?

For example:

$$\begin{bmatrix} 2 \\ -3 \end{bmatrix} = -\frac{1}{3} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \frac{8}{3} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$



The Span of a Set of Vectors

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ be vectors in \mathbb{R}^n . The set of all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ is called the **subset of \mathbb{R}^n spanned by $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$** .

The set **$\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$** is the collection of all vectors in \mathbb{R}^n that can be written in the form

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \sum_{i=1}^p c_i\mathbf{v}_i.$$

The Span of a Set of Vectors

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ be vectors in \mathbb{R}^n . The set of all linear combinations of the vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$ is called the **subset of \mathbb{R}^n spanned by $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$** .

The set **$\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$** is the collection of all vectors in \mathbb{R}^n that can be written in the form

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \sum_{i=1}^p c_i\mathbf{v}_i.$$

To determine whether a vector \mathbf{y} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$:

- ▶ We try to solve $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_p\mathbf{v}_p = \mathbf{y}$.
- ▶ In other words, we have the augmented matrix

$$\begin{bmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_p & \mathbf{y} \end{bmatrix}.$$

The Span of a Set of Vectors in \mathbb{R}^2

Describe the span of the set of vectors in \mathbb{R}^2 .

a) $\text{Span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$

b) $\text{Span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$

c) $\text{Span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -4 \\ -2 \end{bmatrix} \right\}$

d) $\text{Span} \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$

The Span of a Set of Vectors in \mathbb{R}^3

Describe the span of the set of vectors in \mathbb{R}^3 .

a) $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}$

b) $\text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$

c) $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$

d) $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}$

Practice

Determine whether the given vector is in $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} \right\}$.

a) $\begin{bmatrix} 6 \\ -2 \\ -10 \end{bmatrix}$

b) $\begin{bmatrix} 4 \\ 5 \\ -1 \end{bmatrix}$

Conceptual Practice

Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ and \mathbf{w} be vectors in \mathbb{R}^n . Prove that if \mathbf{w} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ then $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

Proof.

Conceptual Practice

Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ and \mathbf{w} be vectors in \mathbb{R}^n . Prove that if \mathbf{w} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ then $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

Proof.

We need to show that the two sets $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\}$ and $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ are the same.

(\supseteq) Suppose that $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Then there exist scalars c_1, c_2, c_3 such that $\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3$. Then $\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + 0\mathbf{w}$, and hence $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\}$.

Conceptual Practice

Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ and \mathbf{w} be vectors in \mathbb{R}^n . Prove that if \mathbf{w} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ then $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$.

Proof.

We need to show that the two sets $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\}$ and $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ are the same.

(\supseteq) Suppose that $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Then there exist scalars c_1, c_2, c_3 such that $\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3$. Then $\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + 0\mathbf{w}$, and hence $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\}$.

(\subseteq) Suppose that $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\}$. Then there exist scalars c_1, c_2, c_3, d such that $\mathbf{y} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + d\mathbf{w}$. Since \mathbf{w} is in $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$, there exist scalars b_1, b_2, b_3 such that $\mathbf{w} = b_1\mathbf{v}_1 + b_2\mathbf{v}_2 + b_3\mathbf{v}_3$. Substituting for \mathbf{w} into the previous equation, we obtain

$$\begin{aligned}\mathbf{y} &= c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + d\mathbf{w} \\ &= c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 + d(b_1\mathbf{v}_1 + b_2\mathbf{v}_2 + b_3\mathbf{v}_3) \\ &= (c_1 + db_1)\mathbf{v}_1 + (c_2 + db_2)\mathbf{v}_2 + (c_3 + db_3)\mathbf{v}_3.\end{aligned}$$

Since $c_1 + db_1$, $c_2 + db_2$, and $c_3 + db_3$ are scalars, \mathbf{y} is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$, and hence $\mathbf{y} \in \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Thus, $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{w}\} = \text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. □

Parametric Vector Form for Solutions of a Linear System

Find the solution set to the linear system

$$\begin{array}{rclcl} x_1 & +2x_2 & & -4x_5 & = & 0 \\ & & x_3 & +7x_5 & = & 1 \\ & & & x_4 & +5x_5 & = & 2 \end{array}$$

Parametric Vector Form for Solutions of a Linear System

Find the solution set to the linear system

$$\begin{array}{rclcl} x_1 & +2x_2 & & -4x_5 & = & 0 \\ & & x_3 & +7x_5 & = & 1 \\ & & & x_4 & +5x_5 & = & 2 \end{array}$$

$$\left\{ \begin{array}{l} x_1 = -2x_2 + 4x_5 \\ x_2 \text{ is free} \\ x_3 = 1 - 7x_5 \\ x_4 = 2 - 5x_5 \\ x_5 \text{ is free} \end{array} \right.$$

Parametric Vector Form for Solutions of a Linear System

Find the solution set to the linear system

$$\begin{array}{rcl} x_1 + 2x_2 & -4x_5 & = 0 \\ & x_3 + 7x_5 & = 1 \\ & x_4 + 5x_5 & = 2 \end{array}$$

$$\left\{ \begin{array}{l} x_1 = -2x_2 + 4x_5 \\ x_2 \text{ is free} \\ x_3 = 1 - 7x_5 \\ x_4 = 2 - 5x_5 \\ x_5 \text{ is free} \end{array} \right.$$

$$\begin{aligned} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} &= \begin{bmatrix} -2x_2 + 4x_5 \\ x_2 \\ 1 - 7x_5 \\ 2 - 5x_5 \\ x_5 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 2 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} 4 \\ 0 \\ -7 \\ -5 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ 0 \\ 1 \\ 2 \\ 0 \end{bmatrix} + s \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} 4 \\ 0 \\ -7 \\ -5 \\ 1 \end{bmatrix}. \end{aligned}$$

The weights s, t are any scalars. This is called **parametric vector form**.