

Markov Chains

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.


This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Monopoly

Monopoly is a board game where tokens are moved around the board. Players start on “Go”.

On each turn, a player rolls two dice, and then advances their token by the number of spaces equal to the sum of the rolled dice.

What is the most landed on spot in Monopoly?



Simplified Monopoly

We'll use a board with 12 spaces (4 on each side), and roll 1 4-sided die (numbered 1,2,3,4) on each turn.

Players start on space 0 ("Go").

If a player lands on space 9 ("Go to Jail"), then when they leave on their next turn, they go to spaces 4, ..., 7.

Where will a player be after their **first turn**?

0	1	2	3
11			4
10			5
9	8	7	6

Simplified Monopoly

We'll use a board with 12 spaces (4 on each side), and roll 1 4-sided die (numbered 1,2,3,4) on each turn.

Players start on space 0 ("Go").

If a player lands on space 9 ("Go to Jail"), then when they leave on their next turn, they go to spaces 4, ..., 7.

Where will a player be after their **first turn**?

A **state** is the current space a player's token is on.

A **state vector** gives a probability distribution on the states.

0	1	2	3
11			4
10			5
9	8	7	6

$$\mathbf{x}_1 = [0 \quad .25 \quad .25 \quad .25 \quad .25 \quad 0 \quad 0 \quad \dots \quad 0]$$

Simplified Monopoly, continued

Where will a player be after their **second turn** if they start at space 0?

Simplified Monopoly, continued

Where will a player be after their **second turn** if they start at space 0?

Let \mathbf{p}_i give the probability distribution on states after one turn if the player **starts** on space i .

$$\text{Then } \mathbf{x}_2 = x_{11}\mathbf{p}_1 + x_{12}\mathbf{p}_2 + \dots + x_{1n}\mathbf{p}_n$$

Simplified Monopoly, continued

Where will a player be after their **second turn** if they start at space 0?

Let \mathbf{p}_i give the probability distribution on states after one turn if the player **starts** on space i .

Then $\mathbf{x}_2 = x_{11}\mathbf{p}_1 + x_{12}\mathbf{p}_2 + \dots + x_{1n}\mathbf{p}_n = P\mathbf{x}_1$, where $P = [\mathbf{p}_1 \quad \mathbf{p}_2 \quad \dots \quad \mathbf{p}_n]$.

$P = [p_{ij}]$ is the **transition matrix**, where p_{ij} is the prob of transitioning from **state j** to **state i** .

Simplified Monopoly, continued

Where will a player be after their **second turn** if they start at space 0?

Let \mathbf{p}_i give the probability distribution on states after one turn if the player **starts** on space i .

Then $\mathbf{x}_2 = x_{11}\mathbf{p}_1 + x_{12}\mathbf{p}_2 + \dots + x_{1n}\mathbf{p}_n = P\mathbf{x}_1$, where $P = [\mathbf{p}_1 \quad \mathbf{p}_2 \quad \dots \quad \mathbf{p}_n]$.

$P = [p_{ij}]$ is the **transition matrix**, where p_{ij} is the prob of transitioning from **state j** to **state i** .

Switch to Jupyter Notebook for calculations.

Definition

A **Markov chain** is a mathematical model used to describe a random process where:

- ▶ At each discrete time $k \geq 0$, the system is in one of n **states**.
- ▶ There are probabilities that describe how likely it is the system transitions from one state to another. These probabilities depend **only** on the **current state**.

(We will only discuss discrete-time Markov chains with a finite number of states.)

Definition

A **Markov chain** is a mathematical model used to describe a random process where:

- ▶ At each discrete time $k \geq 0$, the system is in one of n **states**.
- ▶ There are probabilities that describe how likely it is the system transitions from one state to another. These probabilities depend **only** on the **current state**.

(We will only discuss discrete-time Markov chains with a finite number of states.)

Since a Markov chain describes a random process, we let the **state vector** \mathbf{x}_k be the probability distribution over which state the system is in at time k .

The **transition matrix** $P = [p_{ij}]$ has p_{ij} = probability of transitioning from **state j** to **state i** .

Definition

A **Markov chain** is a mathematical model used to describe a random process where:

- ▶ At each discrete time $k \geq 0$, the system is in one of n **states**.
- ▶ There are probabilities that describe how likely it is the system transitions from one state to another. These probabilities depend **only** on the **current state**.

(We will only discuss discrete-time Markov chains with a finite number of states.)

Since a Markov chain describes a random process, we let the **state vector** \mathbf{x}_k be the probability distribution over which state the system is in at time k .

The **transition matrix** $P = [p_{ij}]$ has p_{ij} = probability of transitioning from **state j** to **state i** .

The sequence $\mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots$ of state vectors satisfies $\mathbf{x}_{k+1} = P\mathbf{x}_k$ for $k \geq 0$,

Definition

A **Markov chain** is a mathematical model used to describe a random process where:

- ▶ At each discrete time $k \geq 0$, the system is in one of n **states**.
- ▶ There are probabilities that describe how likely it is the system transitions from one state to another. These probabilities depend **only** on the **current state**.

(We will only discuss discrete-time Markov chains with a finite number of states.)

Since a Markov chain describes a random process, we let the **state vector** \mathbf{x}_k be the probability distribution over which state the system is in at time k .

The **transition matrix** $P = [p_{ij}]$ has p_{ij} = probability of transitioning from **state j** to **state i** .

The sequence $\mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots$ of state vectors satisfies $\mathbf{x}_{k+1} = P\mathbf{x}_k$ for $k \geq 0$, and hence

$$\mathbf{x}_{k+1} = P\mathbf{x}_k = P(P\mathbf{x}_{k-1}) = P^2\mathbf{x}_{k-1} = \dots = P^{k+1}\mathbf{x}_0.$$

A Simple Weather Model


- ▶ Imagine that there are two possible states for weather: no rain (sunny) or rain.
- ▶ You can always directly observe the current weather state.
- ▶ The current weather has some bearing on what the next day's weather will be.

You want to be able to predict what the weather will be like tomorrow.

A Simple Weather Model

- ▶ Imagine that there are two possible states for weather: no rain (sunny) or rain.
- ▶ You can always directly observe the current weather state.
- ▶ The current weather has some bearing on what the next day's weather will be.

You want to be able to predict what the weather will be like tomorrow.




We can build a model to predict tomorrow's weather based on the weather **today**.

$$P = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$$

- ▶ We organize the probabilities of transitioning from one state to another in a **transition matrix**.
- ▶ p_{ij} is the probability of transitioning from **state j** to **state i** .
- ▶ This will be a **stochastic matrix**. The entries in each column vector add up to 1.

A Simple Weather Model




Today the weather is **sunny**.

state 1 = **sunny**, state 2 = **rainy**

$$\mathbf{x}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

What is the probability that it **rains** three days from now?

A Simple Weather Model



Today the weather is **sunny**.

state 1 = **sunny**, state 2 = **rainy**

$$\mathbf{x}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$


What is the probability that it **rains** three days from now?

$$\begin{aligned} \mathbf{x}_3 &= P\mathbf{x}_2 = P^2\mathbf{x}_1 = P^3\mathbf{x}_0 \\ &= \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}^3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\ &= \begin{bmatrix} 0.844 \\ 0.156 \end{bmatrix} \end{aligned}$$

So the probability it **rains** in three days, given that today is **sunny**, is 0.156.

Modeling Election Outcomes

Consider the following model:



With initial state vector $\mathbf{x}_0 = \begin{bmatrix} 0.50 \\ 0.45 \\ 0.05 \end{bmatrix}$

1. What is the predicted outcome in the next election cycle?
2. What is the predicted outcome in two election cycles?

$$\begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix}$$

$$\begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} = \begin{bmatrix} 0.369 \\ 0.4965 \\ 0.1345 \end{bmatrix}$$

Long Term Predictions

Often we are interested in making predictions far out in time.

- ▶ Based on migration patterns, what is the long term prediction for the population of a certain species?
- ▶ Based on current infection rates, what is the long term prediction for the total number of infected people?

$$\mathbf{x}_k = P^k \mathbf{x}_0 \quad \text{for large } k$$

Long Term Predictions

Often we are interested in making predictions far out in time.

- ▶ Based on migration patterns, what is the long term prediction for the population of a certain species?
- ▶ Based on current infection rates, what is the long term prediction for the total number of infected people?

$$\mathbf{x}_k = P^k \mathbf{x}_0 \quad \text{for large } k$$

Under some reasonable assumptions, the limiting distribution \mathbf{x}_∞ will satisfy

$$\mathbf{x}_\infty = P\mathbf{x}_\infty.$$

Long Term Predictions

Often we are interested in making predictions far out in time.

- ▶ Based on migration patterns, what is the long term prediction for the population of a certain species?
- ▶ Based on current infection rates, what is the long term prediction for the total number of infected people?

$$\mathbf{x}_k = P^k \mathbf{x}_0 \quad \text{for large } k$$

Under some reasonable assumptions, the limiting distribution \mathbf{x}_∞ will satisfy

$$\mathbf{x}_\infty = P\mathbf{x}_\infty.$$

The limiting distribution is an **eigenvector** of P corresponding to the **eigenvalue 1**!

Long Term Predictions

Often we are interested in making predictions far out in time.

- ▶ Based on migration patterns, what is the long term prediction for the population of a certain species?
- ▶ Based on current infection rates, what is the long term prediction for the total number of infected people?

$$\mathbf{x}_k = P^k \mathbf{x}_0 \quad \text{for large } k$$

Under some reasonable assumptions, the limiting distribution \mathbf{x}_∞ will satisfy

$$\mathbf{x}_\infty = P\mathbf{x}_\infty.$$

The limiting distribution is an **eigenvector** of P corresponding to the **eigenvalue 1**!

The limiting distribution does **not** depend on the initial state vector \mathbf{x}_0 .

Long Term Predictions

1. What is the predicted outcome in the next election cycle?
2. What is the predicted outcome in two election cycles?
3. What is the predicted political demographics in the long-run?

Long Term Predictions

1. What is the predicted outcome in the next election cycle?
2. What is the predicted outcome in two election cycles?
3. What is the predicted political demographics in the long-run?

$$\mathbf{x}_1 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} = \begin{bmatrix} 0.369 \\ 0.4965 \\ 0.1345 \end{bmatrix}$$

Long Term Predictions

1. What is the predicted outcome in the next election cycle?
2. What is the predicted outcome in two election cycles?
3. What is the predicted political demographics in the long-run?

$$\mathbf{x}_1 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} = \begin{bmatrix} 0.369 \\ 0.4965 \\ 0.1345 \end{bmatrix}$$

$$\mathbf{x}_{10} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix}^{10} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.3221 \\ 0.5350 \\ 0.1429 \end{bmatrix} \quad \mathbf{x}_{20} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix}^{20} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

Long Term Predictions

1. What is the predicted outcome in the next election cycle?
2. What is the predicted outcome in two election cycles?
3. What is the predicted political demographics in the long-run?

$$\mathbf{x}_1 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.41 \\ 0.475 \\ 0.115 \end{bmatrix} = \begin{bmatrix} 0.369 \\ 0.4965 \\ 0.1345 \end{bmatrix}$$

$$\mathbf{x}_{10} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix}^{10} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.3221 \\ 0.5350 \\ 0.1429 \end{bmatrix} \quad \mathbf{x}_{20} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix}^{20} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

$$\mathbf{x}_{30} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix}^{30} \begin{bmatrix} 0.5 \\ 0.45 \\ 0.05 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

Steady State Vectors

If P is a stochastic matrix, then a **steady state vector** (or **equilibrium vector**) for P is a **probability vector** \mathbf{q} (nonnegative entries that sum to 1) such that

$$P\mathbf{q} = \mathbf{q}.$$

- ▶ Note that \mathbf{q} is an eigenvector of P corresponding to $\lambda = 1$.
- ▶ Thus if probability vector \mathbf{q} is an eigenvector of $\lambda = 1$, it is a steady state vector.

Steady State Vectors

If P is a stochastic matrix, then a **steady state vector** (or **equilibrium vector**) for P is a **probability vector** \mathbf{q} (nonnegative entries that sum to 1) such that

$$P\mathbf{q} = \mathbf{q}.$$

- ▶ Note that \mathbf{q} is an eigenvector of P corresponding to $\lambda = 1$.
- ▶ Thus if probability vector \mathbf{q} is an eigenvector of $\lambda = 1$, it is a steady state vector.

Elections example

$$\begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

Steady State Vectors

If P is a stochastic matrix, then a **steady state vector** (or **equilibrium vector**) for P is a **probability vector** \mathbf{q} (nonnegative entries that sum to 1) such that

$$P\mathbf{q} = \mathbf{q}.$$

- ▶ Note that \mathbf{q} is an eigenvector of P corresponding to $\lambda = 1$.
- ▶ Thus if probability vector \mathbf{q} is an eigenvector of $\lambda = 1$, it is a steady state vector.

Elections example

$$\begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

Weather example

$$\mathbf{x}_{\infty} = \begin{bmatrix} 0.8333 \\ 0.1667 \end{bmatrix}$$

It is **sunny** 83.3% of the days.

Steady State Vectors

If P is a stochastic matrix, then a **steady state vector** (or **equilibrium vector**) for P is a **probability vector** \mathbf{q} (nonnegative entries that sum to 1) such that

$$P\mathbf{q} = \mathbf{q}.$$

- ▶ Note that \mathbf{q} is an eigenvector of P corresponding to $\lambda = 1$.
- ▶ Thus if probability vector \mathbf{q} is an eigenvector of $\lambda = 1$, it is a steady state vector.

Elections example

$$\begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.2 & 0.8 & 0.3 \\ 0.1 & 0.1 & 0.4 \end{bmatrix} \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix} = \begin{bmatrix} 0.3214 \\ 0.5357 \\ 0.1429 \end{bmatrix}$$

Weather example

$$\mathbf{x}_{\infty} = \begin{bmatrix} 0.8333 \\ 0.1667 \end{bmatrix}$$

It is **sunny** 83.3% of the days.

(Monopoly example in Jupyter Notebook.)

Stochastic Matrices

Theorem

If P is a **stochastic matrix** (columns sum to 1), then 1 is an eigenvalue of P .

Stochastic Matrices

Theorem

If P is a **stochastic matrix** (columns sum to 1), then 1 is an eigenvalue of P .

Proof.

P and P^T have the **same eigenvalues**.

Stochastic Matrices

Theorem

If P is a **stochastic matrix** (columns sum to 1), then 1 is an eigenvalue of P .

Proof.

P and P^T have the **same eigenvalues**.

Since P is a stochastic matrix, then the entries in each row of P^T **add up to 1**.

Stochastic Matrices

Theorem

If P is a **stochastic matrix** (columns sum to 1), then 1 is an eigenvalue of P .

Proof.

P and P^T have the **same eigenvalues**.

Since P is a stochastic matrix, then the entries in each row of P^T **add up to 1**.

Consider the column vector of all 1's that we denote \mathbf{e} . Then we have

$$P^T \mathbf{e} = \begin{bmatrix} \sum_{j=1}^n p_{j1} \\ \sum_{j=1}^n p_{j2} \\ \vdots \\ \sum_{j=1}^n p_{jn} \end{bmatrix} = \mathbf{e}.$$

Therefore, \mathbf{e} is an eigenvector of P^T corresponding to the eigenvalue $\lambda = 1$. □