

Applications of Determinants

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

Using Determinants to Solve Systems of Equations

$$\begin{array}{rclcl} 3x_1 & - & 2x_2 & = & 6 \\ -5x_1 & + & 4x_2 & = & 8 \end{array}$$

For a square matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$ and column vector \mathbf{b} , if we replace the j^{th} column of A with \mathbf{b} , we denote the new matrix

$$A_j(\mathbf{b}) = A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_{j-1} \ \mathbf{b} \ \mathbf{a}_{j+1} \ \dots \ \mathbf{a}_n] .$$

Using Determinants to Solve Systems of Equations

$$\begin{array}{rcl} 3x_1 & - & 2x_2 = 6 \\ -5x_1 & + & 4x_2 = 8 \end{array}$$

For a square matrix $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$ and column vector \mathbf{b} , if we replace the j^{th} column of A with \mathbf{b} , we denote the new matrix

$$A_j(\mathbf{b}) = A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_{j-1} \ \mathbf{b} \ \mathbf{a}_{j+1} \ \dots \ \mathbf{a}_n] .$$

Let $A = \begin{bmatrix} 3 & -2 \\ -5 & 4 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$. Then

$$A_1(\mathbf{b}) = \begin{bmatrix} 6 & -2 \\ 8 & 4 \end{bmatrix} \quad \text{and} \quad A_2(\mathbf{b}) = \begin{bmatrix} 3 & 6 \\ -5 & 8 \end{bmatrix} .$$

Cramer's Rule

Let A be an invertible $n \times n$ matrix. For any \mathbf{b} in \mathbb{R}^n , the unique solution to $A\mathbf{x} = \mathbf{b}$ is \mathbf{x} , where the j^{th} entry of \mathbf{x} is given by

$$x_j = \frac{\det A_j(\mathbf{b})}{\det A} \quad \text{for } j = 1, 2, \dots, n.$$

Cramer's Rule

Let A be an invertible $n \times n$ matrix. For any \mathbf{b} in \mathbb{R}^n , the unique solution to $A\mathbf{x} = \mathbf{b}$ is \mathbf{x} , where the j^{th} entry of \mathbf{x} is given by

$$x_j = \frac{\det A_j(\mathbf{b})}{\det A} \quad \text{for } j = 1, 2, \dots, n.$$

$$\begin{array}{rclcl} 3x_1 & - & 2x_2 & = & 6 \\ -5x_1 & + & 4x_2 & = & 8 \end{array}$$

$$A = \begin{bmatrix} 3 & -2 \\ -5 & 4 \end{bmatrix}$$

$$A_1(\mathbf{b}) = \begin{bmatrix} 6 & -2 \\ 8 & 4 \end{bmatrix}$$

$$A_2(\mathbf{b}) = \begin{bmatrix} 3 & 6 \\ -5 & 8 \end{bmatrix}$$

$$\det A = 2$$

$$\mathbf{x} = \begin{bmatrix} \frac{40}{2} \\ \frac{54}{2} \end{bmatrix} = \begin{bmatrix} 20 \\ 27 \end{bmatrix}$$

$$\det A_1(\mathbf{b}) = 40$$

$$\det A_2(\mathbf{b}) = 54$$

Proof of Cramer's Rule

The unique solution to $A\mathbf{x} = \mathbf{b}$ is \mathbf{x} , where the $x_j = \frac{\det A_j(\mathbf{b})}{\det A}$.

Proof of Cramer's Rule

The unique solution to $A\mathbf{x} = \mathbf{b}$ is \mathbf{x} , where the $x_j = \frac{\det A_j(\mathbf{b})}{\det A}$.

Proof.

Let $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$ be an invertible $n \times n$ matrix, where each \mathbf{a}_i denotes a column vector of A . Let \mathbf{x} be the unique solution to the equation $A\mathbf{x} = \mathbf{b}$. We denote $I_j(\mathbf{x}) = [\mathbf{e}_1 \ \dots \ \mathbf{e}_{j-1} \ \mathbf{x} \ \mathbf{e}_{j+1} \ \dots \ \mathbf{e}_n]$, and consider the product

$$\begin{aligned} AI_j(\mathbf{x}) &= A[\mathbf{e}_1 \ \dots \ \mathbf{e}_{j-1} \ \mathbf{x} \ \mathbf{e}_{j+1} \ \dots \ \mathbf{e}_n] \\ &= [A\mathbf{e}_1 \ \dots \ A\mathbf{e}_{j-1} \ A\mathbf{x} \ A\mathbf{e}_{j+1} \ \dots \ A\mathbf{e}_n] \\ &= [\mathbf{a}_1 \ \dots \ \mathbf{a}_{j-1} \ \mathbf{b} \ \mathbf{a}_{j+1} \ \dots \ \mathbf{a}_n] = A_j(\mathbf{b}). \end{aligned}$$

Proof of Cramer's Rule

The unique solution to $A\mathbf{x} = \mathbf{b}$ is \mathbf{x} , where the $x_j = \frac{\det A_j(\mathbf{b})}{\det A}$.

Proof.

Let $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_n]$ be an invertible $n \times n$ matrix, where each \mathbf{a}_i denotes a column vector of A . Let \mathbf{x} be the unique solution to the equation $A\mathbf{x} = \mathbf{b}$. We denote $I_j(\mathbf{x}) = [\mathbf{e}_1 \ \dots \ \mathbf{e}_{j-1} \ \mathbf{x} \ \mathbf{e}_{j+1} \ \dots \ \mathbf{e}_n]$, and consider the product

$$\begin{aligned} AI_j(\mathbf{x}) &= A[\mathbf{e}_1 \ \dots \ \mathbf{e}_{j-1} \ \mathbf{x} \ \mathbf{e}_{j+1} \ \dots \ \mathbf{e}_n] \\ &= [A\mathbf{e}_1 \ \dots \ A\mathbf{e}_{j-1} \ A\mathbf{x} \ A\mathbf{e}_{j+1} \ \dots \ A\mathbf{e}_n] \\ &= [\mathbf{a}_1 \ \dots \ \mathbf{a}_{j-1} \ \mathbf{b} \ \mathbf{a}_{j+1} \ \dots \ \mathbf{a}_n] = A_j(\mathbf{b}). \end{aligned}$$

$$\begin{aligned} \det A_j(\mathbf{b}) &= \det (AI_j(\mathbf{x})) \\ &= (\det A) (\det I_j(\mathbf{x})) = (\det A) x_j. \end{aligned}$$

Therefore, we have $x_j = \frac{\det A_j(\mathbf{b})}{\det A}$.

Using the multiplicative property of determinants, we have

□

Formula for the Inverse of a Matrix

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$, when $\det A = ad - bc \neq 0$.

Is there a similar formula for A^{-1} for larger matrices?

Formula for the Inverse of a Matrix

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$, when $\det A = ad - bc \neq 0$.

Is there a similar formula for A^{-1} for larger matrices?

Let A be an $n \times n$ matrix. The **adjugate** $\text{adj } A$ of A (also called *classical adjoint*) is defined by

$$\text{adj } A = \begin{bmatrix} C_{11} & C_{21} & \dots & C_{i1} & \dots & C_{n1} \\ C_{12} & C_{22} & \dots & C_{i2} & \dots & C_{n2} \\ \vdots & \vdots & \ddots & \vdots & & \vdots \\ C_{1i} & C_{2i} & \dots & C_{ji} & \dots & C_{ni} \\ \vdots & \vdots & & \vdots & \ddots & \vdots \\ C_{1n} & C_{2n} & \dots & C_{jn} & \dots & C_{nn} \end{bmatrix},$$

where the **cofactor** $C_{ij} = (-1)^{i+j} \det A_{ij}$.

Note the **reversed indices!!!**

The adjugate is the **transpose**
of the matrix of cofactors.

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Proof 1.

Let $D = A^{-1}$. Then $AD = I$. To find the j th column \mathbf{d}_j of D , we solve $A\mathbf{d}_j = \mathbf{e}_j$. To find the i th entry of \mathbf{d}_j , we apply [Cramer's Rule](#):

$$\begin{aligned} i\text{th entry of } \mathbf{d}_j &= \frac{\det A_i(\mathbf{e}_j)}{\det A} = \frac{1}{\det A} \det [\mathbf{a}_1 \quad \dots \quad \mathbf{a}_{i-1} \quad \mathbf{e}_j \quad \mathbf{a}_{i+1} \quad \dots \quad \mathbf{a}_n] \\ &= \frac{1}{\det A} (-1)^{j+i} \det A_{ji} \quad \text{by expanding about column } i \\ &= \frac{1}{\det A} C_{ji}. \end{aligned}$$

Hence, the ij th entry of A^{-1} is $\frac{1}{\det A} C_{ji}$.

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Proof 2.

We show that $A(\operatorname{adj} A) = (\det A)I_n$.

The i, j entry of $A(\operatorname{adj} A)$ is $\sum_{k=1}^n a_{ik}(\operatorname{adj} A)_{kj} = \sum_{k=1}^n a_{ik} C_{jk} = \sum_{k=1}^n a_{ik} (-1)^{j+k} \det A_{jk}$.

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Proof 2.

We show that $A(\operatorname{adj} A) = (\det A)I_n$.

The i, j entry of $A(\operatorname{adj} A)$ is $\sum_{k=1}^n a_{ik}(\operatorname{adj} A)_{kj} = \sum_{k=1}^n a_{ik}C_{jk} = \sum_{k=1}^n a_{ik}(-1)^{j+k} \det A_{jk}$.

This is the determinant of A after replacing the j th row with the i th row, which is:

$\det A$, if $i = j$; 0, if $i \neq j$ (since there is a repeated row).

Formula for the Inverse of a Matrix

Let A be an invertible $n \times n$ matrix. Then $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Proof 2.

We show that $A(\operatorname{adj} A) = (\det A)I_n$.

The i, j entry of $A(\operatorname{adj} A)$ is $\sum_{k=1}^n a_{ik}(\operatorname{adj} A)_{kj} = \sum_{k=1}^n a_{ik}C_{jk} = \sum_{k=1}^n a_{ik}(-1)^{j+k} \det A_{jk}$.

This is the determinant of A after replacing the j th row with the i th row, which is:

$\det A$, if $i = j$; 0, if $i \neq j$ (since there is a repeated row).

Since A is invertible, $A^{-1}A(\operatorname{adj} A) = A^{-1}(\det A)I_n$. Thus, $A^{-1} = \frac{1}{\det A} \operatorname{adj} A$.

Example

Compute the inverse of $A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$.

Example

Compute the inverse of $A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$.

$$\det A = -2 + 0 + 0 + 6 - 6 - 0 = -2.$$

Example

Compute the inverse of $A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$.

$$\det A = -2 + 0 + 0 + 6 - 6 - 0 = -2.$$

$$\operatorname{adj} A = \begin{bmatrix} -8 & -9 & 6 \\ -2 & -2 & 2 \\ 2 & 3 & -2 \end{bmatrix}.$$

Example

Compute the inverse of $A = \begin{bmatrix} 1 & 0 & 3 \\ 0 & -2 & -2 \\ 1 & -3 & 1 \end{bmatrix}$.

$$\det A = -2 + 0 + 0 + 6 - 6 - 0 = -2.$$

$$\operatorname{adj} A = \begin{bmatrix} -8 & -9 & 6 \\ -2 & -2 & 2 \\ 2 & 3 & -2 \end{bmatrix}.$$

$$\begin{aligned} (\operatorname{adj} A)_{21} &= C_{12} = (-1)^{1+2} \det A_{12} \\ &= (-1)^3 [0 \cdot 1 - 1 \cdot (-2)] = -2. \end{aligned}$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A = \frac{1}{-2} \begin{bmatrix} -8 & -9 & 6 \\ -2 & -2 & 2 \\ 2 & 3 & -2 \end{bmatrix} = \begin{bmatrix} 4 & 9/2 & -3 \\ 1 & 1 & -1 \\ -1 & -3/2 & 1 \end{bmatrix}.$$

Volumes

Volume of 2d Parallelogram

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Let $\mathbf{u} = (a, b)$ and $\mathbf{v} = (c, d)$.

The origin, \mathbf{u} , \mathbf{v} , and $\mathbf{u} + \mathbf{v}$ form a parallelogram \mathcal{P} .

Volume of 2d Parallelogram

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Let $\mathbf{u} = (a, b)$ and $\mathbf{v} = (c, d)$.

The origin, \mathbf{u} , \mathbf{v} , and $\mathbf{u} + \mathbf{v}$ form a **parallelogram** \mathcal{P} .

Theorem

The area of the parallelogram \mathcal{P} in \mathbb{R}^2 is $|\det(A)|$.

<https://www.desmos.com/calculator/iksjsllpxz>

Volume of 2d Parallelogram

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Let $\mathbf{u} = (a, b)$ and $\mathbf{v} = (c, d)$.

The origin, \mathbf{u} , \mathbf{v} , and $\mathbf{u} + \mathbf{v}$ form a **parallelogram** \mathcal{P} .

Theorem

The area of the parallelogram \mathcal{P} in \mathbb{R}^2 is $|\det(A)|$.

<https://www.desmos.com/calculator/iksjsllpxz>

Proof sketch:

1. Rotate the parallelogram so that \mathbf{u} is on the positive x -axis.
2. Flip the parallelogram (if necessary) so that \mathbf{v} is above the x -axis.
3. Horizontally scale the parallelogram so that $\mathbf{u} = (1, 0)$.
4. Horizontally shear the parallelogram so that \mathbf{v} is on the y -axis.
5. Vertically scale the parallelogram so that $\mathbf{v} = (0, 1)$.

Volume of 3d Parallelopiped

Three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in \mathbb{R}^3 form a **parallelopiped** with vertices $\mathbf{0}, \mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{w}, \mathbf{v} + \mathbf{w}, \mathbf{u} + \mathbf{v} + \mathbf{w}$.

<https://www.geogebra.org/m/VvR6MeR5>

Volume of 3d Parallelopiped

Three vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in \mathbb{R}^3 form a **parallelopiped** with vertices $\mathbf{0}, \mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{w}, \mathbf{v} + \mathbf{w}, \mathbf{u} + \mathbf{v} + \mathbf{w}$.

<https://www.geogebra.org/m/VvR6MeR5>

Theorem

The volume of the parallelopiped \mathcal{P} in \mathbb{R}^3 is $|\det(A)|$, where A has rows \mathbf{u} , \mathbf{v} , and \mathbf{w} .

This works in higher dimensions: for an $n \times n$ matrix A , the volume of the **parallelotope** in \mathbb{R}^n corresponding to the rows of A is given by $|\det(A)|$.

Linear Transformations

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation where $\mathbf{x} \mapsto A\mathbf{x}$.

Consider the unit parallelotope \mathcal{U} with volume 1 determined by the origin and $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

What is the **volume** of $T(\mathcal{U})$?

Linear Transformations

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation where $\mathbf{x} \mapsto A\mathbf{x}$.

Consider the unit parallelotope \mathcal{U} with volume 1 determined by the origin and $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

What is the **volume** of $T(\mathcal{U})$?

$T(\mathcal{U})$ is a parallelotope determined by the columns of $AI = A$. So its volume is **$|\det A|$** .

Linear Transformations

Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation where $\mathbf{x} \mapsto A\mathbf{x}$.

Consider the unit parallelotope \mathcal{U} with volume 1 determined by the origin and $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$.

What is the **volume** of $T(\mathcal{U})$?

$T(\mathcal{U})$ is a parallelotope determined by the columns of $AI = A$. So its volume is **$|\det A|$** .



Since T is a linear transformation, the volume of **any shape** is scaled by $|\det A|$,

ie, if \mathcal{R} is a shape in \mathbb{R}^n with volume **r** , then $T(\mathcal{R})$ has volume **$|\det A|r$** .

Example: Area of an Ellipse

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, with associated matrix A .

The image of the **unit circle** centered at the origin is an **ellipse**. What is the **area** of the ellipse?





$$A = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix}.$$

Example: Area of an Ellipse

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation, with associated matrix A .

The image of the **unit circle** centered at the origin is an **ellipse**. What is the **area** of the ellipse?



$$A = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix}.$$

The area of the **unit circle** is $\pi r^2 = \pi$.

The area of the **ellipse** is $|\det A|\pi = 4\pi$.

Change of Variable in Integration

Consider the integral $\int f(x) \, dx$, and a change of variables $x = g(u)$.

Change of Variable in Integration

Consider the integral $\int f(x) dx$, and a change of variables $x = g(u)$.

$$\text{Then } \int f(x) dx = \int f(g(u)) g'(u) du.$$

Change of Variable in Integration



Consider the integral $\int f(x) dx$, and a change of variables $x = g(u)$.

$$\text{Then } \int f(x) dx = \int f(g(u)) g'(u) du.$$

Consider $\int_a^b f(x) dx = \int_4^9 \sqrt{x} dx$. Let $x = u^2$.

$$\text{Then } = \int_{\sqrt{4}}^{\sqrt{9}} \sqrt{u^2} 2u du = \int_2^3 2u^2 du = \left. \frac{2}{3} u^3 \right|_2^3 = \frac{2}{3} (3^3 - 2^3) = \frac{38}{3}.$$

Change of Variable in Integration



Polar Coordinates in 2 dimensions

Consider the integral $\iint f(x, y) \, dx \, dy$, and a change of variables $\begin{matrix} x = r \cos \theta \\ y = r \sin \theta \end{matrix}$.

<https://www.desmos.com/calculator/bfn2hgtvqp>

Polar Coordinates in 2 dimensions

Consider the integral $\iint f(x, y) \, dx \, dy$, and a change of variables $\begin{matrix} x = r \cos \theta \\ y = r \sin \theta \end{matrix}$.

<https://www.desmos.com/calculator/bfn2hgtvqp>

The correction needed describes how the area of a little rectangle $dx \times dy$ changes with respect to r and θ , and is given by the determinant of the **Jacobian matrix**:

$$\mathbf{J}(r, \theta) = \begin{bmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix}$$

Polar Coordinates in 2 dimensions

Consider the integral $\iint f(x, y) \, dx \, dy$, and a change of variables $\begin{matrix} x = r \cos \theta \\ y = r \sin \theta \end{matrix}$.

<https://www.desmos.com/calculator/bfn2hgtvqp>

The correction needed describes how the area of a little rectangle $dx \times dy$ changes with respect to r and θ , and is given by the determinant of the **Jacobian matrix**:

$$\mathbf{J}(r, \theta) = \begin{bmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix}$$

$$\begin{aligned} \text{Then } \iint f(x, y) \, dx \, dy &= \iint f(r \cos \theta, r \sin \theta) \det \mathbf{J} \, dr \, d\theta \\ &= \iint f(r \cos \theta, r \sin \theta) \, r \, dr \, d\theta. \end{aligned}$$

Spherical Coordinates in 3 dimensions

Consider the integral $\iiint f(x, y, z) \, dx \, dy \, dz$, and a change of variables $\begin{matrix} x = \rho \sin \varphi \cos \theta \\ y = \rho \sin \varphi \sin \theta \\ z = \rho \cos \varphi \end{matrix}$.

<https://www.geogebra.org/m/h9xS5ZZs>

Spherical Coordinates in 3 dimensions

Consider the integral $\iiint f(x, y, z) \, dx \, dy \, dz$, and a change of variables $\begin{matrix} x = \rho \sin \varphi \cos \theta \\ y = \rho \sin \varphi \sin \theta \\ z = \rho \cos \varphi \end{matrix}$.

<https://www.geogebra.org/m/h9xS5ZZs>

$$\mathbf{J}(r, \theta) = \begin{bmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \varphi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \varphi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \varphi} & \frac{\partial z}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \sin \varphi \cos \theta & \rho \cos \varphi \cos \theta & -\rho \sin \varphi \sin \theta \\ \sin \varphi \sin \theta & \rho \cos \varphi \sin \theta & \rho \sin \varphi \cos \theta \\ \cos \varphi & -\rho \sin \varphi & 0 \end{bmatrix}$$

Spherical Coordinates in 3 dimensions

Consider the integral $\iiint f(x, y, z) \, dx \, dy \, dz$, and a change of variables $\begin{matrix} x = \rho \sin \varphi \cos \theta \\ y = \rho \sin \varphi \sin \theta \\ z = \rho \cos \varphi \end{matrix}$.

<https://www.geogebra.org/m/h9xS5ZZs>

$$\mathbf{J}(\rho, \varphi, \theta) = \begin{bmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \varphi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \varphi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \varphi} & \frac{\partial z}{\partial \theta} \end{bmatrix} = \begin{bmatrix} \sin \varphi \cos \theta & \rho \cos \varphi \cos \theta & -\rho \sin \varphi \sin \theta \\ \sin \varphi \sin \theta & \rho \cos \varphi \sin \theta & \rho \sin \varphi \cos \theta \\ \cos \varphi & -\rho \sin \varphi & 0 \end{bmatrix}$$

$$\begin{aligned} \text{Then } \iiint f(x, y, z) \, dx \, dy \, dz &= \iiint f(x, y, z) \det \mathbf{J} \, d\rho \, d\varphi \, d\theta \\ &= \iiint f(\rho \sin \varphi \cos \theta, \rho \sin \varphi \sin \theta, \rho \cos \varphi) \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta. \end{aligned}$$