

Inner Product Spaces

Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

The Dot Product

Let \mathbf{u} and \mathbf{v} denote two vectors in \mathbb{R}^n . The dot product of \mathbf{u} and \mathbf{v} is

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v} = \begin{bmatrix} u_1 & u_2 & \dots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n = \sum_{i=1}^n u_i v_i.$$

- ▶ The dot product is used to find **lengths** of vectors and **distances** between vectors in \mathbb{R}^n .
- ▶ The dot product is used to determine if vectors in \mathbb{R}^n are **orthogonal**.
- ▶ Let's extend this concept more generally to any vector space V .

Inner Product

Definition

An **inner product** on a vector space V is a function that associates to each pair of vectors \mathbf{u} and \mathbf{v} in V a **real number** that we denote $\langle \mathbf{u}, \mathbf{v} \rangle$. An inner product satisfies the following properties for every \mathbf{u} , \mathbf{v} , and \mathbf{w} in V and scalar c :

1. $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$.
2. $\langle \mathbf{u} + \mathbf{w}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle$.
3. $\langle c\mathbf{u}, \mathbf{v} \rangle = c\langle \mathbf{u}, \mathbf{v} \rangle$.
4. $\langle \mathbf{u}, \mathbf{u} \rangle \geq 0$, and $\langle \mathbf{u}, \mathbf{u} \rangle = 0$ if and only if $\mathbf{u} = \mathbf{0}$.

A vector space with an inner product is called an **inner product space**.

Example. The vector space \mathbb{R}^n with the usual dot product is an inner product space.

Verify Properties Example 1

Let $V = \mathbb{R}^2$ and consider the function given by $\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2$. Show V with this function is an inner product space.

Verify Properties Example 1

Let $V = \mathbb{R}^2$ and consider the function given by $\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2$. Show V with this function is an inner product space.

We verify the properties of an **inner product**:

1. $\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2 = 2v_1u_1 + 3v_2u_2 = \langle \mathbf{v}, \mathbf{u} \rangle$.
2. $\langle \mathbf{u} + \mathbf{w}, \mathbf{v} \rangle = 2(u_1 + w_1)v_1 + 3(u_2 + w_2)v_2 = 2u_1v_1 + 2w_1v_1 + 3u_2v_2 + 3w_2v_2 = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle$.
3. $\langle c\mathbf{u}, \mathbf{v} \rangle = 2(cu_1)v_1 + 3(cu_2)v_2 = c(2u_1v_1 + 3u_2v_2) = c\langle \mathbf{u}, \mathbf{v} \rangle$.
4. $\langle \mathbf{u}, \mathbf{u} \rangle = 2u_1^2 + 3u_2^2 \geq 0$, and equality if and only if $u_1 = u_2 = 0$.

Verify Properties Example 2

Let $V = \mathbb{R}^2$ and consider the function given by $\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_2 + u_2 v_1$. Determine whether V with this function is an inner product space.

Verify Properties Example 2

Let $V = \mathbb{R}^2$ and consider the function given by $\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_2 + u_2 v_1$. Determine whether V with this function is an inner product space.

1. $\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_2 + u_2 v_1 = v_1 u_2 + v_2 u_1 = \langle \mathbf{v}, \mathbf{u} \rangle$.
2. $\langle \mathbf{u} + \mathbf{w}, \mathbf{v} \rangle = (u_1 + w_1)v_2 + (u_2 + w_2)v_1 = (u_1 v_2 + u_2 v_1) + (w_1 v_2 + w_2 v_1) = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle$.
3. $\langle c\mathbf{u}, \mathbf{v} \rangle = (cu_1)v_2 + (cu_2)v_1 = c(u_1 v_2 + u_2 v_1) = c\langle \mathbf{u}, \mathbf{v} \rangle$.

Verify Properties Example 2

Let $V = \mathbb{R}^2$ and consider the function given by $\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_2 + u_2 v_1$. Determine whether V with this function is an inner product space.

1. $\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_2 + u_2 v_1 = v_1 u_2 + v_2 u_1 = \langle \mathbf{v}, \mathbf{u} \rangle$.
2. $\langle \mathbf{u} + \mathbf{w}, \mathbf{v} \rangle = (u_1 + w_1)v_2 + (u_2 + w_2)v_1 = (u_1 v_2 + u_2 v_1) + (w_1 v_2 + w_2 v_1) = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle$.
3. $\langle c\mathbf{u}, \mathbf{v} \rangle = (cu_1)v_2 + (cu_2)v_1 = c(u_1 v_2 + u_2 v_1) = c\langle \mathbf{u}, \mathbf{v} \rangle$.
4. Consider the vector $\mathbf{u} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. We have

$$\langle \mathbf{u}, \mathbf{u} \rangle = (1)(0) + (0)(1) = 0 \quad \text{but} \quad \mathbf{u} \neq \mathbf{0}.$$

The last property is **not** satisfied.

\mathbb{R}^2 with the operation defined above is **NOT** an inner product space.

Inner Products on Polynomial Vector Spaces

Recall \mathbb{P}_2 denotes the vector space of all polynomials of degree at most 2. For any two vectors $p(t)$ and $q(t)$ in \mathbb{P}_2 , let

$$\langle p(t), q(t) \rangle = p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Determine whether \mathbb{P}_2 with this function is an **inner product** space.

Inner Products on Polynomial Vector Spaces

Recall \mathbb{P}_2 denotes the vector space of all polynomials of degree at most 2. For any two vectors $p(t)$ and $q(t)$ in \mathbb{P}_2 , let

$$\langle p(t), q(t) \rangle = p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Determine whether \mathbb{P}_2 with this function is an **inner product** space.

1. $\langle p(t), q(t) \rangle = p(0)q(0) + p(1)q(1) + p(2)q(2) = q(0)p(0) + q(1)p(1) + q(2)p(2) = \langle q(t), p(t) \rangle.$
2.
$$\begin{aligned}\langle p(t) + r(t), q(t) \rangle &= (p(0) + r(0))q(0) + (p(1) + r(1))q(1) + (p(2) + r(2))q(2) \\ &= (p(0)q(0) + p(1)q(1) + p(2)q(2)) + (r(0)q(0) + r(1)q(1) + r(2)q(2)) \\ &= \langle p(t), q(t) \rangle + \langle r(t), q(t) \rangle.\end{aligned}$$

Inner Products on Polynomial Vector Spaces

Recall \mathbb{P}_2 denotes the vector space of all polynomials of degree at most 2. For any two vectors $p(t)$ and $q(t)$ in \mathbb{P}_2 , let

$$\langle p(t), q(t) \rangle = p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Determine whether \mathbb{P}_2 with this function is an **inner product** space.

$$\begin{aligned} 3. \quad \langle cp(t), q(t) \rangle &= (cp(0))q(0) + (cp(1))q(1) + (cp(2))q(2) \\ &= c(p(0)q(0) + p(1)q(1) + p(2)q(2)) \\ &= c\langle p(t), q(t) \rangle \end{aligned}$$

$$4. \quad \langle p(t), p(t) \rangle = p(0)p(0) + p(1)p(1) + p(2)p(2) = (p(0))^2 + (p(1))^2 + (p(2))^2 \geq 0.$$

Inner Products on Polynomial Vector Spaces

Recall \mathbb{P}_2 denotes the vector space of all polynomials of degree at most 2. For any two vectors $p(t)$ and $q(t)$ in \mathbb{P}_2 , let

$$\langle p(t), q(t) \rangle = p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Determine whether \mathbb{P}_2 with this function is an **inner product** space.

$$\begin{aligned} 3. \quad \langle cp(t), q(t) \rangle &= (cp(0))q(0) + (cp(1))q(1) + (cp(2))q(2) \\ &= c(p(0)q(0) + p(1)q(1) + p(2)q(2)) \\ &= c\langle p(t), q(t) \rangle \end{aligned}$$

$$4. \quad \langle p(t), p(t) \rangle = p(0)p(0) + p(1)p(1) + p(2)p(2) = (p(0))^2 + (p(1))^2 + (p(2))^2 \geq 0.$$

We have equality when $p(0) = p(1) = p(2) = 0$. If a polynomial with at most degree two has three distinct zeros, then the polynomial must be equal to zero for all t .

This means $\langle p(t), p(t) \rangle = 0$ if and only if $p(t) = 0 + 0t + 0t^2 = \mathbf{0}$.

Lengths, Distances, and Orthogonality

Definition

Let V be an inner product space.

- ▶ We define the **length** (or **norm**) of a vector \mathbf{v} to be $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$.
- ▶ A **unit vector** is a vector whose length is 1.
- ▶ The **distance between vectors** \mathbf{u} and \mathbf{v} is $\|\mathbf{u} - \mathbf{v}\|$.
- ▶ Vectors \mathbf{u} and \mathbf{v} are **orthogonal** if and only if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

Our vector space has been equipped with a **geometric** structure (we have defined distances and angles in our space).

Nearly everything discussed about dot products and orthogonality for \mathbb{R}^n carries over to inner product spaces more generally (including **orthonormal bases** and the **Gram–Schmidt process**).

Polynomial Vector Space Example

Let $V = \mathbb{P}_4$ with the inner product

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Compute lengths and orthogonality of vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$ in \mathbb{P}_4 .

Polynomial Vector Space Example

Let $V = \mathbb{P}_4$ with the inner product

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

Compute lengths and orthogonality of vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$ in \mathbb{P}_4 .

We create column vectors by evaluating the polynomials at the values $t = -2, -1, 0, 1, 2$:

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\|p_1(t)\| = \sqrt{\langle 1, 1 \rangle} = \sqrt{1^2 + 1^2 + 1^2 + 1^2 + 1^2} = \sqrt{5}$$

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\|p_1(t)\| = \sqrt{\langle 1, 1 \rangle} = \sqrt{1^2 + 1^2 + 1^2 + 1^2 + 1^2} = \sqrt{5}$$

2. Are the vectors $p_1(t) = 1$ and $p_2(t) = t$ orthogonal?

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\|p_1(t)\| = \sqrt{\langle 1, 1 \rangle} = \sqrt{1^2 + 1^2 + 1^2 + 1^2 + 1^2} = \sqrt{5}$$

2. Are the vectors $p_1(t) = 1$ and $p_2(t) = t$ orthogonal?

$$\langle 1, t \rangle = (1)(-2) + (1)(-1) + (1)(0) + (1)(1) + (1)(2) = 0 \quad \Rightarrow \text{orthogonal}$$

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\|p_1(t)\| = \sqrt{\langle 1, 1 \rangle} = \sqrt{1^2 + 1^2 + 1^2 + 1^2 + 1^2} = \sqrt{5}$$

2. Are the vectors $p_1(t) = 1$ and $p_2(t) = t$ orthogonal?

$$\langle 1, t \rangle = (1)(-2) + (1)(-1) + (1)(0) + (1)(1) + (1)(2) = 0 \quad \Rightarrow \text{orthogonal}$$

3. Are the vectors $p_1(t) = 1$ and $p_3(t) = t^2$ orthogonal?

$$\langle p(t), q(t) \rangle = p(-2)q(-2) + p(-1)q(-1) + p(0)q(0) + p(1)q(1) + p(2)q(2).$$

$$p_1(t) = 1 \text{ has } \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad p_2(t) = t \text{ has } \begin{bmatrix} -2 \\ -1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \quad p_3(t) = t^2 \text{ has } \begin{bmatrix} 4 \\ 1 \\ 0 \\ 1 \\ 4 \end{bmatrix}$$

1. Compute the length of the vector $p_1(t) = 1$.

$$\|p_1(t)\| = \sqrt{\langle 1, 1 \rangle} = \sqrt{1^2 + 1^2 + 1^2 + 1^2 + 1^2} = \sqrt{5}$$

2. Are the vectors $p_1(t) = 1$ and $p_2(t) = t$ orthogonal?

$$\langle 1, t \rangle = (1)(-2) + (1)(-1) + (1)(0) + (1)(1) + (1)(2) = 0 \quad \Rightarrow \text{orthogonal}$$

3. Are the vectors $p_1(t) = 1$ and $p_3(t) = t^2$ orthogonal?

$$\langle 1, t^2 \rangle = (1)(4) + (1)(1) + (1)(0) + (1)(1) + (1)(4) = 10 \quad \Rightarrow \text{not orthogonal}$$

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

First we note the following inner products (using the inner product on \mathbb{P}_4):

$$\langle 1, 1 \rangle = 5, \quad \langle t, t \rangle = 10, \quad \langle t^2, t^2 \rangle = 34, \quad \langle 1, t \rangle = 0, \quad \langle 1, t^2 \rangle = 10, \quad \langle t, t^2 \rangle = 0.$$

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

First we note the following inner products (using the inner product on \mathbb{P}_4):

$$\langle 1, 1 \rangle = 5, \quad \langle t, t \rangle = 10, \quad \langle t^2, t^2 \rangle = 34, \quad \langle 1, t \rangle = 0, \quad \langle 1, t^2 \rangle = 10, \quad \langle t, t^2 \rangle = 0.$$

► Our first vector is $\mathbf{v}_1 = p_1(t) = 1$.

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

First we note the following inner products (using the inner product on \mathbb{P}_4):

$$\langle 1, 1 \rangle = 5, \quad \langle t, t \rangle = 10, \quad \langle t^2, t^2 \rangle = 34, \quad \langle 1, t \rangle = 0, \quad \langle 1, t^2 \rangle = 10, \quad \langle t, t^2 \rangle = 0.$$

- ▶ Our first vector is $\mathbf{v}_1 = p_1(t) = 1$.
- ▶ Next $\mathbf{v}_2 = p_2(t) - \text{proj}_{\mathbf{v}_1} p_2(t) = t - 0 = t$.

$$\text{proj}_{\mathbf{v}_1} p_2(t) = c_1 \cdot 1 = 0 \quad \text{where} \quad c_1 = \frac{\langle t, 1 \rangle}{\langle 1, 1 \rangle} = 0.$$

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

First we note the following inner products (using the inner product on \mathbb{P}_4):

$$\langle 1, 1 \rangle = 5, \quad \langle t, t \rangle = 10, \quad \langle t^2, t^2 \rangle = 34, \quad \langle 1, t \rangle = 0, \quad \langle 1, t^2 \rangle = 10, \quad \langle t, t^2 \rangle = 0.$$

- ▶ Our first vector is $\mathbf{v}_1 = p_1(t) = 1$.
- ▶ Next $\mathbf{v}_2 = p_2(t) - \text{proj}_{\mathbf{v}_1} p_2(t) = t - 0 = t$.

$$\text{proj}_{\mathbf{v}_1} p_2(t) = c_1 \cdot 1 = 0 \quad \text{where} \quad c_1 = \frac{\langle t, 1 \rangle}{\langle 1, 1 \rangle} = 0.$$

- ▶ The third (and final) vector $\mathbf{v}_3 = p_3(t) - \text{proj}_{\{\mathbf{v}_1, \mathbf{v}_2\}} p_3(t) = t^2 - 2$.

$$\text{proj}_{\{1, t\}} p_3(t) = c_1 \cdot 1 + c_2 \cdot t = (2)(1) - (0)(t) = 2 \quad \text{where} \quad c_1 = \frac{\langle t^2, 1 \rangle}{\langle 1, 1 \rangle} = \frac{10}{5}, \quad c_2 = \frac{\langle t^2, t \rangle}{\langle t, t \rangle} = 0.$$

Producing an Orthogonal Basis for a Polynomial Vector Space

Apply the **Gram–Schmidt process** to produce an **orthogonal basis** for the subspace V of \mathbb{P}_4 spanned by the vectors $p_1(t) = 1$, $p_2(t) = t$, and $p_3(t) = t^2$.

First we note the following inner products (using the inner product on \mathbb{P}_4):

$$\langle 1, 1 \rangle = 5, \quad \langle t, t \rangle = 10, \quad \langle t^2, t^2 \rangle = 34, \quad \langle 1, t \rangle = 0, \quad \langle 1, t^2 \rangle = 10, \quad \langle t, t^2 \rangle = 0.$$

- ▶ Our first vector is $\mathbf{v}_1 = p_1(t) = 1$.
- ▶ Next $\mathbf{v}_2 = p_2(t) - \text{proj}_{\mathbf{v}_1} p_2(t) = t - 0 = t$.

$$\text{proj}_{\mathbf{v}_1} p_2(t) = c_1 \cdot 1 = 0 \quad \text{where} \quad c_1 = \frac{\langle t, 1 \rangle}{\langle 1, 1 \rangle} = 0.$$

- ▶ The third (and final) vector $\mathbf{v}_3 = p_3(t) - \text{proj}_{\{\mathbf{v}_1, \mathbf{v}_2\}} p_3(t) = t^2 - 2$.

$$\text{proj}_{\{1, t\}} p_3(t) = c_1 \cdot 1 + c_2 \cdot t = (2)(1) - (0)(t) = 2 \quad \text{where} \quad c_1 = \frac{\langle t^2, 1 \rangle}{\langle 1, 1 \rangle} = \frac{10}{5}, \quad c_2 = \frac{\langle t^2, t \rangle}{\langle t, t \rangle} = 0.$$

Our **orthogonal basis** is $\{1, t, t^2 - 2\}$. Note this depends on the **inner product of \mathbb{P}_4** .

An Inner Product on Functional Vector Spaces

Let $\mathcal{C}[a, b]$ denote the set of all **continuous functions** on the closed interval $a \leq t \leq b$.

- ▶ Using the typical addition of functions and scalar multiplication, we can verify that $\mathcal{C}[a, b]$ is a vector space.
- ▶ For two functions f and g in $\mathcal{C}[a, b]$, we define the following inner product:

$$\langle f, g \rangle = \int_a^b f(t)g(t) dt$$

Show $\mathcal{C}[a, b]$ with the operation defined above is an **inner product space**.

An Inner Product on Functional Vector Spaces

Let $\mathcal{C}[a, b]$ denote the set of all **continuous functions** on the closed interval $a \leq t \leq b$.

- ▶ Using the typical addition of functions and scalar multiplication, we can verify that $\mathcal{C}[a, b]$ is a vector space.
- ▶ For two functions f and g in $\mathcal{C}[a, b]$, we define the following inner product:

$$\langle f, g \rangle = \int_a^b f(t)g(t) dt$$

Show $\mathcal{C}[a, b]$ with the operation defined above is an **inner product space**.

1. $\langle f, g \rangle = \int_a^b f(t)g(t) dt = \int_a^b g(t)f(t) dt = \langle g, f \rangle.$
2. $\langle \mathbf{f} + \mathbf{h}, g \rangle = \int_a^b ((\mathbf{f}(t) + \mathbf{h}(t))g(t)) dt = \int_a^b \mathbf{f}(t)g(t) dt + \int_a^b \mathbf{h}(t)g(t) dt = \langle \mathbf{f}, g \rangle + \langle \mathbf{h}, g \rangle.$
3. $\langle \mathbf{c}f, g \rangle = \int_a^b (\mathbf{c}f(t))g(t) dt = \mathbf{c} \int_a^b f(t)g(t) dt = \mathbf{c} \langle f, g \rangle.$
4. $\langle f, f \rangle = \int_a^b (f(t))^2 dt \geq 0$ and equality if and only if $f(t) = 0$.

Computing Inner Products of Continuous Functions

Consider the inner product space $\mathcal{C}[0, 1]$ with the usual inner product $\langle f, g \rangle = \int_a^b f(t)g(t) dt$.

Compute $\langle 1 + t^2, 4t \rangle$.

Computing Inner Products of Continuous Functions

Consider the inner product space $\mathcal{C}[0, 1]$ with the usual inner product $\langle f, g \rangle = \int_a^b f(t)g(t) dt$. Compute $\langle 1 + t^2, 4t \rangle$.

$$\begin{aligned}\langle 1 + t^2, 4t \rangle &= \int_0^1 \left((1 + t^2)4t \right) dt \\ &= \int_0^1 (4t + 4t^3) dt \\ &= (2t^2 + t^4) \Big|_0^1 \\ &= 3.\end{aligned}$$

Thus, $1 + t^2$ and $4t$ are **not orthogonal** in $\mathcal{C}[0, 1]$.

Fourier Analysis

A **basis** for $\mathcal{C}[0, 2\pi]$ is $\{1\} \cup \{\cos(mt) : m = 1, 2, \dots\} \cup \{\sin(mt) : m = 1, 2, \dots\}$.

Fourier Analysis

A **basis** for $\mathcal{C}[0, 2\pi]$ is $\{1\} \cup \{\cos(mt) : m = 1, 2, \dots\} \cup \{\sin(mt) : m = 1, 2, \dots\}$.

In fact, this is an **orthogonal basis**. For example, where $m \neq n$ are positive integers,

$$\begin{aligned}\langle \cos(mt), \cos(nt) \rangle &= \int_0^{2\pi} \cos(mt) \cos(nt) \, dt \\ &= \frac{1}{2} \int_0^{2\pi} [\cos(mt + nt) + \cos(mt - nt)] \, dt \\ &= \frac{1}{2} \left[\frac{\sin(mt + nt)}{m + n} + \frac{\sin(mt - nt)}{m - n} \right] \Big|_0^{2\pi} = 0.\end{aligned}$$

Fourier Analysis

A **basis** for $\mathcal{C}[0, 2\pi]$ is $\{1\} \cup \{\cos(mt) : m = 1, 2, \dots\} \cup \{\sin(mt) : m = 1, 2, \dots\}$.

In fact, this is an **orthogonal basis**. Any $f \in \mathcal{C}[0, 2\pi]$ can be represented by a **Fourier series**

$$f(t) = \frac{a_0}{2} + \sum_{m=1}^{\infty} a_m \cos(mt) + \sum_{m=1}^{\infty} b_m \sin(mt),$$

where for $m \geq 1$,

$$a_m = \frac{\langle f, \cos(mt) \rangle}{\langle \cos(mt), \cos(mt) \rangle} = \frac{1}{\pi} \int_0^{2\pi} f(t) \cos(mt) dt, \quad \text{etc.}$$

Fourier Analysis

A **basis** for $\mathcal{C}[0, 2\pi]$ is $\{1\} \cup \{\cos(mt) : m = 1, 2, \dots\} \cup \{\sin(mt) : m = 1, 2, \dots\}$.

In fact, this is an **orthogonal basis**. Any $f \in \mathcal{C}[0, 2\pi]$ can be represented by a **Fourier series**

$$f(t) = \frac{a_0}{2} + \sum_{m=1}^{\infty} a_m \cos(mt) + \sum_{m=1}^{\infty} b_m \sin(mt),$$

where for $m \geq 1$,

$$a_m = \frac{\langle f, \cos(mt) \rangle}{\langle \cos(mt), \cos(mt) \rangle} = \frac{1}{\pi} \int_0^{2\pi} f(t) \cos(mt) dt, \quad \text{etc.}$$

Let W be spanned by $\{1\} \cup \{\cos(mt) : m = 1, \dots, n\} \cup \{\sin(mt) : m = 1, \dots, n\}$.

What is the **best approximation** of f by elements of W ?

Python examples in Jupyter Notebook.