

# Orthogonal Diagonalization and Symmetric Matrices

## Linear Algebra

These materials were created by Adam Spiegler, Stephen Hartke, and others, and are licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

This work was initially funded by an Institutional OER Grant from the Colorado Department of Higher Education (CDHE). For similar OER materials in other courses funded by this project in the Department of Mathematical and Statistical Sciences at the University of Colorado Denver, visit <https://github.com/CU-Denver-MathStats-OER>

# Diagonalizing with an Orthonormal Basis

**Thm.** An  $n \times n$  matrix  $A$  is **diagonalizable** if and only if there exist  $n$  linearly independent eigenvectors.

# Diagonalizing with an Orthonormal Basis

**Thm.** An  $n \times n$  matrix  $A$  is **diagonalizable** if and only if  
there exist  $n$  linearly independent eigenvectors.  
 $\Leftrightarrow$  there is a **basis** for  $\mathbb{R}^n$  consisting of eigenvectors.

**Ques.** When does there exist an **orthonormal basis** for  $\mathbb{R}^n$  consisting of eigenvectors?

# Symmetric Matrix

## Definition

A **symmetric matrix** is a square matrix  $A$  such that  $A^T = A$ .

- ▶ Entries on the main diagonal can be anything.
- ▶ Entries above and below the main diagonal come in mirrored pairs,  $a_{ij} = a_{ji}$ .

Here are some examples of symmetric matrices:

$$\begin{bmatrix} 1 & 2 \\ 2 & -4 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & -2 & 7 \\ -2 & 12 & 5 \\ 7 & 5 & -17 \end{bmatrix}.$$

Here are some examples of matrices that are not symmetric:

$$\begin{bmatrix} 1 & 2 \\ -2 & -4 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}.$$

## Review: Diagonalizing a Matrix

Diagonalize the matrix  $A = \begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix}$ .

1. Find the eigenvalues of  $A$  by solving  $\det(A - \lambda I) = 0$ . We have  $\lambda = 8, 6, 3$ .

## Review: Diagonalizing a Matrix

Diagonalize the matrix  $A = \begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix}$ .

1. Find the eigenvalues of  $A$  by solving  $\det(A - \lambda I) = 0$ . We have  $\lambda = 8, 6, 3$ .
2. If possible, find  $n$  linearly independent eigenvectors.

We have  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

## Review: Diagonalizing a Matrix

Diagonalize the matrix  $A = \begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix}$ .

1. Find the eigenvalues of  $A$  by solving  $\det(A - \lambda I) = 0$ . We have  $\lambda = 8, 6, 3$ .
2. If possible, find  $n$  linearly independent eigenvectors.

We have  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

3. Then  $A = PDP^{-1}$ . We have  $P = \begin{bmatrix} -1 & -1 & 1 \\ 1 & -1 & 1 \\ 0 & 2 & 1 \end{bmatrix}$  and  $D = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ .

## Choosing An Orthonormal Basis

We have  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.



## Choosing An Orthonormal Basis

We have  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

We can instead construct column vectors for  $P$  that are **orthonormal**:

We have  $\mathbf{u}_1 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1/\sqrt{6} \\ -1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix}$ , and  $\mathbf{u}_3 = \begin{bmatrix} 1/\sqrt{3} \\ 1/\sqrt{3} \\ 1/\sqrt{3} \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

## Choosing An Orthonormal Basis

We have  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

We can instead construct column vectors for  $P$  that are **orthonormal**:

We have  $\mathbf{u}_1 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1/\sqrt{6} \\ -1/\sqrt{6} \\ 2/\sqrt{6} \end{bmatrix}$ , and  $\mathbf{u}_3 = \begin{bmatrix} 1/\sqrt{3} \\ 1/\sqrt{3} \\ 1/\sqrt{3} \end{bmatrix}$  corresponding to  $\lambda_1 = 8$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 3$ , respectively.

Now we have  $P = \begin{bmatrix} -1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 0 & 2/\sqrt{6} & 1/\sqrt{3} \end{bmatrix}$  and  $D = \begin{bmatrix} 8 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ .

# Orthogonal Matrices

## Definition

A matrix  $P$  is called **orthogonal** if it is a square matrix with **orthonormal columns**. If  $P$  is an orthogonal matrix, we have shown that  $P^T P = I$  and  $PP^T = I$ , therefore  $P^{-1} = P^T$ .

$$A = PDP^{-1} = PDP^T$$

$$\begin{bmatrix} 6 & -2 & -1 \\ -2 & 6 & -1 \\ -1 & -1 & 5 \end{bmatrix} = \begin{bmatrix} -1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 1/\sqrt{2} & -1/\sqrt{6} & 1/\sqrt{3} \\ 0 & 2/\sqrt{6} & 1/\sqrt{3} \end{bmatrix} \begin{bmatrix} 8 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} & 0 \\ -1/\sqrt{6} & -1/\sqrt{6} & 2/\sqrt{6} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \end{bmatrix}$$

# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

## Proof.

Let  $\mathbf{v}_1$  and  $\mathbf{v}_2$  be eigenvectors that correspond to different eigenvalues  $\lambda_1$  and  $\lambda_2$ , respectively. We need to show that  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .

# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

## Proof.

Let  $\mathbf{v}_1$  and  $\mathbf{v}_2$  be eigenvectors that correspond to different eigenvalues  $\lambda_1$  and  $\lambda_2$ , respectively. We need to show that  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .

$$\lambda_1 \mathbf{v}_1 \cdot \mathbf{v}_2 = (A\mathbf{v}_1) \cdot \mathbf{v}_2 = (A\mathbf{v}_1)^T \mathbf{v}_2 \quad \text{by def of eigenvector}$$

# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

## Proof.

Let  $\mathbf{v}_1$  and  $\mathbf{v}_2$  be eigenvectors that correspond to different eigenvalues  $\lambda_1$  and  $\lambda_2$ , respectively. We need to show that  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .

$$\begin{aligned}\lambda_1 \mathbf{v}_1 \cdot \mathbf{v}_2 &= (A\mathbf{v}_1) \cdot \mathbf{v}_2 = (A\mathbf{v}_1)^T \mathbf{v}_2 && \text{by def of eigenvector} \\ &= \mathbf{v}_1^T A^T \mathbf{v}_2 = \mathbf{v}_1^T A \mathbf{v}_2 && \text{since } A \text{ is symmetric}\end{aligned}$$

# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

## Proof.

Let  $\mathbf{v}_1$  and  $\mathbf{v}_2$  be eigenvectors that correspond to different eigenvalues  $\lambda_1$  and  $\lambda_2$ , respectively. We need to show that  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .

$$\begin{aligned}\lambda_1 \mathbf{v}_1 \cdot \mathbf{v}_2 &= (A\mathbf{v}_1) \cdot \mathbf{v}_2 = (A\mathbf{v}_1)^T \mathbf{v}_2 && \text{by def of eigenvector} \\ &= \mathbf{v}_1^T A^T \mathbf{v}_2 = \mathbf{v}_1^T A \mathbf{v}_2 && \text{since } A \text{ is symmetric} \\ &= \mathbf{v}_1^T \lambda_2 \mathbf{v}_2 = \lambda_2 \mathbf{v}_1 \cdot \mathbf{v}_2. && \text{by def of eigenvector}\end{aligned}$$



# Orthogonality of Eigenvectors

## Theorem

If  $A$  is **symmetric**, then any two eigenvectors from different eigenspaces are **orthogonal**.

## Proof.

Let  $\mathbf{v}_1$  and  $\mathbf{v}_2$  be eigenvectors that correspond to different eigenvalues  $\lambda_1$  and  $\lambda_2$ , respectively. We need to show that  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ .

$$\begin{aligned}\lambda_1 \mathbf{v}_1 \cdot \mathbf{v}_2 &= (A\mathbf{v}_1) \cdot \mathbf{v}_2 = (A\mathbf{v}_1)^T \mathbf{v}_2 && \text{by def of eigenvector} \\ &= \mathbf{v}_1^T A^T \mathbf{v}_2 = \mathbf{v}_1^T A \mathbf{v}_2 && \text{since } A \text{ is symmetric} \\ &= \mathbf{v}_1^T \lambda_2 \mathbf{v}_2 = \lambda_2 \mathbf{v}_1 \cdot \mathbf{v}_2. && \text{by def of eigenvector}\end{aligned}$$

Therefore,  $(\lambda_1 - \lambda_2)\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$ , which implies  $\mathbf{v}_1 \cdot \mathbf{v}_2 = 0$  since we assumed  $\lambda_1$  and  $\lambda_2$  are not equal. □

# Orthogonally Diagonalizable

## Definition

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if there are an **orthogonal** matrix  $P$  and a **diagonal** matrix  $D$  such that  $A = PDP^T$ .

# Orthogonally Diagonalizable

## Definition

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if there are an **orthogonal** matrix  $P$  and a **diagonal** matrix  $D$  such that  $A = PDP^T$ .

## Theorem (The Spectral Theorem)

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if and only if  $A$  is a **symmetric** matrix.

# Orthogonally Diagonalizable

## Definition

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if there are an **orthogonal** matrix  $P$  and a **diagonal** matrix  $D$  such that  $A = PDP^T$ .

## Theorem (The Spectral Theorem)

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if and only if  $A$  is a **symmetric** matrix.

**Proof.**

$(\Rightarrow)$  Assume that  $A$  is orthogonally diagonalizable. Then we know  $A = PDP^T$ .

# Orthogonally Diagonalizable

## Definition

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if there are an **orthogonal** matrix  $P$  and a **diagonal** matrix  $D$  such that  $A = PDP^T$ .

## Theorem (The Spectral Theorem)

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if and only if  $A$  is a **symmetric** matrix.

## Proof.

( $\Rightarrow$ ) Assume that  $A$  is orthogonally diagonalizable. Then we know  $A = PDP^T$ . Taking the transpose of both sides we have

$$A^T = (PDP^T)^T = (P^T)^T D^T P^T = PDP^T = A.$$

Since  $A^T = A$ , we see that  $A$  is a symmetric matrix.

# Orthogonally Diagonalizable

## Definition

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if there are an **orthogonal** matrix  $P$  and a **diagonal** matrix  $D$  such that  $A = PDP^T$ .

## Theorem (The Spectral Theorem)

An  $n \times n$  matrix  $A$  is **orthogonally diagonalizable** if and only if  $A$  is a **symmetric** matrix.

## Proof.

( $\Rightarrow$ ) Assume that  $A$  is orthogonally diagonalizable. Then we know  $A = PDP^T$ . Taking the transpose of both sides we have

$$A^T = (PDP^T)^T = (P^T)^T D^T P^T = PDP^T = A.$$

Since  $A^T = A$ , we see that  $A$  is a symmetric matrix.

( $\Leftarrow$ ) Converse is **difficult**! Hard part:  $\dim$  of each eigenspace equals the algebraic multiplicity.



## Example

Orthogonally diagonalize the matrix  $A = \begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix}$ .

## Example

Orthogonally diagonalize the matrix  $A = \begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix}$ .

1. First we find the eigenvalues:  $\lambda_1 = -3$  (with multiplicity 2) and  $\lambda_2 = 15$ .



## Example

Orthogonally diagonalize the matrix  $A = \begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix}$ .

1. First we find the eigenvalues:  $\lambda_1 = -3$  (with multiplicity 2) and  $\lambda_2 = 15$ .
2. Find a basis for the eigenspace of each eigenvector.

$$\lambda_1 = -3 \text{ has } \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \right\} \text{ and } \lambda_2 = 15 \text{ has } \left\{ \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix} \right\}$$

## Example

Orthogonally diagonalize the matrix  $A = \begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix}$ .

1. First we find the eigenvalues:  $\lambda_1 = -3$  (with multiplicity 2) and  $\lambda_2 = 15$ .
2. Find a basis for the eigenspace of each eigenvector.

$$\lambda_1 = -3 \text{ has } \left\{ \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \right\} \text{ and } \lambda_2 = 15 \text{ has } \left\{ \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix} \right\}$$

3. Using the Gram–Schmidt process, find an orthogonal basis for the eigenspaces.

# Making Things Orthogonal

We have  $\mathbf{w}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{w}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$ , and  $\mathbf{w}_3 = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix}$ .

►  $\mathbf{w}_1$  and  $\mathbf{w}_2$  are NOT orthogonal. We set  $\mathbf{v}_1 = \mathbf{w}_1$ .

# Making Things Orthogonal

We have  $\mathbf{w}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{w}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$ , and  $\mathbf{w}_3 = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix}$ .

- ▶  $\mathbf{w}_1$  and  $\mathbf{w}_2$  are NOT orthogonal. We set  $\mathbf{v}_1 = \mathbf{w}_1$ .
- ▶ We find the component of  $\mathbf{w}_2$  orthogonal to  $\mathbf{v}_1$

$$\text{proj}_{\mathbf{v}_1} \mathbf{w}_2 = \frac{\mathbf{w}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = \frac{-1}{2} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/2 \\ -1/2 \\ 0 \end{bmatrix}$$

Our next vector is  $\mathbf{v}_2 = \mathbf{w}_2 - \text{proj}_{\mathbf{v}_1} \mathbf{w}_2 = \begin{bmatrix} 1/2 \\ 1/2 \\ 2 \end{bmatrix}$ .

# Making Things Orthogonal

We have  $\mathbf{w}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{w}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$ , and  $\mathbf{w}_3 = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix}$ .

- ▶  $\mathbf{w}_1$  and  $\mathbf{w}_2$  are NOT orthogonal. We set  $\mathbf{v}_1 = \mathbf{w}_1$ .
- ▶ We find the component of  $\mathbf{w}_2$  orthogonal to  $\mathbf{v}_1$

$$\text{proj}_{\mathbf{v}_1} \mathbf{w}_2 = \frac{\mathbf{w}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = \frac{-1}{2} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/2 \\ -1/2 \\ 0 \end{bmatrix}$$

Our next vector is  $\mathbf{v}_2 = \mathbf{w}_2 - \text{proj}_{\mathbf{v}_1} \mathbf{w}_2 = \begin{bmatrix} 1/2 \\ 1/2 \\ 2 \end{bmatrix}$ .

- ▶  $\mathbf{w}_3$  is orthogonal to  $\mathbf{v}_1$  and  $\mathbf{v}_2$ , so we set  $\mathbf{v}_3 = \mathbf{w}_3$ .

## Normalizing the Vectors

We now have orthogonal eigenvectors  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 1/2 \\ 1/2 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix}$ .

## Normalizing the Vectors

We now have orthogonal eigenvectors  $\mathbf{v}_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 1/2 \\ 1/2 \\ 2 \end{bmatrix}$ , and  $\mathbf{v}_3 = \begin{bmatrix} -2 \\ -2 \\ 1 \end{bmatrix}$ .

4. Normalize each column vector to find possible columns of  $P$ .

$$P = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \sqrt{2}/6 & -2/3 \\ \frac{1}{\sqrt{2}} & \sqrt{2}/6 & -2/3 \\ 0 & 2\sqrt{2}/3 & 1/3 \end{bmatrix}$$

We can check that  $A = PDP^T$ :

$$A = \begin{bmatrix} 5 & 8 & -4 \\ 8 & 5 & -4 \\ -4 & -4 & -1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{2}} & \sqrt{2}/6 & -2/3 \\ \frac{1}{\sqrt{2}} & \sqrt{2}/6 & -2/3 \\ 0 & 2\sqrt{2}/3 & 1/3 \end{bmatrix} \begin{bmatrix} -3 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & 15 \end{bmatrix} \begin{bmatrix} -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \sqrt{2}/6 & \sqrt{2}/6 & 2\sqrt{2}/3 \\ -2/3 & -2/3 & 1/3 \end{bmatrix}$$

# The Spectral Theorem

The set of eigenvalues of  $A$  is sometimes called the **spectrum** of  $A$ .

## Theorem (The Spectral Theorem)

If  $A$  is a **symmetric**  $n \times n$  matrix, then  $A$  is **orthogonally diagonalizable** (with real eigenvalues).

There exist **orthogonal**  $P = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n]$  and **diagonal**  $D = \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix}$  such that

$$A = \mathbf{P} \mathbf{D} \mathbf{P}^T = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n] \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix} \begin{bmatrix} \mathbf{u}_1^T \\ \vdots \\ \mathbf{u}_n^T \end{bmatrix}$$



# The Spectral Theorem

The set of eigenvalues of  $A$  is sometimes called the **spectrum** of  $A$ .

## Theorem (The Spectral Theorem)

If  $A$  is a **symmetric**  $n \times n$  matrix, then  $A$  is **orthogonally diagonalizable** (with real eigenvalues).

There exist **orthogonal**  $P = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n]$  and **diagonal**  $D = \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix}$  such that

$$A = P D P^T = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n] \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix} \begin{bmatrix} \mathbf{u}_1^T \\ \vdots \\ \mathbf{u}_n^T \end{bmatrix} = \lambda_1 \mathbf{u}_1 \mathbf{u}_1^T + \lambda_2 \mathbf{u}_2 \mathbf{u}_2^T + \dots + \lambda_n \mathbf{u}_n \mathbf{u}_n^T.$$

# The Spectral Theorem

The set of eigenvalues of  $A$  is sometimes called the **spectrum** of  $A$ .

## Theorem (The Spectral Theorem)

If  $A$  is a **symmetric**  $n \times n$  matrix, then  $A$  is **orthogonally diagonalizable** (with real eigenvalues).

There exist **orthogonal**  $P = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n]$  and **diagonal**  $D = \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix}$  such that

$$A = \textcolor{red}{P} \textcolor{blue}{D} \textcolor{red}{P}^T = [\mathbf{u}_1 \ \dots \ \mathbf{u}_n] \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \lambda_n \end{bmatrix} \begin{bmatrix} \mathbf{u}_1^T \\ \vdots \\ \mathbf{u}_n^T \end{bmatrix} = \lambda_1 \mathbf{u}_1 \mathbf{u}_1^T + \lambda_2 \mathbf{u}_2 \mathbf{u}_2^T + \dots + \lambda_n \mathbf{u}_n \mathbf{u}_n^T.$$

The matrices  $\mathbf{u}_i \mathbf{u}_i^T$  are  $n \times n$  matrices, rank 1, and are orthogonal projection matrices.

The product  $\mathbf{u} \mathbf{v}^T$  of  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$  is sometimes called the *outer product*, and has rank  $\leq 1$ .