

Introduction to the Laplace Transform

One of the basic problem solving techniques in mathematics is to

- transform a difficult problem into an easier one,
- solve the easier problem, and
- then use its solution to obtain a solution of the original problem.

For example, the reverse product rule (method of integrating factor) is used to transform a linear first order differential equation into an easier problem we can solve. In this chapter we study the method of **Laplace transforms**, which is one example of this technique. Like the method of integrating factors, Laplace transforms are **integral operators**. Solving by the method of Laplace transforms:

- Can be used to solve higher order linear differential equations.
- Can be applied for more complicated forcing functions.
- Requires initial conditions.

The **improper integral** of g over $[a, \infty)$ is defined as

$$\int_a^\infty g(t) dt = \lim_{N \rightarrow \infty} \int_a^N g(t) dt.$$

- We say the improper integral **converges** if the limit exists.
- Otherwise we say the improper integral **diverges**.

1. Determine whether $\int_0^\infty e^{-2t} dt$ converges or diverges.

Definition of the Laplace Transform

Let $f(t)$ be a function on $[0, \infty)$. The **Laplace transform** of f is the function F defined by

$$\mathcal{L}\{f\} = F(s) = \int_0^{\infty} e^{-st} f(t) dt.$$

- The domain of $F(s)$ is all values of s for which the integral converges.
- The functions f and F form a **transform pair**.

2. Find and state the domain of the Laplace transform $F(s) = \mathcal{L}\{f(t)\}$.

(a) $f(t) = 2, t \geq 0$

(b) $f(t) = t$

(c) $f(t) = e^{3t}$

(d) $g(t) = \cos(bt)$ where $b \neq 0$ is a constant.

$$(e) \ f(t) = \begin{cases} 5 & 0 < t < 2 \\ e^{8t} & t > 2 \end{cases}$$

Common Laplace Transforms

$f(t)$	$F(s) = \mathcal{L}\{f(t)\}$
$f(t) = 1$	$F(s) = \frac{1}{s}, s > 0$
$f(t) = e^{at}$	$F(s) = \frac{1}{s-a}, s > a$
$f(t) = t^n, n = 1, 2, \dots$	$F(s) = \frac{n!}{s^{n+1}}, s > 0$
$f(t) = \sin(bt)$	$F(s) = \frac{b}{s^2 + b^2}, s > 0$
$f(t) = \cos(bt)$	$F(s) = \frac{s}{s^2 + b^2}, s > 0$