

Uniform Convergence for Sequences of Iterates

Butler University

August 25, 2020

Iterates

Definition

Let $f : X \rightarrow X$; define the n^{th} iterate of f as

$$f^n := \overbrace{f \circ f \circ \dots \circ f}^n$$

Example:

If $f(z) = z^2$, then $f^3(z) = z^2 \circ z^2 \circ z^2 = (((z^2)^2)^2) = z^8$

Iterates

Definition

Let $f : X \rightarrow X$; define the n^{th} iterate of f as

$$f^n := \overbrace{f \circ f \circ \dots \circ f}^n$$

Example:

If $f(z) = z^2$, then $f^3(z) = z^2 \circ z^2 \circ z^2 = (((z^2)^2)^2) = z^8$

Definition

We say $x \in X$ is a fixed point of $f: X \rightarrow X$ if $f(x) = x$.

Example:

If $f(z) = z^2$ then $f(0) = 0$. Therefore 0 is a fixed point.

Old Theorems

Denjoy-Wolff Theorem

If $\varphi: \mathbb{D} \rightarrow \mathbb{D}$ is analytic but neither the identity nor a rotation, then there exists a unique point $a \in \overline{\mathbb{D}}$ so that φ^n converges to the constant function a uniformly on compact subsets of \mathbb{D} .

Old Theorems

Denjoy-Wolff Theorem

If $\varphi: \mathbb{D} \rightarrow \mathbb{D}$ is analytic but neither the identity nor a rotation, then there exists a unique point $a \in \overline{\mathbb{D}}$ so that φ^n converges to the constant function a uniformly on compact subsets of \mathbb{D} .

We call a the Denjoy-Wolff point:

- ▶ If $a \in \mathbb{D}$, then $\varphi(a) = a$, and f has no other fixed points;
- ▶ If $a \in \partial\overline{\mathbb{D}}$, then $\lim_{r \rightarrow 1^-} \varphi(ra) = a$.

New Theorems

Theorem 1 (Cowen, Ko, Thompson, Tiang 2014)

Suppose $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and continuous on the boundary, $\partial\mathbb{D}$. If the Denjoy-Wolff point $a \in \mathbb{D}$, then $\varphi^n \rightarrow a$ uniformly *on all of* \mathbb{D} if and only if there is $N > 0$ such that $\varphi^N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$.

New Theorems

Theorem 1 (Cowen, Ko, Thompson, Tiang 2014)

Suppose $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and continuous on the boundary, $\partial\mathbb{D}$. If the Denjoy-Wolff point $a \in \mathbb{D}$, then $\varphi^n \rightarrow a$ uniformly *on all of* \mathbb{D} if and only if there is $N > 0$ such that $\varphi^N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$.

Theorem 2 (Cowen, Ko, Thompson, Tiang 2014)

Suppose $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and continuous on the boundary, $\partial\mathbb{D}$, and has Denjoy-Wolff point a with $|a| = 1$ and $\varphi'(a) < 1$. If $\varphi^N(\mathbb{D}) \subseteq \mathbb{D} \cup \{a\}$ for some $N > 0$, then $\varphi^n \rightarrow a$ uniformly on all of \mathbb{D} .

New Theorems

Theorem 1 (Cowen, Ko, Thompson, Tiang 2014)

Suppose $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and continuous on the boundary, $\partial\mathbb{D}$. If the Denjoy-Wolff point $a \in \mathbb{D}$, then $\varphi^n \rightarrow a$ uniformly on all of \mathbb{D} if and only if there is $N > 0$ such that $\varphi^N(\overline{\mathbb{D}}) \subseteq \mathbb{D}$.

Theorem 2 (Cowen, Ko, Thompson, Tiang 2014)

Suppose $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and continuous on the boundary, $\partial\mathbb{D}$, and has Denjoy-Wolff point a with $|a| = 1$ and $\varphi'(a) < 1$. If $\varphi^N(\mathbb{D}) \subseteq \mathbb{D} \cup \{a\}$ for some $N > 0$, then $\varphi^n \rightarrow a$ uniformly on all of \mathbb{D} .

Definition

We say UCI (*uniformly convergent iterates*) holds for $f: X \rightarrow X$ if f^n converges uniformly on all of X to a constant a .

An Example

$\varphi(z) = \frac{1}{2}z + \frac{1}{2}$ has fixed point at 1 with $\varphi'(1) < 1$. Has UCI.

An Example

$\varphi(z) = \frac{1}{2}z + \frac{1}{2}$ has fixed point at 1 with $\varphi'(1) < 1$. Has UCI.

Another Example

$\varphi(z) = 1/(2 - z)$ has fixed point at 1 with $\varphi'(1) = 1$. Has UCI.

An Example

$\varphi(z) = \frac{1}{2}z + \frac{1}{2}$ has fixed point at 1 with $\varphi'(1) < 1$. Has UCI.

Another Example

$\varphi(z) = 1/(2 - z)$ has fixed point at 1 with $\varphi'(1) = 1$. Has UCI.

Generalized Example

If φ is a linear fractional map with DW point $|a| = 1$ and $\varphi'(a) = 1$ but not an automorphism, then UCI.

An Example

$\varphi(z) = \frac{1}{2}z + \frac{1}{2}$ has fixed point at 1 with $\varphi'(1) < 1$. Has UCI.

Another Example

$\varphi(z) = 1/(2 - z)$ has fixed point at 1 with $\varphi'(1) = 1$. Has UCI.

Generalized Example

If φ is a linear fractional map with DW point $|a| = 1$ and $\varphi'(a) = 1$ but not an automorphism, then UCI.

An Annoying Example

$\varphi(z) = \frac{(z+1)^k + (z-1)}{(z+1)^k - (z-1)}$ for $k > 1$ has $\varphi(1) = 1$ and $\varphi(-1) = -1$. NO UCI.

More New Theorems

Theorem 3 (Cowen, Ko, Thompson, Tiang 2014)

If

- ▶ $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} ,
- ▶ φ is continuous on $\partial\mathbb{D}$, and
- ▶ $\varphi^n(z) \rightarrow a$ uniformly on \mathbb{D} and $|a| = 1$,

then a is the only fixed point of φ on the closed disk.

Theorem 4 (Glickfield, Kaschner 2018)

If

- ▶ $\varphi: \overline{\mathbb{D}} \rightarrow \mathbb{D}$ is analytic on \mathbb{D} and
- ▶ $\varphi^n(z) \rightarrow a$ uniformly on \mathbb{D} and $|a| = 1$,

then a is the only fixed point of φ on the closed unit disk.

Problem(s)

When does UCI hold?

Problem(s)

When does UCI hold?
When is the fixed point unique?

Problem(s)

When does UCI hold?
When is the fixed point unique?

What about $\varphi(z) = \frac{1}{2}z^2 + \frac{1}{2}$?

Problem(s)

When does UCI hold?
When is the fixed point unique?

What about $\varphi(z) = \frac{1}{2}z^2 + \frac{1}{2}$?

Let's explore different domains:

- ▶ \mathbb{D}^2 (bi-disk)
- ▶ $\{\vec{z} \in \mathbb{C}^2: \|\vec{z}\| < 1\}$ (unit ball)

Problem(s)

When does UCI hold?
When is the fixed point unique?

What about $\varphi(z) = \frac{1}{2}z^2 + \frac{1}{2}$?

Let's explore different domains:

- ▶ \mathbb{D}^2 (bi-disk)
- ▶ $\{\vec{z} \in \mathbb{C}^2: \|\vec{z}\| < 1\}$ (unit ball)
- ▶ bi-complex bi-disk and unit ball

Problem(s)

When does UCI hold?
When is the fixed point unique?

What about $\varphi(z) = \frac{1}{2}z^2 + \frac{1}{2}$?

Let's explore different domains:

- ▶ \mathbb{D}^2 (bi-disk)
- ▶ $\{\vec{z} \in \mathbb{C}^2: \|\vec{z}\| < 1\}$ (unit ball)
- ▶ bi-complex bi-disk and unit ball
- ▶ $(-1, 1)$
- ▶ $(-1, 1)^2$ (real bi-disk or unit square)
- ▶ $\{\vec{z} \in \mathbb{C}^2: \|\vec{z}\| < 1\}$ (real unit ball)