
Julia's Lemma and Wolff's Theorem For J^* -Algebras

Author(s): Kazimierz Włodarczyk

Source: *Proceedings of the American Mathematical Society*, Mar., 1987, Vol. 99, No. 3 (Mar., 1987), pp. 472-476

Published by: American Mathematical Society

Stable URL: <https://www.jstor.org/stable/2046348>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*

JSTOR

JULIA'S LEMMA AND WOLFF'S THEOREM FOR J^* -ALGEBRAS

KAZIMIERZ WŁODARCZYK

ABSTRACT. Julia's lemma and Wolff's theorem are established for (Fréchet-) holomorphic maps of bounded symmetric homogeneous domains in infinite dimensional complex Banach spaces called J^* -algebras.

1. Introduction. In [11] V. P. Potapov extended the classical lemma of G. Julia [8, p. 87] to matrix-valued holomorphic maps of a complex variable. Next, I. Glicksberg [6] and K. Fan [5] proved the versions of Julia's lemma for function algebras and for holomorphic maps of proper contraction operators in the sense of functional calculus, respectively.

In another paper [4], K. Fan extended the classical theorem of J. Wolff [14] on iterates of self-maps to holomorphic maps of proper contraction operators in the sense of functional calculus. Similar extensions of Wolff's theorem to (Fréchet-) holomorphic maps of the open unit ball and the generalized upper half-plane in \mathbb{C}^N were given by G. N. Chen [1]. Furthermore, Y. Kubota [9] and B. D. MacCluer [10], using differnet methods, proved some results on iterates of Wolff-Denjoy type [14, 2] in \mathbb{C}^N .

The main results of this paper are of the above two types (see §2). We first prove a general version of Julia's lemma for (Fréchet-)holomorphic maps of bounded symmetric homogeneous domains in infinite dimensional complex Banach spaces called J^* -algebras. We next prove, as an application of this, the extension of Wolff's theorem. In particular, our results imply Julia's lemma and Wolff's theorem for arbitrary complex Hilbert spaces (see §5), B^* -algebras, C^* -algebras, and ternary algebras.

2. Notions. Main results. Let H and K be Hilbert spaces over \mathbb{C} , let $\mathcal{L}(H, K)$ denote the Banach space of all bounded linear operators X from H to K with the operator norm, and let $\mathfrak{A} \in \mathcal{L}(H, K)$ be a J^* -algebra (see L. A. Harris [7]), i.e. a normed complex linear subspace of $\mathcal{L}(H, K)$ closed under the operation $X \rightarrow XX^*X$.

Let

$$\mathfrak{A}_0 = \{X \in \mathfrak{A} : \|X\| < 1\}, \quad \mathfrak{A}_1 = \{X \in \mathfrak{A} : \|X\| \leq 1\},$$

and, for $X \in \mathfrak{A}_0$, $Z \in \mathfrak{A}_1$, and $\alpha \geq 1$, let

$$W_Z(X) = (I_H - Z^*X)A_X^{-1}(I_H - X^*Z),$$

$$c_\alpha(Z, X) = \{I_H\alpha\|W_Z(X)\| + ZZ^*\}^{-1},$$

Received by the editors January 27, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46G20.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

$$r_\alpha(Z, X) = \|c_\alpha(Z, X)\|^{1/2} \|I_H\{\alpha\|W_Z(X)\| - 1\} + Z^*c_\alpha(Z, X)Z\|^{1/2},$$

where I_H is the identity map on H and $A_X = I_H - X^*X$.

We shall use these notations in proving the following Julia-type lemma for \mathfrak{A} .

LEMMA 2.1. *Let $F: \mathfrak{A}_0 \rightarrow \mathfrak{A}_0$ be a holomorphic map in \mathfrak{A}_0 . If $\{z_m\} \subset \mathfrak{A}_0$ is such that*

$$(2.1) \quad \lim_{m \rightarrow \infty} \|Z_m - V\| = 0$$

and

$$(2.2) \quad \lim_{m \rightarrow \infty} \|F(Z_m) - V\| = 0$$

for some $V \in \mathfrak{A}_1 \setminus \mathfrak{A}_0$, and if

$$(2.3) \quad \lim_{m \rightarrow \infty} \frac{\|A_{F(Z_m)}\|}{1 - \|Z_m\|^2} = \alpha \neq \infty,$$

then

$$(2.4) \quad \|X - c_1(V, X)V\| \leq r_1(V, X),$$

$$(2.5) \quad \|W_V[F(X)]\| \leq \alpha \|W_V(X)\|,$$

and

$$(2.6) \quad \|F(X) - c_\alpha(V, X)V\| \leq r_\alpha(V, X)$$

hold for all $X \in \mathfrak{A}_0$.

We can also use Lemma 2.1 to obtain a Wolff-type theorem for \mathfrak{A} .

THEOREM 2.2. *Let $F: \mathfrak{A}_0 \rightarrow \mathfrak{A}_0$ be a compact holomorphic map having no fixed point in \mathfrak{A}_0 and let $F^{[n]}$ denote the n th iterate of F (i.e., $F^{[1]} = F$ and $F^{[n]} = F \circ F^{[n-1]}$ for $n \geq 2$). Then there exist $\{Z_m\} \subset \mathfrak{A}_0$ and $V \in \mathfrak{A}_1 \setminus \mathfrak{A}_0$ such that*

$$\lim_{m \rightarrow \infty} \|Z_m - V\| = 0 \quad \text{and} \quad F(V) = V.$$

Moreover, $\|X - c_1(V, X)V\| \leq r_1(V, X)$ for all $X \in \mathfrak{A}_0$, and if

$$(2.7) \quad \lim_{m \rightarrow \infty} \frac{\|A_{Z_m}\|}{1 - \|Z_m\|^2} = \alpha \neq \infty,$$

then

$$(2.8) \quad \|W_V[F^{[n]}(X)]\| \leq \alpha \|W_V(X)\|$$

and

$$(2.9) \quad \|F^{[n]}(X) - c_\alpha(V, X)V\| \leq r_\alpha(V, X)$$

hold for all $x \in \mathfrak{A}_0$ and $n = 1, 2, \dots$

For the special case when $\mathfrak{A} = H$, see §5.

3. Proof of Lemma 2.1.

LEMMA 3.1. *If $F: \mathfrak{A}_0 \rightarrow \mathfrak{A}_0$ is a holomorphic map in \mathfrak{A}_0 , then*

$$(3.1) \quad \|W_{F(Z)}[F(X)]\| \leq \frac{\|A_{F(Z)}\|}{1 - \|Z\|^2} \|W_Z(X)\|$$

for all $X, Z \in \mathfrak{A}_0$.

PROOF. Let $X, Z \in \mathfrak{A}_0$. It follows from [13, Theorem 1(a)] that

$$(3.2) \quad \|A_{F(Z)}^{-1/2} W_{F(Z)}[F(X)] A_{F(Z)}^{-1/2}\| \leq \|A_Z^{-1/2} W_Z(X) A_Z^{-1/2}\|.$$

But

$$(3.3) \quad \|A_Z^{-1/2}\|^2 = \|A_Z^{-1}\| = (1 - \|Z\|^2)^{-1}$$

and

$$(3.4) \quad I_H \|A_{F(Z)}^{1/2}\|^{-1} \leq A_{F(Z)}^{-1/2} \quad \text{since } I_H \leq A_{F(Z)}^{-1/2}.$$

Thus, using (3.3) and (3.4), from (3.2) we get (3.1). This completes the proof.

LEMMA 3.2. *If $X \in \mathfrak{A}_0$, $Z \in \mathfrak{A}_1$, and D satisfy*

$$(3.5) \quad \|W_Z(X)\| \leq D,$$

then

$$(3.6) \quad \begin{aligned} & \|X - (I_H D + Z Z^*)^{-1} Z\| \\ & \leq \|(I_H D + Z Z^*)^{-1}\|^{1/2} \|I_H(D - 1) + Z^*(I_H D + Z Z^*)^{-1} Z\|^{1/2}. \end{aligned}$$

PROOF. Inequality (3.5) implies

$$\|A_X^{-1/2} (I_H - X^* Z) (I_H - Z^* X) A_X^{-1/2}\| \leq D$$

or, equivalently,

$$(I_H - X^* Z)(I_H - Z^* X) \leq A_X D.$$

Further, the above inequality is identical to

$$\begin{aligned} & \{X^* - Z^*(I_H D + Z Z^*)^{-1}\} (I_H D + Z Z^*) \{X - (I_H D + Z Z^*)^{-1} Z\} \\ & \leq I_H(D - 1) + Z^*(I_H D + Z Z^*)^{-1} Z. \end{aligned}$$

Consequently,

$$\|(I_H D + Z Z^*)^{1/2} \{X - (I_H D + Z Z^*)^{-1} Z\}\|^2 \leq \|I_H(D - 1) + Z^*(I_H D + Z Z^*)^{-1} Z\|.$$

Hence (3.6) follows. This completes the proof.

Now, we assume that $\{Z_m\} \subset \mathfrak{A}_0$ satisfies (2.1)–(2.3). Then, from Lemma 3.1 it follows that (2.5) holds, and (2.4) and (2.6) follow from the relations $\|W_Z(X)\| = D_1$ and $\|W_V[F(X)]\| \leq D_\alpha$, $D_\alpha = \alpha \|W_V(X)\|$, respectively, if we use Lemma 3.2.

4. Proof of Theorem 2.2.

LEMMA 4.1. *If $F: \mathfrak{A}_0 \rightarrow \mathfrak{A}_0$ is a compact holomorphic map having no fixed point in \mathfrak{A}_0 , then there exist $\{Z_m\} \subset \mathfrak{A}_0$ and a fixed point $V \in \mathfrak{A}_1 \setminus \mathfrak{A}_0$ of F such that*

$$\lim_{m \rightarrow \infty} \|Z_m - V\| = 0.$$

PROOF. Let $F_m = \alpha_m F$, $0 < \alpha_m < 1$, $m = 1, 2, \dots$, and let $\lim_{m \rightarrow \infty} \alpha_m = 1$. Let $\{Z_m\} \subset \mathfrak{A}_0$ be such that $F_m(Z_m) = Z_m$, $m = 1, 2, \dots$; such a sequence exists by the Earle-Hamilton theorem [3]. Since $F(\mathfrak{A}_0)$ is contained in a compact subset of \mathfrak{A} , F has no fixed points in \mathfrak{A}_0 and $F(Z_m) = Z_m/\alpha_m$, $m = 1, 2, \dots$, we may assume that $\lim_{m \rightarrow \infty} \|Z_m - V\| = 0$ and $\lim_{m \rightarrow \infty} \|F(Z_m) - V\| = \lim_{m \rightarrow \infty} \|Z_m/\alpha_m - V\| = 0$ for some $V \in \mathfrak{A}_1 \setminus \mathfrak{A}_0$. This completes the proof.

Let now $X \in \mathfrak{A}_0$ and let $F_m^{[n]}$ denote the n th iterate of F_m , $m = 1, 2, \dots$. By Lemma 4.1 and (3.1),

$$(4.1) \quad \|W_{Z_m}[F_m^{[n]}(X)]\| \leq \frac{\|A_{Z_m}\|}{1 - \|Z_m\|^2} \|W_{Z_m}(X)\|.$$

Furthermore (see [12, formula (4.6), p. 158]),

$$(4.2) \quad \lim_{m \rightarrow \infty} \|F_m^{[n]}(X) - F^{[n]}(X)\| = 0.$$

Thus, if (2.7) holds, from (4.1) and (4.2) we obtain (2.8). By Lemma 3.2, inequality (2.8) implies (2.9).

5. Concluding remarks. Let

$$H_0 = \{x \in H: \|x\| < 1\}, \quad H_1 = \{x \in H: \|x\| \leq 1\},$$

and, for $x \in H_0$, $v \in H_1 \setminus H_0$, and $\alpha \geq 1$, let

$$c_\alpha(v, x) = \frac{1 - \|x\|^2}{\alpha|1 - \langle x, v \rangle|^2 + 1 - \|x\|^2},$$

$$r_\alpha(v, x) = \frac{\alpha|1 - \langle x, v \rangle|^2}{\alpha|1 - \langle x, v \rangle|^2 + 1 - \|x\|^2}.$$

Identifying H with $\mathcal{L}(\mathbf{C}, H)$, as corollaries from our main results and their proofs we get the following Julia's lemma and Wolff's theorem for arbitrary complex Hilbert spaces.

LEMMA 5.1. *Let $F: H_0 \rightarrow H_0$ be a holomorphic map in H_0 . If $\{z_m\} \subset H_0$ is such that*

$$\lim_{m \rightarrow \infty} \|z_m - v\| = 0 \quad \text{and} \quad \lim_{m \rightarrow \infty} \|F(z_m) - v\| = 0$$

for some $v \in H_1 \setminus H_0$, and if

$$\lim_{m \rightarrow \infty} \frac{1 - \|F(z_m)\|^2}{1 - \|z_m\|^2} = \alpha \neq \infty,$$

then

$$\|x - c_1(v, x)v\| = r_1(v, x),$$

$$\frac{|1 - \langle F(x), v \rangle|^2}{1 - \|F(x)\|^2} \leq \alpha \frac{|1 - \langle x, v \rangle|^2}{1 - \|x\|^2},$$

and

$$\|F(x) - c_\alpha(v, x)v\| \leq r_\alpha(v, x)$$

for all $x \in H_0$.

THEOREM 5.2. *Let $F: H_0 \rightarrow H_0$ be a compact holomorphic map having no fixed points in H_0 and let $F^{[n]}$ denote the n th iterate of F . Then there exists $v \in H_1 \setminus H_0$ such that $F(v) = v$,*

$$\begin{aligned} \|x - c_1(v, x)v\| &= r_1(v, x), \\ \frac{|1 - \langle F^{[n]}(x), v \rangle|^2}{1 - \|F^{[n]}(x)\|^2} &\leq \frac{|1 - \langle x, v \rangle|^2}{1 - \|x\|^2}, \end{aligned}$$

and

$$\|F^{[n]}(x) - c_1(v, x)v\| \leq r_1(v, x)$$

for all $x \in H_0$ and $n = 1, 2, \dots$.

REMARKS. If $H = \mathbf{C}$, Lemma 5.1 and Theorem 5.2 imply the results of Julia [8, p. 87] and Wolff [14], respectively.

REFERENCES

1. G. N. Chen, *Iteration for holomorphic maps of the open unit ball and the generalized upper half-plane in \mathbf{C}^n* , J. Math. Anal. Appl. **98** (1984), 305–313.
2. A. Denjoy, *Sur l'iteration des fonctions analytiques*, C. R. Acad. Sci. Paris **182** (1926), 255–257.
3. C. J. Earle and R. S. Hamilton, *A fixed point theorem for holomorphic mappings*, Global Analysis (Berkeley, Calif., 1968), Proc. Sympos. Pure Math., vol. 16, Amer. Math. Soc., Providence, R.I., 1970, pp. 61–65.
4. K. Fan, *Iteration of analytic functions of operators*, Math. Z. **179** (1982), 293–298.
5. ———, *Julia's lemma for operators*, Math. Ann. **239** (1979), 241–245.
6. I. Glicksberg, *Julia's lemma for function algebras*, Duke Math. J. **43** (1976), 277–284.
7. L. A. Harris, *Bounded symmetric homogeneous domains in infinite dimensional spaces*, Lecture Notes in Math., vol. 364, Springer-Verlag, Berlin, Heidelberg and New York, 1974, pp. 13–40.
8. G. Julia, *Principes geometriques d'analyse*, Gauthier-Villars, Paris, 1930.
9. Y. Kubota, *Iteration of holomorphic maps of the unit ball into itself*, Proc. Amer. Math. Soc. **88** (1983), 476–480.
10. B. D. MacCluer, *Iterates of holomorphic self-maps of the unit ball in \mathbf{C}^N* , Michigan Math. J. **30** (1983), 97–106.
11. V. P. Potapov, *The multiplicative structure of J -contractive matrix functions*, Amer. Math. Soc. Transl. **15** (1960), 131–243.
12. K. Włodarczyk, *Iterations of holomorphic maps of infinite dimensional homogeneous domains*, Monatsh. Math. **99** (1985), 153–160.
13. ———, *Pick-Julia theorems for holomorphic maps in J^* -algebras and Hilbert spaces*, J. Math. Anal. Appl. **448** (1986).
14. J. Wolff, *Sur une généralisation d'un théorème de Schwarz*, C. R. Acad. Sci. Paris **182** (1926), 918–920; **183** (1926), 500–502.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ŁÓDŹ, BANACHA 22, P-90-238 ŁÓDŹ, POLAND