

A detailed illustration of a Red Porgy (Lutjanus campechanus) fish, shown in profile facing left. The fish has a robust, deep-bodied shape. Its coloration is primarily a vibrant reddish-orange, with lighter, almost white, areas on the belly and around the eyes. The dorsal fin is prominent, featuring a series of sharp, spiny rays. The pectoral fins are also visible, showing a similar reddish hue. The head is large with a prominent eye and a slightly open mouth. The overall appearance is that of a healthy, mature specimen.

Lee Cronin-Fine²

³University of Washington, School of Aquatic and Fishery Sciences

Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by NOAA Fisheries. It does not represent and should not be construed to represent any agency determination or policy.

Status of Pacific ocean perch (*Sebastes alutus*) along the US west coast in 2017

Contents

Executive Summary	1
Stock	1
Landings	1
Data and Assessment	2
Stock Biomass	4
Recruitment	7
Exploitation status	9
Ecosystem Considerations	12
Reference Points	12
Management Performance	13
Unresolved Problems And Major Uncertainties	13
Decision Table(s) (groundfish only)	14
Research And Data Needs	19
Rebuilding Projections	19
1 Introduction	20
1.1 Basic Information	20
1.2 Map	21
1.3 Life History and Ecosystem Considerations	22
1.4 Fishery Information	22
1.5 Summary of Management History	22
1.6 Management Performance	22
1.7 Fisheries off Canada, Alaska, and/or Mexico	22

42	2	Assessment	23
43	2.1	Data	23
44	2.1.1	Commercial Fishery Landings	23
45	2.1.2	Abundance Indices	25
46	2.1.3	Fishery-Dependent Data:	25
47	2.1.4	Fishery-Independent Data:	25
48	2.1.5	Biological Parameters and Data	27
49	2.1.6	Environmental Or Ecosystem Data Included In The Assessment . . .	29
50	2.2	History Of Modeling Approaches Used For This Stock	29
51	2.2.1	Previous Assessments	29
52	2.2.2	Previous Assessment Recommendations	29
53	2.3	Model Description	30
54	2.3.1	Transition To The Current Stock Assessment	30
55	2.3.2	Definition of Fleets and Areas	30
56	2.3.3	Summary of Data for Fleets and Areas	30
57	2.3.4	Modeling Software	30
58	2.3.5	Data Weighting	30
59	2.3.6	Priors	31
60	2.3.7	General Model Specifications	31
61	2.3.8	Estimated And Fixed Parameters	31
62	2.4	Model Selection and Evaluation	31
63	2.4.1	Key Assumptions and Structural Choices	31
64	2.4.2	Alternate Models Considered	31
65	2.4.3	Convergence	31
66	2.5	Response To The Current STAR Panel Requests	32
67	2.6	Model 1	32
68	2.6.1	Model 1 Base Case Results	32
69	2.6.2	Model 1 Uncertainty and Sensitivity Analyses	32
70	2.6.3	Model 1 Retrospective Analysis	33
71	2.6.4	Model 1 Likelihood Profiles	33
72	2.6.5	Model 1 Harvest Control Rules (CPS only)	33
73	2.6.6	Model 1 Reference Points (groundfish only)	33

74	3 Harvest Projections and Decision Tables	33
75	4 Regional Management Considerations	33
76	5 Research Needs	34
77	6 Acknowledgments	34
78	7 Tables	35
79	8 Figures	62
80	References	

81 ???

82 Executive Summary

executive-summary

83 Stock

stock

84 This assessment reports the status of the Pacific ocean perch (*Sebastes alutus*) species off
85 rockfish off the U.S. West Coast from Northern California to the Canadian Border using
86 data through 2017. Pacific ocean perch are most abundant in the Gulf of Alaska and have
87 observed off of Japan, in the Bering Sea, and south to Baja California, although they are
88 sparse south of Oregon and rare in southern California. Composition data indicate that
89 good recruitment years coincide in Oregon and Washington. To date, no significant genetic
90 differences have been found in the range covered by this assessment.

91 Landings

landings

92 The first year that harvest of Pacific ocean perch exceeded 1 mt off the U.S. West Coast
93 first occurred in 1929. Catches ramped up in the 1940s with large removals in Washington
94 waters. During the 1950s the removals primarily occurred in Oregon waters with catches from
95 Washington declining following the 1940s. The largest removals in 1966-1968 were largely a
96 result of harvest by foreign vessels. The fishery proceeded with more moderate removals ranging
97 between 1,200 to 2,600 metric tons per year between 1969 to 1980. Removals generally
98 declined from 1981 to 1994 to between 1,000 and 1,700 metric tons per year. Pacific ocean
99 perch was declared overfished in 1999 resulting in large reduction in harvest in recent years
100 since the declaration.

Table a: Landings (mt) for the past 10 years for Pacific ocean perch by fleet.

Year	California	Oregon	Washington	At-sea Hake	Research	<code>tab:Exec_catch</code> Total
						Landings
2007	0.15	83.65	45.12	4.05	0.58	133.55
2008	0.39	58.64	16.61	15.93	0.80	92.37
2009	0.92	58.75	33.22	1.56	2.72	97.17
2010	0.14	58.00	22.29	16.87	1.68	98.98
2011	0.12	30.26	19.66	9.17	1.94	61.14
2012	0.18	30.41	21.79	4.52	1.62	58.51
2013	0.08	34.86	14.83	5.41	1.71	56.89
2014	0.18	33.92	15.82	3.92	0.57	54.41
2015	0.12	38.12	11.41	8.71	1.59	59.95
2016	0.19	34.15	13.12	10.30	0.12	57.87

Data and Assessment

`data-and-assessment`

This a new full assessment for Pacific ocean perch which was last assessed in 2011. In this assessment, all aspects of the model including catches, data, and modelling assumptions were re-evaluated as much as possible. The assessment was conducted using the length- and age-structured modeling software Stock Synthesis (version 3.30). The coastwide population was modeled assuming separate growth and mortality parameters for each sex (a two-sex model) from 1918 to 2017, and forecasted beyond 2017.




Figure a: Landings of Pacific ocean perch for California, Oregon, Washington, the Foreign fishery (1966-1976), At-Sea Hake fishery, and fishery independent surveys. fig:Exec_catch1

Stock Biomass

stock-biomass

Include: trends and current levels relative to virgin or historic levels, description of uncertainty-include table for last 10 years and graph with long term estimates.

Spawning output Figure: Figure [b](#)

Spawning output Table(s): Table [b](#)

Relative depletion Figure: Figure [c](#)

Example text (remove Models 2 and 3 if not needed - if using, remove the # in-line comments!!!)
The estimated relative depletion level (spawning output relative to unfished spawning output) of the the base-case model in 2017 is 38% (~95% asymptotic interval: \pm -18.2%-94.1%) (Figure [c](#)).

Table b: Recent trend in estimated spawning output (million eggs) and relative spawning output.

tab:SpawningDeplete_mod1				
Year	Spawning Output (million eggs)	~ 95% confidence interval	Estimated depletion	~ 95% confidence interval
2008	1100.00	-590 - 2790	0.19	-0.084 - 0.472
2009	1155.00	-628 - 2939	0.20	-0.090 - 0.498
2010	1203.00	-664 - 3070	0.21	-0.095 - 0.520
2011	1245.00	-698 - 3189	0.22	-0.101 - 0.540
2012	1288.00	-726 - 3302	0.23	-0.105 - 0.560
2013	1336.00	-757 - 3429	0.24	-0.110 - 0.581
2014	1442.00	-820 - 3704	0.26	-0.119 - 0.628
2015	1656.00	-946 - 4258	0.29	-0.137 - 0.722
2016	1919.00	-1104 - 4942	0.34	-0.161 - 0.838
2017	2151.00	-1245 - 5547	0.38	-0.182 - 0.941




Figure b: Time series of spawning output trajectory (circles and line: median; light broken lines: 95% credibility intervals) for the base case assessment model. fig:Spawnbio_all




Figure c: Estimated relative depletion with approximate 95% asymptotic confidence intervals (dashed lines) for the base case assessment model. fig:RelDeplete_all

Recruitment

recruitment

Include: trends and current levels relative to virgin or historic levels-include table for last 10 years and graph with long term estimates.

Recruitment Figure: (Figure d)

Recruitment Tables: (Tables c)

Table c: Recent estimated trend in recruitment with approximate 95confidence intervals determined from the base model

Year	Estimated Recruitment	~ 95% confidence interval	Estimated Recruitment Devs.	~ 95% confidence interval
2008	70862.00	17357 - 289304	2.98	2.605 - 3.358
2009	1947.00	414 - 9162	-0.67	-1.598 - 0.261
2010	3171.00	642 - 15655	-0.23	-1.319 - 0.862
2011	4069.00	826 - 20047	-0.02	-1.090 - 1.045
2012	3956.00	781 - 20047	-0.09	-1.262 - 1.075
2013	5973.00	1159 - 30775	0.27	-0.973 - 1.520
2014	4345.00	852 - 22172	-0.12	-1.383 - 1.150
2015	5358.00	1046 - 27437	-0.00	-1.372 - 1.372
2016	5888.00	1193 - 29070	0.00	-1.372 - 1.372
2017	6312.00	1575 - 25298	0.00	-0.970 - 0.970




Figure d: Time series of estimated Pacific ocean perch recruitments for the base-case model with 95% confidence or credibility intervals. `fig:Recruits_all`

Exploitation status

exploitation-status

Include: exploitation rates (i.e., total catch divided by exploitable biomass, or the annual SPR harvest rate) include a table with the last 10 years of data and a graph showing the trend in fishing mortality relative to the target (y-axis) plotted against the trend in biomass relative to the target (x-axis).

Exploitation Tables: Table d, Table ??, Table ?? Exploitation Figure: Figure e).

A summary of Pacific ocean perch exploitation histories for base model is provided as Figure f.

Table d: Recent trend in spawning potential ratio (1-SPR) and summary exploitation rate for Pacific ocean perch.

Year	Fishing intensity	~ 95% confidence interval	Exploitation rate	tab:SPR_Exploit_mod1 ~ 95% confidence interval
2007	0.279	-0.097 - 0.655	0.006	-0.003 - 0.015
2008	0.257	-0.096 - 0.611	0.005	-0.003 - 0.014
2009	0.266	-0.101 - 0.633	0.006	-0.003 - 0.015
2010	0.250	-0.099 - 0.598	0.006	-0.003 - 0.014
2011	0.099	-0.049 - 0.247	0.002	-0.001 - 0.004
2012	0.090	-0.045 - 0.226	0.001	-0.001 - 0.004
2013	0.083	-0.042 - 0.208	0.001	-0.001 - 0.003
2014	0.071	-0.037 - 0.178	0.001	-0.001 - 0.003
2015	0.068	-0.036 - 0.172	0.001	-0.001 - 0.003
2016	0.058	-0.032 - 0.149	0.001	-0.001 - 0.003




Figure e: Estimated spawning potential ratio (SPR) for the base-case model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the $SPR_{50\%}$ harvest rate. The last year in the time series is 2016. fig:SPR_all




Figure f: Phase plot of estimated relative (1-SPR) vs. relative spawning biomass for the base case model. The relative (1-SPR) is (1-SPR) divided by 50% (the SPR target). Relative depletion is the annual spawning biomass divided by the unfished spawning biomass. fig:Phase_all

Ecosystem Considerations

ecosystem-considerations

In this assessment, ecosystem considerations were....

Reference Points

reference-points

Include: management targets and definition of overfishing, including the harvest rate that brings the stock to equilibrium at $B_{40\%}$ (the B_{MSY} proxy) and the equilibrium stock size that results from fishing at the default harvest rate (the F_{MSY} proxy). Include a summary table that compares estimated reference points for SSB, SPR, Exploitation Rate and Yield based on SSBproxy for MSY, SPRproxy for MSY, and estimated MSY values

Write intro paragraph

This stock assessment estimates that Pacific ocean perch in the Base model are below the biomass target, but above the minimum stock size threshold. **Add sentence about spawning output trend.** The estimated relative depletion level for **Model 1** in 2017 is 38% (~95% asymptotic interval: \pm -18.2%-94.1%, corresponding to an unfished spawning output of 2151 million eggs (~95% asymptotic interval: -1245.35559241101-5547.095592411 million eggs) of spawning output in the base model (Table e). Unfished age 3+ biomass was estimated to be 122166 mt in the base case model. The target spawning output based on the biomass target ($SB_{40\%}$) is 2265.8 million eggs, which gives a catch of 1176.1 mt. Equilibrium yield at the proxy F_{MSY} harvest rate corresponding to $SPR_{50\%}$ is 1018 mt.

Table e: Summary of reference points and management quantities for the base case.

Quantity	Estimate	tab:Ref_pts_mod1
		95% Confidence Interval
Unfished spawning output (million eggs)	5664.4	4293.9 - 7034.9
Unfished age 3+ biomass (mt)	122166	94382.3 - 149949.7
Unfished recruitment (R0, thousands)	10014.3	7894.9 - 12702.7
Spawning output(2017 million eggs)	2150.9	-1245.356 - 5547.1
Depletion (2017)	0.38	-0.182 - 0.941
Reference points based on SB_{40%}		
Proxy spawning output ($B_{40\%}$)	2265.8	1717.6 - 2814
SPR resulting in $B_{40\%}$ ($SPR_{B40\%}$)	0.615	0.443 - 0.788
Exploitation rate resulting in $B_{40\%}$	0.022	0.008 - 0.036
Yield with $SPR_{B40\%}$ at $B_{40\%}$ (mt)	1176.1	287.7 - 2064.5
Reference points based on SPR proxy for MSY		
Spawning output	1245	-785.915 - 3275.9
SPR_{proxy}	0.5	
Exploitation rate corresponding to SPR_{proxy}	0.033	0.033 - 0.034
Yield with SPR_{proxy} at SB_{SPR} (mt)	1018	-630.883 - 2666.8
Reference points based on estimated MSY values		
Spawning output at MSY (SB_{MSY})	2176.7	1496.9 - 2856.5
SPR_{MSY}	0.605	0.374 - 0.837
Exploitation rate at MSY	0.023	0.003 - 0.043
MSY (mt)	1177.3	275.5 - 2079.1

Management Performance

management-performance

Include: catches in comparison to OFL, ABC and OY/ACL values for the most recent 10 years (when available), overfishing levels, actual catch and discard. Include OFL(encountered), OFL(retained) and OFL(dead) if different due to discard and discard mortality.

Management performance table: Table f

Unresolved Problems And Major Uncertainties

unresolved-problems-and-major-uncertainties

TBD after STAR panel

Table f: Recent trend in total catch and commercial landings (mt) relative to the management guidelines. Estimated total catch reflect the commercial landings plus the model estimated discarded biomass.

Year	OFL (mt; ABC prior to 2011)	ABC (mt)	ACL (mt; OY prior to 2011)	tab:mnmgt_perform	
				Total landings (mt)	Estimated total catch (mt)
2007	-	-	150	134	158
2008	-	-	150	92	151
2009	-	-	189	97	169
2010	-	-	200	99	161
2011	-	-	180	61	61
2012	-	-	183	59	59
2013	-	-	150	57	58
2014	-	-	153	54	55
2015	-	-	158	60	60
2016	-	-	164	58	58

Decision Table(s) (groundfish only)

decision-tables-groundfish-only

Include: projected yields (OFL, ABC and ACL), spawning biomass, and stock depletion levels for each year. Not required in draft assessments undergoing review.

OFL projection table: Table [g](#)

Decision table(s) Table [h](#), Table ??, Table ??

Yield curve: Figure [\ref{fig:Yield_all}](#)




Figure g: Equilibrium yield curve for the base case model. Values are based on the 2016 fishery selectivity and with steepness fixed at... `fig:Yield_all`

Table g: Projections of potential OFL (mt) and ACL (mt) and the estimated spawning output and relative biomass.

tab:OFL_projection				
Year	OFL	ACL	Spawning Output (million eggs)	Relative Biomass
2017	1899	1783	2151	0.380
2018	1988	1898	2257	0.398
2019	2018	1929	2312	0.408
2020	2008	1919	2337	0.413
2021	1975	1888	2342	0.413
2022	1935	1850	2333	0.412
2023	1894	1811	2316	0.409
2024	1858	1776	2296	0.405
2025	1827	1747	2274	0.402
2026	1801	1718	2252	0.398
2027	1779	1692	2230	0.394
2028	1760	1669	2210	0.390

Table h: Summary of 10-year projections beginning in 2019 for alternate states of nature based on an axis of uncertainty for the base model. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. An entry of "–" indicates that the stock is driven to very low abundance under the particular scenario.

tab:Decision_table_mod1

		States of nature					
		Low M 0.05		Base M 0.07		High M 0.09	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
40-10 Rule, Low M	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
40-10 Rule	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
40-10 Rule, High M	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
Average Catch	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-

Table i: Base model results summary.

Quantity	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	tab:base summary
Landings (mt)	-	-	-	-	-	-	-	-	-	-	-
Total Est. Catch (mt)	150	189	200	180	183	150	153	158	164	164	281
OFL (mt)	92	97	99	61	59	57	54	60	58	58	58
ACL (mt)	151	169	161	61	59	58	55	60	58	58	58
(1-SPR)(1-SPR _{50%})	0.26	0.27	0.25	0.10	0.09	0.08	0.07	0.07	0.06	0.06	0.06
Exploitation rate	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Age 3+ biomass (mt)	27737.5	28463.0	29077.8	37251.2	41154.9	44993.1	48570.4	51699.7	54560.3	56851.5	56851.5
Spawning Output	1100	1155	1203	1245	1288	1336	1442	1656	1919	2151	2151
95% CI	-590 - 2790	-628 - 2939	-664 - 3070	-698 - 3189	-726 - 3302	-757 - 3429	-820 - 3704	-946 - 4258	-1104 - 4942	-1245 - 5547	-1245 - 5547
Depletion	0.194	0.204	0.212	0.220	0.227	0.236	0.255	0.292	0.339	0.380	0.380
95% CI	-0.084 - 0.472	-0.090 - 0.498	-0.095 - 0.520	-0.101 - 0.540	-0.105 - 0.560	-0.110 - 0.581	-0.119 - 0.628	-0.137 - 0.722	-0.161 - 0.838	-0.182 - 0.941	-0.182 - 0.941
Recruits	70862	1947	3171	4069	3956	5973	4345	5358	5888	6312	6312
95% CI	17357 - 289304	414 - 9162	642 - 15655	826 - 20047	781 - 20047	1159 - 30775	852 - 22172	1046 - 27437	1193 - 29070	1575 - 25298	1575 - 25298

Research And Data Needs

research-and-data-needs

Include: identify information gaps that seriously impede the stock assessment.

We recommend the following research be conducted before the next assessment:

1. List item No. 1 in the list
2. List item No. 2 in the list, etc.

Rebuilding Projections

rebuilding-projections

Include: reference to the principal results from rebuilding analysis if the stock is overfished. This section should be included in the Final/SAFE version assessment document but is not required for draft assessments undergoing review. See Rebuilding Analysis terms of reference for detailed information on rebuilding analysis requirements.

1 Introduction

introduction

1.1 Basic Information

basic-information

Pacific ocean perch (*Sebastes alutus*) are most abundant in the Gulf of Alaska, and have been observed off of Japan, in the Bering Sea, and south to Baja California, although they are sparse south of Oregon and rare in southern California. While genetic studies have found three populations of Pacific ocean perch off of British Columbia (Seeb and Gunderson 1988, Withler et al. 2001) with, notably, a separate stock off of Vancouver Island, no significant genetic differences have been found in the range covered by this assessment. Pacific ocean perch show dimorphic growth, with females reaching a slightly large size than males. Males and females are equally abundant on rearing grounds at age 1.5.

The Pacific ocean perch population has been modeled as a single stock off of the U.S. West Coast (essentially northern California to the Canadian border, since Pacific ocean perch are seen extremely rarely in central and southern California). Good recruitments show up in size-composition data throughout all portions of this area, which supports the single stock hypothesis. This assessment includes landings and catch data for Pacific ocean perch from the states of Washington, Oregon and California, along with records from foreign fisheries, the at-sea hake fleet, and surveys.

Prior to 1966, the Pacific ocean perch resource off of the northern portion of the U.S. West Coast was harvested almost entirely by Canadian and United States vessels. Harvest was negligible prior to 1940, reached 1,300 mt in 1950, 3,200 mt in 1961 and exceeded 7,600 mt in 1965. Catches increased dramatically after 1965, with the introduction of large distant-water fishing fleets from the Soviet Union and Japan. Both nations employed large factory stern trawlers as their primary method for harvesting Pacific ocean perch. Peak removals by all foreign nations combined are estimated at over 15,000 mt in 1966 and remained over 12,000 mt in 1967. These numbers are based upon a re-analysis of the foreign catch data (Rogers 2003), which focused on deriving a more realistic species composition for catches previously identified only as Pacific ocean perch. Catches declined rapidly following these peak years, and Pacific ocean perch stocks were considered to be severely depleted throughout the Oregon-Vancouver Island region by 1969 (Gunderson 1977, Gunderson et al. 1977). Landed harvest averaged 1,350 mt over the period 1977-94. Landings have continued to decline since 1994, primarily due to more restrictive management.

Prior to 1977, Pacific ocean perch in the northeast Pacific were managed by the Canadian Government in its waters and by the individual states in waters off of the United States. With implementation of the Magnuson Fishery Conservation and Management Act (MFCMA) in 1977, U.S. territorial waters were extended to 200 miles from shore, and primary responsibility for management of the groundfish stocks off Washington, Oregon and California shifted from the states to the Pacific Fishery Management Council (PFMC) and the National Marine Fisheries Service (NMFS). At that time, however, a Fishery Management Plan (FMP) for the

west coast groundfish stocks had not yet been approved. In the interim, the state agencies worked with the PFMC to address conservation issues. In 1981, the PFMC adopted a management strategy to rebuild the depleted Pacific ocean perch stocks to levels that would produce Maximum Sustainable Yield (MSY) within 20 years. On the basis of cohort analysis (Gunderson 1978), the PFMC set Acceptable Biological Catch (ABC) levels at 600 mt for the US portion of the Vancouver INPFC area and 950 mt for the Columbia INPFC area. To implement this strategy, the states of Oregon and Washington each established landing limits for Pacific ocean perch. Trawl trip limits of various forms remained in effect through 2010 (1).

Age estimates for Pacific ocean perch prior to the 1980s were made via surface ageing of otoliths, which misses the very tight annuli at the edge of the otolith once the fish reaches near maximum size. Ages are biased by around age 10-12, and maximum age was estimated to be in the 20s, which lead to an overestimate of the natural mortality rate and the productivity of the stock. Using break and burn methods, Pacific ocean perch have been aged to over 100 years, and we now know that the underlying assumptions of the early models were overly optimistic about productivity. Research surveys have been used to provide fishery-independent information about the abundance, distribution, and biological characteristics of Pacific ocean perch. A coast-wide survey of the rockfish resource was conducted in 1977 (Gunderson and Sample 1980) and was repeated every three years through 2004 (referred to as the ‘Triennial Survey’). The National Marine Fisheries Service (NMFS) coordinated a cooperative research survey of the Pacific ocean perch stocks off Washington and Oregon with the Washington Department of Fisheries (WDFW) and the Oregon Department of Fish and Wildlife (ODFW) in March-May 1979 (Wilkins and Golden 1983). This survey was repeated in 1985 (referred to as the Pacific ocean perch Survey). Two slope surveys have been conducted on the West Coast in recent years, one using the research vessel Miller Freeman, which ended in 2001 (referred to as the ‘AFSC Slope Survey’), and another ongoing cooperative survey using commercial fishing vessels which began in 1998 as a DTS (Dover sole, thornyhead and sablefish) survey, was expanded to other groundfish in 1999 (referred to as the ‘NWFSC Slope Survey’). In 2003, this survey was expanded spatially to include the shelf. This last survey, conducted by the NWFSC, continues to cover depths from 30-700 fathoms (55-1280 meters) on an annual basis (referred to as the ‘NWFSC shelf-slope Survey’).

1.2 Map

map

A map showing the scope of the assessment and depicting boundaries for fisheries or data collection strata is provided in Figure 1.

1.3 Life History and Ecosystem Considerations

life-history-and-ecosystem-considerations

Include: Ecosystem considerations (e.g., ecosystem role and trophic relationships of the species, habitat requirements/preferences, relevant data on ecosystem processes that may affect stock or parameters used in the stock assessment, and/or cross-FMP interactions with other fisheries). This section should note if environmental correlations or food web interactions were incorporated into the assessment model. The length and depth of this section would depend on availability of data and reports from the IEA, expertise of the STAT, and whether ecosystem factors are informational to contribute quantitative information to the assessment.

1.4 Fishery Information

fishery-information

Include: Important features of current fishery and relevant history of fishery.

1.5 Summary of Management History

summary-of-management-history

Include: Summary of management history (e.g., changes in mesh sizes, trip limits, or other management actions that may have significantly altered selection, catch rates, or discards).

1.6 Management Performance

management-performance-1

Include: Management performance, including a table or tables comparing Overfishing Limit (OFL), Annual Catch Limit (ACL), Harvest Guideline (HG) [CPS only], landings, and catch (i.e., landings plus discard) for each area and year.

Management performance table: (Table f)

A summary of these values as well as other base case summary results can be found in Table i.

1.7 Fisheries off Canada, Alaska, and/or Mexico

fisheries-off-canada-alaska-andor-mexico

Include if necessary.

2 Assessment

assessment

2.1 Data

data

Data used in the Pacific ocean perch assessment are summarized in Figure 2. A description of each data source is provided below.

2.1.1 Commercial Fishery Landings

commercial-fishery-landings

Washington

Historical commercial fishery landings of Pacific ocean perch from Washington for the years 1918-2016 were obtained from Theresa Tsou (WDFW) and Phillip Weyland (WDFW). This assessment is the first Pacific ocean perch assessment to include a state provide historical catch reconstruction and hence, the historical catches for Washington vary markedly from those used in the 2011 assessment. Due to Recent landings (1981-2016) were obtained directly from Washington state rather than from PacFIN (Pacific Fisheries Information Network (PacFIN) due to identified missing catches not available within PacFIN for Pacific ocean perch.

Oregon

Historical commercial fishery landings of Pacific ocean perch from Oregon for the years 1892-1986 were obtained from Alison Dauble (ODFW). A description of the methods can be found in Karnowski et al. (2014). Recent landings (1987-2016) were obtained from PacFIN retrieval dated March 3, 2015, Pacific States Marine Fisheries Commission, Portland, Oregon; www.psmfc.org). The catch data in from the POP and POP2 categories contained within PacFIN for Pacific ocean perch were used for this assessment. Additional catches from 1987-1999 for Pacific ocean perch under the UROCK category not yet available in PacFIN were received directly from the state and combined with the catch data available for that period within PacFIN.

California

Historical commercial fishery landings of Pacific ocean perch were obtained from the on-line database of the California Cooperative Groundfish Survey, also known as CALCOM (128.114.3.187) for the years 1916-1980. A description of the methods can be found in (Ralston et al. 2010). Recent landings (1981-2016) were obtained from PacFIN (Pacific Fisheries Information Network (PacFIN) retrieval dated March 3, 2015, Pacific States Marine Fisheries Commission, Portland, Oregon; www.psmfc.org).

At-sea fishery

Catches of Pacific ocean perch are monitored aboard the vessel by observers in the At-Sea hake Observer program (ASHOP) and were available for the years of 1975-2016. Observers use a spatial sample design, based on weight, to randomly choose a portion of the haul to sample for species composition. For the last decade, this is typically 30-50% of the total weight. The total weight of the sample is determined by all catch passing over a flow scale. All species other than hake are removed and weighed, by species, on a motion compensated flatbed scale. Observers record the weights of all non-hake species. Non-hake species total weights are expanded in the database by using the proportion of the haul sampled to the total weight of the haul. The catches of non-hake species in unsampled hauls is determined using bycatch rates determined from sampled hauls. Since 2001, more than 97% of the hauls have been observed and sampled.

Foreign

From the 1960s through the early 1970s, foreign trawling enterprises harvested considerable amounts of rockfish off Washington and Oregon, and along with the domestic trawling fleet, landed large quantities of Pacific ocean perch. Foreign catches of individual species were estimated by Rogers (2003) and attributed to INPFC areas for the years of 1966-1976 for Pacific ocean perch. The foreign catches were combined across areas for a coastwide removal total.

Discards

Data on discards of Pacific ocean perch are available from two different data sources. The earliest source is called the Pikitch data and comes from a study organized by Ellen Pikitch that collected trawl discards from 1985-1987 (Pikitch et al. 1988). The northern and southern boundaries of the study were 48°42' N latitude and 42°60' N. latitude respectively, which is primarily within the Columbia INPFC area (Pikitch et al. 1988 , Rogers and Pikitch 1992). Participation in the study was voluntary and included vessels using bottom, midwater, and shrimp trawl gears. Observers of normal fishing operations on commercial vessels collected the data, estimated the total weight of the catch by tow and recorded the weight of species retained and discarded in the sample. Results of the Pikitch data were obtained from John Wallace (NWFSC, personal communication) in the form of ratios of discard weight to retained weight of Pacific ocean perch and sex-specific length frequencies. Discard estimates are shown in Table 3. Length compositions for discards show a wide range of sizes being discarded, with a peak are XX cm (Cite figure).

The second source is from the West Coast Groundfish Observer Program (WCGOP). This program is part of the NWFSC and has been recording discard observations since 2003. Table ??{tab:Discard} shows the discard ratios of Pacific ocean perch from the WCGOP. Since 2011, when the trawl rationalization proram was implemented, observer coverage rates increased to nearly 100% for all the limited entry trawl vessels in the program and discard rates declined compared to pre-2011 rates. Discard rates were obtained for both the catch-share and the non-catch share sector for Pacific ocean perch. A single discard rate was calculated by weighting discard rates based on the commercial landings by each sector. Discard length

composition for the trawl fleet varied by year, with larger fish being discarded prior to 2011 (Figure 3).

2.1.2 Abundance Indices

abundance-indices

2.1.3 Fishery-Dependent Data:

fishery-dependent-data

Historical Commercial Catch-per-unit effort

Data on catch-per-unit-effort (CPUE) in mt/hr from the domestic fishery were combined for the INPFC Vancouver and Columbia areas (15 from Gunderson (1977)). Although these data reflect catch rates for the US fleet, the highest catch rates coincided with the beginning of removals by the foreign fleet. This suggest that, barring unaccounted changes in fishing efficiency during this period, the level of abundance was high at that time. A CV of 0.40 was used in this assessment to be consistent with the CV observed in the survey data.

2.1.4 Fishery-Independent Data:

fishery-independent-data

Northwest Fisheries Science Center (NWFSC) shelf-slope survey

The NWFSC shelf-slope survey is based on a random-grid design; covering the coastal waters from a depth of 55 m to 1,280 m (Bradburn et al. 2011). This design uses four chartered industry vessels in most years, assigned to a roughly equal number of randomly selected grid cells. The survey, which has been conducted from late-May to early-October each year, is divided into two 2-vessel passes of the coast, which are executed from north to south. This design therefore incorporates both vessel-to-vessel differences in catchability as well as variance associated with selecting a relatively small number (~700) of cells from a very large population of possible cells (greater than 11,000) distributed from the Mexican to the Canadian border.

The data from the NWFSC shelf-slope survey was analyzed using a spatial delta-generalized linear mixed model (delta-GLMM) (Thorson and Barnett 2017). Predicted fish biomass density is derived as the product of a “delta” portion for the probability of a non-zero catch and a second portion for the magnitude of the non-zero catches. Further, the geostatistical GLMM framework can accommodate spatial autocorrelation. Additional information about the approach and the software package it is implemented in are available from www.fishstats.org. describe VAST

The estimated index of abundance is shown in Table 16.

Northwest Fisheries Science Center (NWFSC) slope survey

The NWFSC slope survey covered waters throughout the summer from 183 m to 1280 m north of 34°30' S, which is near Point Conception. The survey strata used to expand the biomass data for this assessment are shown in Table 5.

The estimated index of abundance is shown in Table 16.

Alaska Fisheries Science Center (AFSC) slope survey

The AFSC slope survey operated during autumn (October-November) aboard the R/V Miller Freeman. Partial survey coverage of the U.S. west coast occurred during 1988-96 and complete coverage (north of 34°30' S) during 1997, 1999, 2000, and 2001. Only the four years of consistent and complete surveys plus 1996, which surveyed north of 43° N latitude to the U.S.-Canada border, were used in this assessment. The number of tows with length data ranged from 19 in 2000 to 48 in 1996 7. Because a large number of positive tows occurred in 1996, it was decided to include that year, which surveyed from 43° N latitude to the U.S.-Canada border. Therefore, only tows from 43° N latitude to the U.S.-Canada border were used.

The estimated index of abundance is shown in Table 16.

Triennial Bottom Trawl Survey

The triennial survey was first conducted by the AFSC in 1977 and spanned the timeframe from 1977-2004. The survey's design and sampling methods are most recently described in (Weinberg et al. 2002). Its basic design was a series of equally-spaced transects from which searches for tows in a specific depth range were initiated (Figure 5). The survey design has changed slightly over the period of time (Table 4, Figure 3). In general, all of the surveys were conducted in the mid-summer through early fall: the 1977 survey was conducted from early July through late September; the surveys from 1980 through 1989 ran from mid-July to late September; the 1992 survey spanned from mid-July through early October; the 1995 survey was conducted from early June to late August; the 1998 survey ran from early June through early August; and the 2001 and 2004 surveys were conducted in May-July (Figure 4).

Haul depths ranged from 91-457 m during the 1977 survey with no hauls shallower than 91 m. The surveys in 1980, 1983, and 1986 covered the West Coast south to 36.8° N latitude and a depth range of 55-366 meters. The surveys in 1989 and 1992 covered the same depth range but extended the southern range to 34.5° N (near Point Conception). From 1995 through 2004, the surveys covered the depth range 55-500 meters and surveyed south to 34.5° N. In the final year of the triennial series (2004), the NWFSC's Fishery Resource and Monitoring division (FRAM) conducted the survey and followed very similar protocols as the AFSC.

Given the different depths surveyed during 1977, the data from that year were not included in this assessment. Water hauls (Zimmermann et al. 2003) and tows located in Canadian waters were also excluded from the analysis of this survey. The survey was analyzed as an early series (1980-1992) and a late series (1995-2004), as has been done in other West Coast rockfish assessments.

Describe whether the time-series was split or retained as one index

The estimated index of abundance is shown in Table 16.

Pacific ocean perch Survey

A survey targeted designed to sample Pacific ocean perch was conducted in 1979 and again in 1985. The estimated index of abundance is shown in Table 16.

2.1.5 Biological Parameters and Data

biological-parameters-and-data

Length And Age Compositions

Include: Sample size information for length and age composition data by area, year, gear, market category, etc., including both the number of trips and fish sampled.

Length compositions were provided from the following sources, by region, with brief descriptions below:

- Commercial fishery - landed: 1966-2016
- Commercial fishery - discard: 2004-2015
- At-sea hake fishery: 2003-2016
- Pacific ocean perch Survey: 1979 and 1985
- Triennial Survey: 1980, 1983, 1986, 1989, 1992, 1995, 1998, 2001, 2004
- AFSC Slope Survey: 1996-2001
- NWFS Slope Survey: 2001-2002
- NWFSC Shelf-Slope Survey: 2003-2016

Commercial: PacFIN

Research: NWFSC shelf-slope survey

Research: NWFSC slope survey

Age Structures

Age structure data were available from the following sources:

Model Region 1

- Source No. 1 (*ex. research, commercial dead fish, live fish, etc.*, date range (ex. 2010-2011))

- Source No. 2 (*ex. research, commercia dead fish, live fish, etc,*
date range (ex. 2010-2011))
- etc...
- Begin sublist if desired
 - Sublist source No. 1
 - Sublist source No. 2
 - etc...
- Back to main list, next Source
- Last Source

Natural mortality

Historic Pacific ocean perch ages determined using scales and surface reading methods of otoliths, resulted in estimates of natural mortality of between 0.10 and 0.20yr⁻¹ with a longevity less than 30 years (Gunderson 1977). Based on break-and-burn method of age determination using otoliths, the maximum age of Pacific ocean perch was revised to be 90 years (Chilton and Beamish 1982). The updated understanding concerning Pacific ocean perch longevity reduced the estimate of natural mortality based on Hoenig's (1983) relationship to 0.059yr⁻¹. The previous assessment applied a prior distribution on natural mortality based upon multiple life history correlates (including Hoenig's method, Gunderson gonadosomatic index (1997), and McCoy and Gillooly's (2008) theoretical relationship) developed separately for female and male Pacific ocean perch. This assessment also applied a prior on natural mortality. However, the prior and standard deviation were generated as a non-linear function of maximum age as developed by Then et al. (2015) and modified by Owen Hamel which greatly improved the fit to the underlying age data to create the 'Hamel-Then' prior. A maximum age of 100 was used in the development of the prior where female natural mortality was set equal to 0.054 and male natural mortality estimated as an offset from females at 0.054.

Sex ratio, maturation, and fecundity

Examining all biological data sources, the sex ratio of young fish are within 5% of 1:1 by either length or age (Figure 12 and 13), and hence this assessment the sex ratio at birth was assumed to be 1:1. This assessment assumed a logistic maturity-at-length curve based on analysis of 537 fish maturity samples collected from the NWFSC shelf-slope survey. This is revised from the previous assessment which assumed maturity-at-age based on the work of Hannah and Parker (Hannah and Parker 2007). Additionally, the new maturity-at-length curve is based on the estimate of functional maturity an approach that classifies rockfish maturity with developing oocytes as mature or immature based on the proportion of vitellogenin in the cytoplasm and the measured frequency of atretic cells (M. Head, personal communication). The 50% size-at-maturity was estimated at 32.1 cm with maturity asymptoting to one for larger fish.

The fecundity-at-age has also been updated from the previous assessment based on new

research. Dick (2017) estimated new fecundity relationships for select West Coast stocks where fecundity for Pacific ocean perch was esimated equal to $0L^{4.98}$ in millions of eggs. Spawning output at length is shown in Figure 14.

Length-weight relationship

The length-weight relationship for Pacific ocean perch was esimated outside the model using all biological data available from fishery and fishery-independant data sources where the female weight-at-length in grams was estimated at $0.0000098L^{3.11}$ and males at $0.0000094L^{3.12}$ where L is length in cm.

Growth (length-at-age)

Write if estiamted or fixed in the final model

Aging Precision And Bias

2.1.6 Environmental Or Ecosystem Data Included In The Assessment environmental-or-ecosystem-data-included-in-the-assessment

2.2 History Of Modeling Approaches Used For This Stock history-of-modeling-approaches-used-for-this-stock

2.2.1 Previous Assessments previous-assessments

2.2.2 Previous Assessment Recommendations previous-assessment-recommendations

Include: Response to STAR panel recommendations from the most recent previous assessment.

Recommendation 1: blah blah blah.

STAT response: blah blah blah....

Recommendation 2: blah blah blah.

STAT response: blah blah blah....

Recommendation 3: blah blah blah., etc.

STAT response: Continue recommendations as needed

498 **2.3 Model Description** model-description

499 **2.3.1 Transition To The Current Stock Assessment** transition-to-the-current-stock-assessment

500 Include: Complete description of any new modeling approaches

501 Below, we describe the most important changes made since the last full assessment and
502 explain rationale for each change.:

503 1. Change No. 1. *Rationale*: blah blah blah.

504 2. Change No. 2. *Rationale*: blah blah blah.

505 3. Change No. 3. *Rationale*: Continue list as needed.

506 **2.3.2 Definition of Fleets and Areas** definition-of-fleets-and-areas

507 We generated data sources for each of the models. Fleets by model include:

508 *Commercial*: The commercial fleets include...

509 *Recreational*: The recreational fleets include...

510 *Research*: Research derived-data include...

511 **2.3.3 Summary of Data for Fleets and Areas** summary-of-data-for-fleets-and-areas

512 **2.3.4 Modeling Software** modeling-software

513 The STAT team used Stock Synthesis version 3.30.01.13 by Dr. Richard Methot at the
514 NWFSC (Methot and Wetzel [2013](#)). This most recent version was used, since it included
515 improvements and corrections to older versions.

516 **2.3.5 Data Weighting** data-weighting

517 Citation for Francis method (Francis and Hilborn [2011](#))

518 Citation for Ianelli-McAllister harmonic mean method (McAllister and Ianelli [1997](#))

519 2.3.6 Priors

priors

520 Citation for Hamel prior on natural mortality (Hamel 2015)

521 2.3.7 General Model Specifications

general-model-specifications

522 Citation for posterior predictive fecundity relationship from Dick (2009) and (2017)

523 Model data, control, starter, and forecast files can be found in Appendices A-D.

524 2.3.8 Estimated And Fixed Parameters

estimated-and-fixed-parameters

525 A full list of all estimated and fixed parameters is provided in Tables... Estimated and fixed
526 parameters tables currently read in from .csv file, EXAMPLE: Table ??

527 2.4 Model Selection and Evaluation

model-selection-and-evaluation

528 2.4.1 Key Assumptions and Structural Choices

key-assumptions-and-structural-choices

529 Include: Evidence of search for balance between model realism and parsimony.

530 Comparison of key model assumptions, include comparisons based on nested models (e.g.,
531 asymptotic vs. domed selectivities, constant vs. time-varying selectivities).

532 2.4.2 Alternate Models Considered

alternate-models-considered

533 Include: Summary of alternate model configurations that were tried but rejected.

534 2.4.3 Convergence

convergence

535 Include: Randomization run results or other evidence of search for global best estimates.

536 Convergence testing through use of dispersed starting values often requires extreme values to
537 actually explore new areas of the multivariate likelihood surface. Jitter is a SS option that
538 generates random starting values from a normal distribution logistically transformed into
539 each parameter's range (Methot and Wetzel 2013). Table 18 shows the results of running 100
540 jitters for each pre-STAR base model...

541 **2.5 Response To The Current STAR Panel Requests**
response-to-the-current-star-panel-requests

542 **Request No. 1: Add after STAR panel.**

543

544 **Rationale:** Add after STAR panel.

545 **STAT Response:** Add after STAR panel.

546 **Request No. 2: Add after STAR panel.**

547

548 **Rationale:** Add after STAR panel.

549 **STAT Response:** Add after STAR panel.

550 **Request No. 3: Add after STAR panel.**

551

552 **Rationale:** Add after STAR panel.

553 **STAT Response:** Add after STAR panel.

554 **Request No. 4: Example of a request that may have a list:**

555

- 556 **• Item No. 1**
557 **• Item No. 2**
558 **• Item No. 3, etc.**

559 **Rationale:** Add after STAR panel.

560 **STAT Response:** Continue requests as needed.

561 **2.6 Model 1**
model-1

562 **2.6.1 Model 1 Base Case Results**
model-1-base-case-results

563 Table ??

564 **2.6.2 Model 1 Uncertainty and Sensitivity Analyses**
model-1-uncertainty-and-sensitivity-analyses

565 Table 19

566 **2.6.3 Model 1 Retrospective Analysis** model-1-retrospective-analysis

567 **2.6.4 Model 1 Likelihood Profiles** model-1-likelihood-profiles

568 **2.6.5 Model 1 Harvest Control Rules (CPS only)** model-1-harvest-control-rules-cps-only

569 **2.6.6 Model 1 Reference Points (groundfish only)** model-1-reference-points-groundfish-only

570 Intro sentence or two. . . (Table [20](#)).

571 Equilibrium yield at the proxy F_{MSY} harvest rate corresponding to $SPR_{50\%}$ is 1018 mt. Table
572 [e](#) shows the full suite of estimated reference points for the northern area model and Figure [g](#)
573 shows the equilibrium yield curve.

574 **3 Harvest Projections and Decision Tables** harvest-projections-and-decision-tables

575 Table [f](#)

576 **Model 1 Projections and Decision Table (groundfish only)** (Table [21](#)

577 Table [h](#)

578 **Model 2 Projections and Decision Table (groundfish only)**

579 **Model 3 Projections and Decision Table (groundfish only)**

580 **4 Regional Management Considerations** regional-management-considerations

- 581 1. For stocks where current practice is to allocate harvests by management area, a
582 recommended method of allocating harvests based on the distribution of biomass should
583 be provided. The MT advisor should be consulted on the appropriate management
584 areas for each stock.
- 585 2. Discuss whether a regional management approach makes sense for the species from a
586 biological perspective.
- 587 3. If there are insufficient data to analyze a regional management approach, what are the
588 research and data needs to answer this question?

5 Research Needs

research-needs

1. Research need No. 1

2. Research need No. 2

3. Research need No. 3

4. etc.

6 Acknowledgments

acknowledgments

Include: STAR panel members and affiliations as well as names and affiliations of persons who contributed data, advice or information but were not part of the assessment team. Not required in draft assessment undergoing review.

Table 1: West Coast history of regulations.

tab:Regs

Date	Area	Regulation
11/10/1983	Columbia	Closed Columbia area to Pacific ocean perch fishing until the end of the year, as 950 mt OY for this species has been reached;
11/10/1983	Vancouver	retained 5,000-pound trip limit or 10% of total trip weight on landings of Pacific ocean perch in the Vancouver area.
1/1/1984	ALL	Continued 5,000-pound trip limit or 10% of total trip weight on Pacific ocean perch as specified in FMP. Fishery to close when area OYs are reached (see action effective November 10, 1983 above).
8/1/1984	Vancouver Columbia	Reduced trip limit for Pacific ocean perch in the Vancouver and Columbia areas to 20% by weight of all fish on board, not to exceed 5,000 pounds per vessel per trip.
8/16/1984	Columbia	Commercial fishing for Pacific ocean perch in the Columbia area closed for remainder of the year.
1/10/1985	Vancouver Columbia	Established Vancouver and Columbia areas Pacific ocean perch trip limit of 20% by weight of all fish on board (no 5,000-pound limit as specified in last half of 1984).
4/28/1985	Vancouver Columbia	Reduced the Vancouver and Columbia areas Pacific ocean perch trip limit to 5,000 pounds or 20% by weight of all fish on board, whichever is less.
4/28/1985	ALL	Landings of Pacific ocean perch less than 1,000 pounds will be unrestricted. The fishery for this species will close when the OY in each area is reached.
6/10/1985	ALL	Landings of Pacific ocean perch up to 1,000 pounds per trip will be unrestricted regardless of the percentage of these fish on board.
1/1/1986	Cape Blanco North	Established the Pacific ocean perch trip limit north of Cape Blanco (4250) at 20% (by weight) of all fish on board or 10,000 pounds whichever is less;
1/1/1986	ALL	landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board; Vancouver area OY = 600 mt; Columbia area OY = 950 mt.
12/1/1986	Vancouver	OY quota for Pacific ocean perch reached in the Vancouver area; fishery closed until January 1, 1987.
1/1/1987	ALL	Established coastwide Pacific ocean perch limit at 20% of all legal fish on board or 5,000 pounds whichever is less (in round weight); landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board; Vancouver area OY = 500 mt; Columbia area OY = 800 mt.
1/1/1988	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 5,000 pounds, whichever is less; landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board;
1/1/1989	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 5,000 pounds whichever is less;
1/1/1989	ALL	landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board (Vancouver area OY = 500 mt; Columbia area OY = 800 mt).
7/26/1989	ALL	Reduced the coastwide trip limit for Pacific ocean perch to 2,000 pounds or 20% of all fish on board, whichever is less, with no trip frequency restriction.
12/13/1989	Columbia	Closed the Pacific ocean perch fishery in the Columbia area because 1,040 mt OY reached.
1/1/1990	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all fish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board. (Vancouver area OY = 500 mt; Columbia area OY = 1,040 mt).
1/1/1991	ALL	Established the coastwide Pacific ocean perch trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,000 mt).
1/1/1992	ALL	For Pacific ocean perch, established the coastwide trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch be unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,550 mt).

Date	Area	Regulation
1/1/1993	Cape Mendocino Coos Bay	For Pacific ocean perch, continued the coastwide trip limit at 20% (by weight) of all groundfish on board or 3,000 pounds whichever is less; landings of Pacific ocean perch unrestricted if less than 1,000 pounds regardless of percentage on board (harvest guideline for combined Vancouver and Columbia areas = 1,550 mt).
1/1/1994	ALL	Pacific Ocean Perch trip limit of 3,000 pounds or 20% of all fish on board, whichever is less, in landings of Pacific ocean perch above 1,000 pounds.
1/1/1995	ALL	For Pacific Ocean Perch, established a cumulative trip limit of 6,000 pounds per month
1/1/1996	ALL	Pacific Ocean Perch cumulative trip limit of 10,000 pounds per two-month period.
7/1/1996	4030 North	Reduced the cumulative 2-month limit for Pacific ocean perch to 8,000 pounds, and established the cumulative 2-month limit for Dover sole north of Cape Mendocino at 38,000 pounds
1/1/1997	ALL	Pacific Ocean Perch limited entry fishery cumulative trip limit of 8,000 pounds per two-month period
1/1/1998	ALL	Pacific Ocean Perch: limited entry fishery Cumulative trip limit of 8,000 pounds per two-month period.
7/1/1998	ALL	Open Access Rockfish: removed overall rockfish monthly limit and replaced it with limits for component rockfish species: for Sebastes complex, monthly cumulative limit is 33,000 pounds, for widow rockfish, monthly cumulative trip limit is 3,000 pounds, for Pacific Ocean Perch, monthly cumulative trip limit is 4,000 pounds.
1/1/1999	ALL	for the limited entry fishery A new three phase cumulative limit period system is introduced for 1999. Phase 1 is a single cumulative limit period that is 3months long, from January 1 - March 31. Phase 2 has 3 separate 2 month cumulative limit periods of April 1 - May 31, June 1 - July 31, and August 1 - September 30. Phase 3 has 3 separate 1 month cumulative limit periods of October 1-31, November 1-30, and December 1-31. For all species except Pacific ocean perch and Bocaccio, there will be no monthly limit within the cumulative landings limit periods. An option to apply cumulative trip limits lagged by 2 weeks (from the 16th to the 15th) was made available to limited entry trawl vessels when their permits were renewed for 1999. Vessels that are authorized to operate in this "B" platoon may take and retain, but may not land, groundfish during January 1-15, 1999.
1/1/1999	ALL	for the limited entry fishery Pacific Ocean Perch: cumulative limit, Phase 1: 4,000 pounds per month; Phase 2: 4,000 pounds per month; Phase 3: 4,000pounds per month.
1/1/1999	ALL	for open access gear: Pacific Ocean Perch: coastwide, 100 pounds per month.
1/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 500 lbs per month
1/1/2000	ALL	Pacific Ocean Perch, Open Access gear except exempted trawl, 100 lbs per month
1/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 500 lbs per month
5/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 2500 lbs per 2 months
5/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 2500 lbs per month
11/1/2000	ALL	Limited entry trawl, Pacific Ocean Perch, 500 lbs per month
11/1/2000	ALL	Pacific Ocean Perch, limited entry fixed gear, 500 lbs per month
1/1/2001	3600 North	Pacific Ocean Perch, open access, 100 lbs per month
1/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 1500 lbs per month
1/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 1500 lbs per month
5/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 2500 lbs per month
5/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 2500 lbs per month
10/1/2001	4010 North	Pacific Ocean Perch, limited entry trawl, 1500 lbs per month
11/1/2001	ALL	Pacific Ocean Perch, limited entry fixed gear, 1500 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, open access, 100 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 2000 lbs per month
1/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 2000 lbs per month
4/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 4000 lbs per month
5/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 4000 lbs per month
11/1/2002	4010 North	Pacific Ocean Perch, limited entry fixed gear, 2000 lbs per month
11/1/2002	4010 North	Pacific Ocean Perch, limited entry trawl, 2000 lbs per month
1/1/2003	3800 South	minor slope rockfish south including pacific ocean perch, open access gear, 10000 lbs per 2 months

Date	Area	Regulation
1/1/2003	3800 South	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 30000 lbs per 2 months
1/1/2003	3800 South	Minor slope rockfish south including Pacific ocean perch , limited entry trawl, 30000 lbs per 2 months
1/1/2003	3800 4010	minor slope rockfish south including pacific ocean perch, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch , limited entry trawl, 1800 lbs per 2 months
1/1/2003	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2003	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2003	4010 North	Pacific Ocean Perch, Limited entry trawl gear, 3000 lbs per 2 months
3/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, no more than 25% of the weight of sablefish landed per trip
11/1/2003	3800 4010	Minor slope rockfish south including Pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2004	3800 South	Minor slope rockfish south including Pacific ocean perch, open access gear, 10000 lbs per 2 months
1/1/2004	3800 South	minor slope rockfish south inclding pacific ocean perch, limited entry fixed gear, 40000 lbs per 2 months
1/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 40000 lbs per 2 months
1/1/2004	3800 4010	Minor slope rockfish south including Pacific ocean perch, open access gear, per trip no more than 25% of the weight of sablefish landed
1/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 7000 lbs per 2 months
1/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 7000 lbs per 2 months
1/1/2004	4010 North	pacific ocean perch, open access gear, 100 lbs per month
1/1/2004	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2004	4010 North	pacific ocean perch, limited entry trawl, 3000 lbs per 2 months
5/1/2004	3800 South	minor slope rockfish south inclding pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
5/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
5/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
5/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
11/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 50000 lbs per 2 months
11/1/2004	3800 South	minor slope rockfish south including pacific ocean perch, limited entry trawl, 50000 lbs per 2 months
11/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry fixed gear, 10000 lbs per 2 months
11/1/2004	3800 4010	minor slope rockfish south including pacific ocean perch, limited entry trawl, 10000 lbs per 2 months
1/1/2005	3800 South	minor slope rockfish south including darkblotched and pacific ocean perch, open access gear, 10000 lbs per 2 months
1/1/2005	3800 South	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, closed
1/1/2005	3800 4010	minor slope rockfish south including darkblotched and pacific ocean perch, open access gear, per trip no more than 25% of weight of sablefish onboard
1/1/2005	3800 4010	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, 4000 lbs per 2 months
1/1/2005	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2005	4010 North	pacific ocean perch, limited entry trawl gear, 3000 lbs per 2 months
1/1/2005	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2005	4010 South	minor slope rockfish south including darkblotched and pacific ocean perch, limited entry fixed gear, 40000 lbs per 2 months
5/1/2005	3800 4010	minor slope rockfish south including darkblotched rockfish and pacific ocean perch, limited entry trawl, 8000 lbs per 2 months

Date	Area	Regulation
1/1/2008	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2008	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
1/1/2009	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2009	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2009	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2009	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2009	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 55000 lbs per 2 months
1/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2009	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
7/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 10000 lbs per 2 months
11/1/2009	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2010	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2010	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2010	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2010	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2010	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2010	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 55000 lbs per 2 months
1/1/2010	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, limited entry trawl, 15000 lbs per 2 months
1/1/2010	4010 North	pacific ocean perch, limited entry trawl, 1500 lbs per 2 months
1/1/2011	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2011	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2011	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2011	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2011	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2011	ALL	Pacific Ocean Perch managed in part by IFQ
1/1/2012	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2012	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months
1/1/2012	3800 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months
1/1/2012	3800 4010	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, per trip no more than 25% (by weight) of sablefish landed
1/1/2012	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2013	4010 North	pacific ocean perch, open access gears, 100 lbs per month
1/1/2013	4010 North	pacific ocean perch, limited entry fixed gear, 1800 lbs per 2 months
1/1/2013	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched, limited entry fixed gear, 40000 lbs per 2 months no more than 1375 lbs may be blackgill
1/1/2013	4010 South	minor slope rockfish south including pacific ocean perch and darkblotched rockfish, open access gear, 10000 lbs per 2 months no more than 475 lbs of which may be blackgill rockfish
1/1/2014	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2014	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1375 lbs may be blackgill rockfish

Date	Area	Regulation
1/1/2014	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2014	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
1/1/2015	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2015	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1375 lbs may be blackgill rockfish
1/1/2015	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2015	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
7/1/2015	4010 South	non-trawl, limited entry, minor slope rockfish and darkblotched rockfish and pacific ocean perch, 40000 lbs per 2 months of which no more than 1600 lbs may be blackgill rockfish
7/1/2015	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 550 lbs may be blackgill rockfish
1/1/2016	4010 North	non-trawl, limited entry, pacific ocean perch, 1800 lbs per 2 months
1/1/2016	4010 North	non-trawl, open access, pacific ocean perch, 100 lbs per month
1/1/2016	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 475 lbs may be blackgill rockfish
7/1/2016	4010 South	non-trawl, open access, minor slope rockfish including darkblotched rockfish and pacific ocean perch, 10000 lbs per 2 months of which no more than 550 lbs may be blackgill rockfish

Table 2: Landings for each state (all gears combined), the At-Sea Hake fishery, the Foreign fleet, and research.

Year	California	Oregon	Washington	At-Sea Hake	Foreign	tab:Comm_Catch
						Research
1892	0.0	0.1	0.0	0.0	0	0.0
1893	0.0	0.1	0.0	0.0	0	0.0
1894	0.0	0.1	0.0	0.0	0	0.0
1895	0.0	0.0	0.0	0.0	0	0.0
1896	0.0	0.0	0.0	0.0	0	0.0
1897	0.0	0.0	0.0	0.0	0	0.0
1898	0.0	0.0	0.0	0.0	0	0.0
1899	0.0	0.0	0.0	0.0	0	0.0
1900	0.0	0.0	0.0	0.0	0	0.0
1901	0.0	0.0	0.0	0.0	0	0.0
1902	0.0	0.0	0.0	0.0	0	0.0
1903	0.0	0.0	0.0	0.0	0	0.0
1904	0.0	0.0	0.0	0.0	0	0.0
1905	0.0	0.0	0.0	0.0	0	0.0
1906	0.0	0.0	0.0	0.0	0	0.0
1907	0.0	0.0	0.0	0.0	0	0.0
1908	0.0	0.0	0.1	0.0	0	0.0
1909	0.0	0.0	0.1	0.0	0	0.0
1910	0.0	0.0	0.1	0.0	0	0.0
1911	0.0	0.0	0.1	0.0	0	0.0
1912	0.0	0.0	0.0	0.0	0	0.0
1913	0.0	0.0	0.0	0.0	0	0.0
1914	0.0	0.0	0.0	0.0	0	0.0
1915	0.0	0.0	0.0	0.0	0	0.0
1916	0.1	0.0	0.4	0.0	0	0.0
1917	0.1	0.0	0.8	0.0	0	0.0
1918	0.1	0.0	1.1	0.0	0	0.0
1919	0.1	0.0	0.4	0.0	0	0.0
1920	0.1	0.0	0.3	0.0	0	0.0
1921	0.1	0.0	0.3	0.0	0	0.0
1922	0.1	0.0	0.1	0.0	0	0.0
1923	0.1	0.0	0.2	0.0	0	0.0
1924	0.1	0.0	0.5	0.0	0	0.0
1925	0.1	0.0	0.6	0.0	0	0.0
1926	0.2	0.0	1.0	0.0	0	0.0
1927	0.1	0.0	1.4	0.0	0	0.0
1928	0.1	0.1	1.2	0.0	0	0.0
1929	0.5	0.1	0.7	0.0	0	0.0
1930	0.4	0.1	0.9	0.0	0	0.0
1931	0.9	0.1	0.4	0.0	0	0.0

Year	California	Oregon	Washington	At-Sea Hake	Foreign	Research
1932	0.6	0.1	0.4	0.0	0	0.0
1933	1.1	0.1	0.5	0.0	0	0.0
1934	0.8	0.0	2.3	0.0	0	0.0
1935	0.7	0.1	7.7	0.0	0	0.0
1936	0.4	0.2	1.6	0.0	0	0.0
1937	0.9	0.4	2.0	0.0	0	0.0
1938	1.2	0.1	5.1	0.0	0	0.0
1939	1.9	0.4	8.7	0.0	0	0.0
1940	1.7	9.1	12.2	0.0	0	0.0
1941	2.6	14.0	13.6	0.0	0	0.0
1942	0.9	26.6	18.6	0.0	0	0.0
1943	2.0	94.3	453.6	0.0	0	0.0
1944	5.6	164.5	739.3	0.0	0	0.0
1945	13.4	247.1	1887.1	0.0	0	0.0
1946	14.6	193.2	845.9	0.0	0	0.0
1947	5.1	167.2	385.3	0.0	0	0.0
1948	7.9	177.8	491.1	0.0	0	0.0
1949	4.0	472.9	409.5	0.0	0	0.0
1950	3.0	690.1	675.7	0.0	0	0.0
1951	4.3	840.1	735.1	0.0	0	0.0
1952	2.9	2030.5	305.6	0.0	0	0.0
1953	145.6	1223.5	361.6	0.0	0	0.0
1954	123.2	1837.5	538.8	0.0	0	0.0
1955	48.8	1346.4	555.6	0.0	0	0.0
1956	3.8	2563.8	548.2	0.0	0	0.0
1957	1.6	2128.1	538.5	0.0	0	0.0
1958	2.9	1564.9	530.4	0.0	0	0.0
1959	1.5	892.6	337.0	0.0	0	0.0
1960	19.6	1358.8	928.1	0.0	0	0.0
1961	1.1	2061.9	1179.8	0.0	0	0.0
1962	0.6	2584.9	1725.2	0.0	0	0.0
1963	32.5	3693.9	2006.0	0.0	0	0.0
1964	46.1	4261.6	1770.7	0.0	0	0.0
1965	34.9	5627.8	1972.1	0.0	0	0.0
1966	5.2	1591.2	1725.5	0.0	15561	0.0
1967	17.8	354.7	1861.0	0.0	12357	0.0
1968	21.9	466.4	2501.2	0.0	6639	0.0
1969	8.4	422.3	1236.0	0.0	469	0.0
1970	8.7	507.4	1293.3	0.0	441	0.0
1971	12.2	290.4	673.6	0.0	902	0.0
1972	11.4	105.3	796.5	0.0	950	0.0
1973	11.9	121.2	713.1	0.0	1773	0.0
1974	15.7	136.7	641.8	0.0	1457	0.0
1975	11.4	181.3	413.9	62.3	496	0.0
1976	17.1	663.7	521.1 ₄₂	31.9	239	0.0

Year	California	Oregon	Washington	At-Sea Hake	Foreign	Research
1977	16.7	457.1	752.0	3.8	0	11.9
1978	42.5	498.7	1391.5	15.4	0	0.0
1979	136.7	735.9	581.4	15.1	0	34.5
1980	19.2	948.6	666.2	47.0	0	4.6
1981	10.8	929.7	390.3	15.4	0	0.0
1982	145.9	584.0	273.0	28.3	0	0.0
1983	102.0	1032.7	437.7	10.9	0	4.4
1984	47.6	750.4	815.7	2.3	0	0.9
1985	70.9	789.5	503.2	11.4	0	13.6
1986	52.8	676.5	588.9	19.8	0	1.4
1987	120.9	550.0	399.4	5.4	0	0.0
1988	75.4	749.8	509.8	4.5	0	0.5
1989	29.5	927.8	466.2	4.3	0	4.2
1990	18.3	567.8	427.2	80.9	0	0.0
1991	8.4	853.2	530.1	46.1	0	0.0
1992	15.3	623.8	435.2	373.3	0	4.9
1993	11.0	797.8	464.7	0.9	0	0.2
1994	6.7	626.4	352.0	83.8	0	0.0
1995	9.2	515.0	289.8	46.6	0	2.8
1996	18.4	531.1	236.7	6.3	0	1.2
1997	15.8	439.1	184.9	6.4	0	0.1
1998	21.6	436.6	172.4	22.3	0	3.8
1999	19.8	326.8	145.8	16.5	0	1.4
2000	6.8	95.1	33.0	10.1	0	0.6
2001	0.5	193.4	51.8	21.0	0	2.8
2002	0.8	107.1	39.5	3.9	0	0.3
2003	0.2	94.6	30.2	6.3	0	3.6
2004	2.1	97.7	22.3	1.1	0	2.5
2005	0.1	51.2	10.4	1.7	0	1.8
2006	0.2	52.2	15.8	3.1	0	1.2
2007	0.2	83.6	45.1	4.0	0	0.6
2008	0.4	58.6	16.6	15.9	0	0.8
2009	0.9	58.7	33.2	1.6	0	2.7
2010	0.1	58.0	22.3	16.9	0	1.7
2011	0.1	30.3	19.7	9.2	0	1.9
2012	0.2	30.4	21.8	4.5	0	1.6
2013	0.1	34.9	14.8	5.4	0	1.7
2014	0.2	33.9	15.8	3.9	0	0.6
2015	0.1	38.1	11.4	8.7	0	1.6
2016	0.2	34.1	13.1	10.3	0	0.1

Table 3: Summary of discard rates used in the model by each data source.

				tab:Discard
Year	Source	Discard	Standard Error	
1985	Pikitch	0.027	0.068	
1986	Pikitch	0.024	0.063	
1987	Pikitch	0.039	0.083	
1992	Management	0.100	0.300	
	Restrictions			
2002	WCGOP	0.150	0.164	
2003	WCGOP	0.183	0.268	
2004	WCGOP	0.203	0.206	
2005	WCGOP	0.175	0.346	
2006	WCGOP	0.148	0.243	
2007	WCGOP	0.171	0.261	
2008	WCGOP	0.362	0.172	
2009	WCGOP	0.504	0.153	
2010	WCGOP	0.487	0.195	
2011	WCGOP	0.015	0.053	
2012	WCGOP	0.028	0.054	
2013	WCGOP	0.027	0.054	
2014	WCGOP	0.035	0.050	
2015	WCGOP	0.010	0.053	

Table 4: Summary of commercial fishery length samples used in the stock assessment.

tab:Comm_Lengths			
Year	Trips	Fish	Sample Size
1966	1	238	7
1967	5	1020	35
1968	3	912	21
1969	4	1213	28
1970	13	1830	92
1971	22	4698	155
1972	23	4561	162
1973	17	4134	120
1974	20	4806	141
1975	19	3637	134
1976	21	3677	148
1977	32	4846	226
1978	52	7715	367
1979	34	3414	240
1980	55	5426	388
1981	40	3921	282
1982	48	4824	339
1983	39	3944	275
1984	31	3103	219
1985	45	4509	318
1986	40	4005	282
1987	43	3056	304
1988	9	602	64
1989	16	798	113
1990	12	599	85
1991	8	216	38
1994	43	2608	304
1995	49	3161	346
1996	64	3085	452
1997	76	3570	537
1998	56	3450	395
1999	58	2812	409
2000	49	2004	326
2001	59	1696	293
2002	50	1666	280

Year	Trips	Fish	Sample Size
2003	68	1685	301
2004	53	1202	219
2005	50	1270	225
2006	59	1486	264
2007	81	2248	391
2008	101	3058	523
2009	108	3208	551
2010	131	2829	521
2011	100	1944	368
2012	97	1873	355
2013	117	2168	416
2014	140	2850	533
2015	107	2459	446
2016	92	1271	267

Table 5: Summary of Pacific ocean perch survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1979	125	2375	303
1985	126	2558	306

tab:POP_Lengths

Table 6: Summary of Triennial survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1980	18	1315	43
1983	40	2820	97
1986	17	877	41
1989	42	1851	102
1992	33	1182	80
1995	71	1136	172
1998	81	1482	196
2001	74	669	179
2004	63	1240	153

tab:TriennialLengths

Table 7: Summary of AFSC slope survey length samples used in the stock assessment.

Year	Tows	Fish	Sample Size
1996	48	1396	116
1997	21	347	51
1999	21	562	51
2000	19	353	46
2001	23	390	55

tab:AFSC_Lengths

Table 8: Summary of NWFSC slope survey length samples used in the stock assessment.

tab:NWslope_Lengths			
Year	Tows	Fish	Sample Size
2001	18	27	43
2002	24	54	58

Table 9: Summary of NWFSC shelf-slope survey length samples used in the stock assessment.

tab:NWcombo_Lengths			
Year	Tows	Fish	Sample Size
2003	46	80	111
2004	34	56	82
2005	38	81	92
2006	33	73	80
2007	50	74	121
2008	39	75	94
2009	46	61	111
2010	53	73	128
2011	53	72	128
2012	50	79	121
2013	45	76	109
2014	52	77	126
2015	69	67	167

Table 10: Summary of commercial fishery age samples used in the stock assessment.

tab:Comm_Ages			
Year	Trips	Fish	Sample Size
1981	11	1027	78
1982	40	2776	282
1983	33	3320	233
1984	27	2625	191
1985	21	2097	148
1986	17	1696	120
1987	24	1196	169
1988	4	200	28
1994	8	238	41
1999	18	863	127
2000	14	677	99
2001	40	1349	226
2002	38	1414	233
2003	41	1333	225
2004	30	854	148
2005	37	1018	177
2006	49	1259	223
2007	63	1825	315
2008	44	1129	200
2009	76	1549	290
2010	53	1258	227
2011	86	1251	259
2012	7	331	49

Table 11: Summary of Pacific ocean perch survey age samples used in the stock assessment.

tab:POP_Ages			
Year	Tows	Fish	Sample Size
1985	29	1635	70

Table 12: Summary of Triennial survey age samples used in the stock assessment.

tab:Triennial_Ages			
Year	Tows	Fish	Sample Size
1989	15	577	36
1992	10	373	24
1995	12	275	29
1998	28	352	68
2001	43	342	104
2004	57	416	138

Table 13: Summary of NWFSC slope survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2001	17	125	41
2002	24	216	58

Table 14: Summary of NWFSC shelf-slope survey age samples used in the stock assessment.

Year	Tows	Fish	Sample Size
2003	45	265	109
2004	34	149	82
2005	38	192	92
2006	33	170	80
2007	50	228	121
2008	39	218	94
2009	45	190	109
2010	53	292	128
2011	53	258	128

Table 15: Summary of the commercial catch-per-unit effort time-series used in the stock assessment.

Year	Obs	SE
1956	0.40	0.40
1957	0.30	0.40
1958	0.32	0.40
1959	0.29	0.40
1960	0.28	0.40
1961	0.31	0.40
1962	0.29	0.40
1963	0.34	0.40
1964	0.35	0.40
1965	0.55	0.40
1966	0.47	0.40
1967	0.30	0.40
1968	0.17	0.40
1969	0.18	0.40
1970	0.17	0.40
1971	0.20	0.40
1972	0.20	0.40
1973	0.11	0.40

Table 16: Summary of the fishery-independant biomass/abundance time-series used in the stock assessment.

tab:Index_Summary										
Year	POP		Triennial		AFSC Slope		NWFSC Slope		NWFSC Shelf-Slope	
	Obs	SE	Obs	SE	Obs	SE	Obs	SE	Obs	SE
1979	56461	0.27	-	-	-	-	-	-	-	-
1980	-	-	10384	0.56	-	-	-	-	-	-
1983	-	-	8974	0.50	-	-	-	-	-	-
1985	34645	0.29	-	-	-	-	-	-	-	-
1986	-	-	2977	0.57	-	-	-	-	-	-
1989	-	-	4873	0.56	-	-	-	-	-	-
1992	-	-	3207	0.55	-	-	-	-	-	-
1995	-	-	2724	0.54	-	-	-	-	-	-
1996	-	-	-	-	7621	0.51	-	-	-	-
1997	-	-	-	-	3807	0.51	-	-	-	-
1998	-	-	4163	0.55	-	-	-	-	-	-
1999	-	-	-	-	4694	0.50	2201	0.48	-	-
2000	-	-	-	-	4243	0.53	2010	0.50	-	-
2001	-	-	1494	0.55	4187	0.49	2290	0.57	-	-
2002	-	-	-	-	-	-	1646	0.58	-	-
2003	-	-	-	-	-	-	-	-	9940	0.64
2004	-	-	2922	0.58	-	-	-	-	4870	0.68
2005	-	-	-	-	-	-	-	-	7782	0.67
2006	-	-	-	-	-	-	-	-	5722	0.70
2007	-	-	-	-	-	-	-	-	5913	0.64
2008	-	-	-	-	-	-	-	-	3710	0.68
2009	-	-	-	-	-	-	-	-	2754	0.65
2010	-	-	-	-	-	-	-	-	4943	0.63
2011	-	-	-	-	-	-	-	-	7417	0.63
2012	-	-	-	-	-	-	-	-	8326	0.63
2013	-	-	-	-	-	-	-	-	7566	0.63
2014	-	-	-	-	-	-	-	-	4720	0.63
2015	-	-	-	-	-	-	-	-	5317	0.60

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)	NA
1	NatM_p_1_Fem_GP_1	0	-2.000	(0.02, 0.1)			Log_Norm	Log_Norm (-2.9, 0.1)
2	L_at_Amin_Fem_GP_1	21	-3.000	(15, 25)			No_prior	None
3	L_at_Amax_Fem_GP_1	42	2.000	(35, 45)	OK	0	No_prior	None
4	VonBert_K_Fem_GP_1	0	3.000	(0.1, 0.4)	OK	0	No_prior	None
5	CV_young_Fem_GP_1	2	5.000	(0.03, 5)	OK	0	No_prior	None
6	CV_old_Fem_GP_1	3	5.000	(0.03, 5)	OK	0	No_prior	None
7	Wtlen_1_Fem	0	-50.000	(0, 3)			No_prior	None
8	Wtlen_2_Fem	3	-50.000	(2, 4)			No_prior	None
9	Mat50%_Fem	32	-50.000	(20, 40)			No_prior	None
10	Mat_slope_Fem	-1	-50.000	(-2, 4)			No_prior	None
11	Eggs_scalar_Fem	0	-50.000	(0, 6)			No_prior	None
12	Eggs_exp_len_Fem	5	-50.000	(-3, 5)			No_prior	None
13	NatM_p_1_Mal_GP_1	0	2.000	(0, 0.3)	OK	0	Normal	Normal (0.05, 0.1)
14	L_at_Amin_Mal_GP_1	21	-2.000	(6, 68)			No_prior	None
15	L_at_Amax_Mal_GP_1	39	2.000	(13, 122)	OK	0	No_prior	None
16	VonBert_K_Mal_GP_1	0	3.000	(0.04, 1.09)	OK	0	No_prior	None
17	CV_young_Mal_GP_1	2	5.000	(0, 742.07)	OK	0	No_prior	None
18	CV_old_Mal_GP_1	2	5.000	(0, 742.07)	OK	0	No_prior	None
19	Wtlen_1_Mal	0	-50.000	(0, 3)			No_prior	None
20	Wtlen_2_Mal	3	-50.000	(2, 4)			No_prior	None
24	CohortGrowDev	1	-50.000	(0, 2)			No_prior	None
25	FracFemale_GP_1	0	-99.000	(0.000001, 0.999999)			No_prior	None
26	SR_LN(R0)	9	1.000	(5, 20)	OK	0	No_prior	None
27	SR_BH_steep	0	2.000	(0.2, 1)	OK	0	Full_Beta	Full_Beta (0.76, 1)
28	SR_sigmaR	1	-6.000	(0.5, 1.2)			No_prior	None
29	SR_regime	0	-50.000	(-5, 5)			No_prior	None

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)	NA
30	SR_autocorr	0	-50.000	(0, 2)			No.prior	None
154	LnQ_base_Fishery(1)	-12	-1.000	(-15, 15)			No.prior	None
155	LnQ_base_POP(4)	0	-1.000	(-15, 15)			No.prior	None
156	LnQ_base_Triennial(5)	-1	-1.000	(-15, 15)			No.prior	None
157	Q_extraSD_Triennial(5)	0	-2.000	(0, 0.5)			No.prior	None
158	LnQ_base_AFSCSlope(6)	-1	-1.000	(-15, 15)			No.prior	None
159	LnQ_base_NWFSCSlope(7)	-2	-1.000	(-15, 15)			No.prior	None
160	LnQ_base_NWFSCcombo(8)	-1	-1.000	(-15, 15)			No.prior	None
161	Q_extraSD_NWFSCcombo(8)	0	-2.000	(0, 0.5)			No.prior	None
162	SizeSel_P1_Fishery(1)	38	2.000	(20, 45)	OK	0	No.prior	None
163	SizeSel_P2_Fishery(1)	-5	-2.000	(-6, 4)			No.prior	None
164	SizeSel_P3_Fishery(1)	4	3.000	(-1, 9)	OK	0	No.prior	None
165	SizeSel_P4_Fishery(1)	-9	3.000	(-9, 9)	LO	0	No.prior	None
166	SizeSel_P5_Fishery(1)	-4	4.000	(-5, 9)	OK	0	No.prior	None
167	SizeSel_P6_Fishery(1)	1	2.000	(-5, 9)	OK	0	No.prior	None
168	Retain_P1_Fishery(1)	29	1.000	(15, 45)	OK	0	No.prior	None
169	Retain_P2_Fishery(1)	1	1.000	(0.1, 10)	OK	0	No.prior	None
170	Retain_P3_Fishery(1)	1	1.000	(-10, 10)	OK	31441	No.prior	None
171	Retain_P4_Fishery(1)	0	-3.000	(0, 0)			No.prior	None
172	SizeSel_P1_ASHOP(2)	50	2.000	(20, 49.5)	HI	0	No.prior	None
173	SizeSel_P2_ASHOP(2)	-5	-2.000	(-6, 4)			No.prior	None
174	SizeSel_P3_ASHOP(2)	5	3.000	(-1, 9)	OK	0	No.prior	None
175	SizeSel_P4_ASHOP(2)	1	3.000	(-1, 9)	OK	6133	No.prior	None
176	SizeSel_P5_ASHOP(2)	-5	4.000	(-9, 9)	OK	3	No.prior	None
177	SizeSel_P6_ASHOP(2)	999	-2.000	(-5, 999)			No.prior	None
178	SizeSel_P1_POP(4)	24	2.000	(20, 70)	OK	3	No.prior	None

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)	NA
179	SizeSel_P2_POP(4)	13	3.000	(0.001, 50)	OK	5	No_prior	None
180	SizeSel_P1_Triennial(5)	29	2.000	(20, 45)	OK	4	No_prior	None
181	SizeSel_P2_Triennial(5)	-5	-2.000	(-6, 4)			No_prior	None
182	SizeSel_P3_Triennial(5)	4	3.000	(-1, 9)	OK	2	No_prior	None
183	SizeSel_P4_Triennial(5)	2	3.000	(-1, 9)	OK	3	No_prior	None
184	SizeSel_P5_Triennial(5)	-1	4.000	(-5, 9)	OK	1	No_prior	None
185	SizeSel_P6_Triennial(5)	-1	2.000	(-5, 9)	OK	1	No_prior	None
186	SizeSel_P1_AFSCSlope(6)	37	2.000	(20, 45)	OK	4	No_prior	None
187	SizeSel_P2_AFSCSlope(6)	-5	-2.000	(-6, 4)			No_prior	None
188	SizeSel_P3_AFSCSlope(6)	5	3.000	(-1, 9)	OK	1	No_prior	None
189	SizeSel_P4_AFSCSlope(6)	1	3.000	(-1, 9)	OK	6133	No_prior	None
190	SizeSel_P5_AFSCSlope(6)	-9	-4.000	(-9, 9)			No_prior	None
191	SizeSel_P6_AFSCSlope(6)	999	-2.000	(-5, 999)			No_prior	None
192	SizeSel_P1_NWFSCSlope(7)	36	2.000	(20, 45)	OK	3	No_prior	None
193	SizeSel_P2_NWFSCSlope(7)	-5	-2.000	(-6, 4)			No_prior	None
194	SizeSel_P3_NWFSCSlope(7)	2	3.000	(-1, 9)	OK	2	No_prior	None
195	SizeSel_P4_NWFSCSlope(7)	1	3.000	(-1, 9)	OK	6133	No_prior	None
196	SizeSel_P5_NWFSCSlope(7)	-9	-4.000	(-9, 9)			No_prior	None
197	SizeSel_P6_NWFSCSlope(7)	999	-2.000	(-5, 999)			No_prior	None
198	SizeSel_P1_NWFSCCombo(8)	50	2.000	(20, 49.5)	HI	0	No_prior	None
199	SizeSel_P2_NWFSCCombo(8)	-5	-2.000	(-6, 4)			No_prior	None
200	SizeSel_P3_NWFSCCombo(8)	7	3.000	(-1, 9)	OK	1	No_prior	None
201	SizeSel_P4_NWFSCCombo(8)	1	3.000	(-1, 9)	OK	6133	No_prior	None
202	SizeSel_P5_NWFSCCombo(8)	-4	4.000	(-9, 9)	OK	6	No_prior	None
203	SizeSel_P6_NWFSCCombo(8)	999	-2.000	(-5, 999)			No_prior	None
204	Retain_P3_Fishery(1)_BLK1repl_1918	4	1.000	(-10, 10)	OK	0	No_prior	None

Continued on next page

Table 17: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)	NA
205	Retain_P3_Fishery(1)_BLK1repl_1992	2	1.000	(-10, 10)	OK	0	No.prior	None
206	Retain_P3_Fishery(1)_BLK1repl_2002	2	1.000	(-10, 10)	OK	0	No.prior	None
207	Retain_P3_Fishery(1)_BLK1repl_2008	0	1.000	(-10, 10)	OK	0	No.prior	None
208	Retain_P3_Fishery(1)_BLK1repl_2011	6	1.000	(-10, 10)	OK	1	No.prior	None
tab:model_params								

Table 18: Results from 100 jitters from the base model.

Status	Base.Model
Returned to base case	-
Found local minimum	-
Found better solution	-
Error in likelihood	-
Total	100

tab:jitter

Table 20: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs) (mt)	Summary biomass 3+	Relative biomass	Age-0 re- cruits	Estimated total catch (mt)	1-SPR	Exp. rate
1918	122166	5664	121512	1.00	10236	0.0	0.00	0
1919	122168	5664	121510	1.00	10242	1.3	0.00	0
1920	122179	5664	121510	1.00	10248	0.5	0.00	0
1921	122204	5664	121534	1.00	10254	0.4	0.00	0
1922	122240	5664	121571	1.00	10260	0.4	0.00	0
1923	122290	5664	121620	1.00	10267	0.3	0.00	0
1924	122350	5665	121680	1.00	10273	0.3	0.00	0
1925	122420	5665	121750	1.00	10280	0.6	0.00	0
1926	122499	5667	121828	1.00	10288	0.8	0.00	0
1927	122586	5670	121914	1.00	10295	1.2	0.00	0
1928	122678	5673	122005	1.00	10302	1.6	0.00	0
1929	122775	5676	122102	1.00	10310	1.4	0.00	0
1930	122876	5680	122203	1.00	10316	1.4	0.00	0
1931	122980	5684	122306	1.00	10322	1.3	0.00	0
1932	123086	5688	122412	1.00	10327	1.4	0.00	0
1933	123194	5693	122520	1.01	10332	1.1	0.00	0
1934	123302	5698	122628	1.01	10335	1.8	0.00	0
1935	123410	5702	122734	1.01	10339	3.2	0.00	0
1936	123511	5707	122836	1.01	10345	8.8	0.00	0
1937	123618	5712	122943	1.01	10355	2.3	0.00	0
1938	123723	5717	123047	1.01	10375	3.4	0.00	0
1939	123825	5722	123148	1.01	10409	6.5	0.00	0
1940	123923	5726	123243	1.01	10875	11.3	0.01	0
1941	124015	5730	123327	1.01	10948	23.6	0.01	0
1942	124119	5734	123407	1.01	11031	31.1	0.02	0
1943	124241	5736	123524	1.01	11106	47.3	0.21	0
1944	123884	5714	123162	1.01	11138	564.3	0.32	0
1945	123211	5675	122485	1.00	11137	933.1	0.65	0.01
1946	121339	5576	120612	0.98	11100	2204.1	0.38	0.02
1947	120678	5534	119951	0.98	11167	1081.5	0.22	0.01
1948	120584	5520	119857	0.97	11381	572.4	0.26	0
1949	120415	5503	119681	0.97	11790	694.7	0.33	0.01
1950	120089	5479	119337	0.97	12481	909.9	0.47	0.01
1951	119350	5434	118566	0.96	13487	1405.1	0.53	0.01
1952	118522	5381	117686	0.95	14613	1621.5	0.72	0.01
1953	117096	5294	116195	0.93	15066	2401.4	0.58	0.02
1954	116531	5240	115571	0.93	14152	1777.2	0.76	0.02

Table 20: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs) (mt)	Summary biomass 3+	Relative biomass	Age-0 re- cruits	Estimated total catch (mt)	1-SPR	Exp. rate
1955	115420	5152	114456	0.91	12587	2567.2	0.65	0.02
1956	115075	5097	114181	0.90	11086	2004.3	0.90	0.02
1957	113632	4993	112840	0.88	9698	3202.2	0.82	0.03
1958	112659	4923	111961	0.87	8570	2743.1	0.70	0.02
1959	112174	4894	111562	0.86	7862	2157.3	0.47	0.02
1960	112390	4920	111842	0.87	7709	1265.6	0.74	0.01
1961	111257	4901	110746	0.87	8012	2370.1	0.93	0.02
1962	108935	4837	108426	0.85	8173	3330.6	1.10	0.03
1963	105336	4714	104812	0.83	7292	4425.5	1.28	0.04
1964	100155	4511	99640	0.80	5986	5883.1	1.33	0.06
1965	94530	4276	94078	0.75	5263	6237.1	1.48	0.07
1966	87216	3953	86838	0.70	4931	7834.4	1.83	0.09
1967	68782	3090	68446	0.55	4484	18970.4	1.83	0.28
1968	54878	2428	54564	0.43	4645	14651.4	1.79	0.27
1969	46071	2007	45774	0.35	5688	9712.5	1.21	0.21
1970	44850	1952	44523	0.34	7661	2183.3	1.25	0.05
1971	43532	1894	43132	0.33	5563	2301.5	1.16	0.05
1972	42682	1852	42223	0.33	3413	1905.5	1.16	0.05
1973	41904	1808	41579	0.32	2731	1888.6	1.36	0.05
1974	40299	1725	40088	0.30	2843	2643.0	1.30	0.07
1975	38949	1658	38767	0.29	3527	2274.1	0.95	0.06
1976	38580	1646	38384	0.29	2880	1182.6	1.09	0.03
1977	37786	1626	37565	0.29	3565	1507.3	1.00	0.04
1978	37143	1619	36945	0.29	3006	1270.3	1.26	0.03
1979	35687	1572	35463	0.28	3077	1999.8	1.13	0.06
1980	34648	1537	34452	0.27	2773	1533.5	1.21	0.04
1981	33360	1485	33163	0.26	3144	1727.0	1.10	0.05
1982	32382	1445	32193	0.26	4091	1381.6	0.97	0.04
1983	31711	1417	31491	0.25	3567	1058.0	1.23	0.03
1984	30492	1361	30235	0.24	3215	1628.0	1.27	0.05
1985	29293	1302	29065	0.23	3869	1660.2	1.21	0.06
1986	28362	1253	28144	0.22	2577	1422.2	1.21	0.05
1987	27508	1206	27277	0.21	2904	1375.3	1.11	0.05
1988	26951	1172	26776	0.21	3230	1105.7	1.25	0.04
1989	26112	1129	25914	0.20	4297	1377.9	1.31	0.05
1990	25218	1084	24988	0.19	4711	1470.7	1.17	0.06
1991	24738	1057	24455	0.19	2790	1123.2	1.34	0.05

Table 20: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs) (mt)	Summary biomass 3+	Relative biomass	Age-0 re- cruits	Estimated total catch (mt)	1-SPR	Exp. rate
1992	23981	1014	23709	0.18	1549	1477.4	1.38	0.06
1993	23175	965	23015	0.17	1266	1570.5	1.36	0.07
1994	22446	926	22344	0.16	3650	1417.8	1.27	0.06
1995	21893	898	21771	0.16	3145	1180.6	1.17	0.05
1996	21538	887	21313	0.16	1582	954.3	1.12	0.04
1997	21276	883	21097	0.16	1587	881.7	1.00	0.04
1998	21121	886	21017	0.16	1796	717.4	1.01	0.03
1999	20887	883	20765	0.16	7288	724.6	0.87	0.03
2000	20824	882	20605	0.16	10246	564.6	0.33	0.03
2001	21315	900	20803	0.16	4587	160.5	0.55	0.01
2002	22000	911	21430	0.16	3508	295.6	0.36	0.01
2003	23016	926	22738	0.16	2113	179.0	0.32	0.01
2004	24028	939	23814	0.17	5103	157.3	0.30	0.01
2005	25034	954	24851	0.17	2657	147.9	0.16	0.01
2006	26075	988	25783	0.17	2767	77.0	0.17	0
2007	27099	1041	26923	0.18	2883	85.5	0.28	0
2008	28097	1100	27738	0.19	70862	157.9	0.26	0.01
2009	29633	1155	28463	0.20	1947	150.9	0.27	0.01
2010	32534	1203	29078	0.21	3171	168.5	0.25	0.01
2011	37402	1245	37251	0.22	4069	161.1	0.10	0
2012	41377	1288	41155	0.23	3956	61.4	0.09	0
2013	45262	1336	44993	0.24	5973	59.1	0.08	0
2014	48859	1442	48570	0.25	4345	57.8	0.07	0
2015	52065	1656	51700	0.29	5358	55.4	0.07	0
2016	54863	1919	54560	0.34	5888	60.1	0.06	0
2017	57212	2151	56852	0.38	6312	58.3	0.96	0
2018	57520	2257	57128	0.40	6493	-	-	-
2019	57415	2312	57000	0.41	6586	-	-	-
2020	57055	2337	56629	0.41	6627	-	-	-
2021	56548	2342	56117	0.41	6634	-	-	-
2022	55971	2333	55538	0.41	6620	-	-	-
2023	55373	2316	54940	0.41	6592	-	-	-
2024	54783	2296	54351	0.41	6559	-	-	-
2025	54215	2274	53785	0.40	6523	-	-	-
2026	53674	2252	53247	0.40	6486	-	-	-
2027	53166	2230	52740	0.39	6449	-	-	-
2028	52690	2210	52267	0.39	6413	-	-	-

Table 20: Time-series of population estimates from the base model.

Year	Total biomass (mt)	Spawning output (million eggs) (mt)	Summary biomass 3+	Relative biomass	Age-0 re- cruits	Estimated total 1-SPR catch (mt)	Exp. rate
tab:Timeseries_mod1							

Table 19: Sensitivity of the base model to dropping or down-weighting data sources and alternative assumptions about growth.

Label	tab:Sensitivity_model1							
	Base (Francis weights)	Harmonic mean weights	Drop index	Drop ages	Down- weight lengths	Free size Age0	Free CV Amin	External growth
TOTAL_like	-	-	-	-	-	-	-	-
Catch_like	-	-	-	-	-	-	-	-
Equil_catch_like	-	-	-	-	-	-	-	-
Survey_like	-	-	-	-	-	-	-	-
Length_comp_like	-	-	-	-	-	-	-	-
Age_comp_like	-	-	-	-	-	-	-	-
Parm_priors_like	-	-	-	-	-	-	-	-
SSB_Unfished_thousand_mt	-	-	-	-	-	-	-	-
TotBio_Unfished	-	-	-	-	-	-	-	-
SmryBio_Unfished	-	-	-	-	-	-	-	-
Recr_Unfished_billions	-	-	-	-	-	-	-	-
SSB_Btgt_thousand_mt	-	-	-	-	-	-	-	-
SPR_Btgt	-	-	-	-	-	-	-	-
Fstd.Btgt	-	-	-	-	-	-	-	-
TotYield.Btgt_thousand_mt	-	-	-	-	-	-	-	-
SSB_SPRtgt_thousand_mt	-	-	-	-	-	-	-	-
Fstd.SPRtgt	-	-	-	-	-	-	-	-
TotYield_SPRtgt_thousand_mt	-	-	-	-	-	-	-	-
SSB_MSY_thousand_mt	-	-	-	-	-	-	-	-
SPR_MSY	-	-	-	-	-	-	-	-
Fstd.MSY	-	-	-	-	-	-	-	-
TotYield_MSY_thousand_mt	-	-	-	-	-	-	-	-
RetYield_MSY	-	-	-	-	-	-	-	-
Bratio.2015	-	-	-	-	-	-	-	-
F_2015	-	-	-	-	-	-	-	-
SPRratio.2015	-	-	-	-	-	-	-	-
Recr.2015	-	-	-	-	-	-	-	-
Recr_Virgin_billions	-	-	-	-	-	-	-	-
L.at_Amin_Fem_GP_1	-	-	-	-	-	-	-	-
L.at_Amax_Fem_GP_1	-	-	-	-	-	-	-	-
VonBert_K_Fem_GP_1	-	-	-	-	-	-	-	-
CV_young_Fem_GP_1	-	-	-	-	-	-	-	-
CV_old_Fem_GP_1	-	-	-	-	-	-	-	-

Table 21: Projection of potential OFL, spawning biomass, and depletion for the base case model.

Year	OFL contriubtion (mt)	ACL landings (mt)	Age 5+ biomass (mt)	Spawning Output	tab:Forecast_mod1 Depletion
2017	1899	1308	56852	2151	0.38
2018	1988	1393	57128	2257	0.40
2019	2018	1419	57000	2312	0.41
2020	2008	1414	56629	2337	0.41
2021	1975	1395	56117	2342	0.41
2022	1935	1369	55538	2333	0.41
2023	1894	1343	54940	2316	0.41
2024	1858	1319	54351	2296	0.41
2025	1827	1298	53785	2274	0.40
2026	1801	1277	53247	2252	0.40
2027	1779	1258	52740	2230	0.39
2028	1760	1241	52267	2210	0.39




Figure 1: Map showing the state boundary lines for management of the recreational fishing fleets. CRFS Districts 1-6 in California are presented as well as the WDFW Recreational Management Areas in Washington. Florence, OR is shown as a potential location of model stratification. fig:boundary_map




Figure 2: Summary of data sources used in the Base model. fig:data_plot




Figure 3: Discard length frequency distributions from the WCGOP for Pacific ocean perch. fig:WCGOP_dis

length comps, retained, Fishery




Figure continued from previous page




Figure 4: length comps, discard, Fishery fig:mod1_1_comp_lenfit_flt1mkt1




Figure 5: Pearson residuals, discard, Fishery (max=4.71)
 Closed bubbles are positive residuals (observed > expected) and open bubbles are negative residuals (observed < expected).
 fig:mod1_2_comp_lenfit_residsfit1mkt1




Figure 6: N-EffN comparison, length comps, discard, Fishery | fig:mod1_3_comp_lenfit_sam




Figure 7: Francis data weighting method TA1.8 Fishery Suggested sample size adjustment (with 95% interval) for len data from Fishery: 0.8226 (0.5299_1.5611) fig:mod1_4_comp_lenfit_data_weig





Figure 8: length comps, retained, Fishery (plot 1 of 2) | fig:mod1_5_comp_lenfit_flt1mk



602

603

Figure continued from previous page




Figure 9: N_EffN comparison, length comps, retained, Fishery | fig:mod1_8_comp_lenfit_san




Figure 10: Francis data weighting method TA1.8 Fishery Suggested sample size adjustment (with 95% interval) for len data from Fishery: 0.8226 (0.5458-1.505) [fig:mod1_9_comp_lenfit_data_weigh




Figure 11: length comps, whole catch, ASHOP fig:mod1_10_comp_lenfit_flt2mkt0




Figure 12: The estimated sex ratio of Pacific ocean perch at length from all biological data sources. `fig:sexratio`




Figure 13: The estimated sex ratio of Pacific ocean perch at age from all biological data sources. `fig:sexratio_Age`




Figure 14: Fecundity at length of Pacific ocean perch in the Base model. ^{fig:fecundity}

604 \dot{I} — *****MODEL 2 REFERENCE POINTS FIGURES – IF NEEDED *****
605 \dot{I}

References

references

- Bradburn, M., Keller, A., and Horness, B. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: Estimates of distribution, abundance, length, and age composition. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine Fisheries Service.
- Chilton, D.E., and Beamish, R.J. 1982. Age determination methods for fishes studied by the Groundfish Program at the Pacific Biological Station. [Ottawa:] Minister of Supply; Services Canada.
- Dick, E., Beyer, S., Mangel, M., and Ralston, S. 2017. A meta-analysis of fecundity in rockfishes (genus *Sebastes*). Fisheries Research **187**: 73–85. doi: [10.1016/j.fishres.2016.11.009](https://doi.org/10.1016/j.fishres.2016.11.009).
- Dick, E.J. 2009. Modeling the Reproductive Potential of Rockfishes (*Sebastes* Spp.). ProQuest. Available from [http://books.google.com/books?hl=en&lr=&id=0d6-3rhfynkC&oi=fnd&pg=PR7&dq=%22Synthesis+of+findings+regarding+the+reproductive%22+%22C:+Linear+interpolation+algorithms%22+%22for+yellowtail+rockfish+\(S.+flavidus\)%22+%22greater+than+zero,+based+on+the+2-level+relative+fecundity%22+%22A:+Methods+for+data+recovery+from+published%22+%22&ots=NR0UylgymD&sig=58IaN_a3pJeYTPYVmJ1NYMABmvE](http://books.google.com/books?hl=en&lr=&id=0d6-3rhfynkC&oi=fnd&pg=PR7&dq=%22Synthesis+of+findings+regarding+the+reproductive%22+%22C:+Linear+interpolation+algorithms%22+%22for+yellowtail+rockfish+(S.+flavidus)%22+%22greater+than+zero,+based+on+the+2-level+relative+fecundity%22+%22A:+Methods+for+data+recovery+from+published%22+%22&ots=NR0UylgymD&sig=58IaN_a3pJeYTPYVmJ1NYMABmvE) [accessed 27 February 2017].
- Francis, R.C., and Hilborn, R. 2011. Data weighting in statistical fisheries stock assessment models. Canadian Journal of Fisheries and Aquatic Sciences **68**(6): 1124–1138. doi: [10.1139/f2011-025](https://doi.org/10.1139/f2011-025).
- Gunderson, D.R. 1977. Population biology of Pacific ocean perch, *Sebastes alutus*, stocks in the Washington/Queen Charlotte Sound region and their response to fishing. Fishery Bulletin **75**: 369–403. Available from <http://fishbull.noaa.gov/75-2/gunderson.pdf> [accessed 27 February 2017].
- Gunderson, D.R. 1978. Results of cohort analysis for Pacific ocean perch stocks off British Columbia, Washington, and Oregon and an evaluation of alternative rebuilding strategies for these stocks. Pacific Fishery Management Council, 7700 Ambassador Place NE, Suite 200, Portland, OR 97220.
- Gunderson, D.R. 1997. Trade-off between reproductive effort and adult survival in oviparous and viviparous fishes. Canadian Journal of Fisheries and Aquatic Sciences **54**(5): 990–998. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f97-019> [accessed 27 February 2017].
- Gunderson, D.R., and Sample, T.M. 1980. Distribution and abundance of rockfish off Washington, Oregon and California during 1977. Northwest; Alaska Fisheries Center, National

Marine Fisheries Service. Available from <http://spo.nmfs.noaa.gov/mfr423-4/mfr423-42.pdf> [accessed 28 February 2017].

Gunderson, D.R., Westrheim, S., Demory, R., and Fraidenburg, M. 1977. The status of Pacific ocean perch (*Sebastes alutus*) stocks off British Columbia, Washington, and Oregon in 1974.

Hamel, O.S. 2015. A method for calculating a meta-analytical prior for the natural mortality rate using multiple life history correlates. ICES Journal of Marine Science: Journal du Conseil **72**(1): 62–69. doi: [10.1093/icesjms/fsu131](https://doi.org/10.1093/icesjms/fsu131).

Hannah, R., and Parker, S. 2007. Age-modulated variation in reproductive development of female Pacific Ocean perch (*Sebastes alutus*) in waters off Oregon. Alaska Sea Grant, University of Alaska Fairbanks. pp. 1–20. doi: [10.4027/bamnpr.2007.01](https://doi.org/10.4027/bamnpr.2007.01).

Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. Fishery Bulletin **82**: 898–903. Available from <http://fishbull.noaa.gov/81-4/hoenig.pdf> [accessed 28 February 2017].

Karnowski, M., Gertseva, V., and Stephens, A. 2014. Historical Reconstruction of Oregon's Commercial Fisheries Landings. Oregon Department of Fish; Wildlife, Salem, OR.

McAllister, M.K., and Ianelli, J.N. 1997. Bayesian stock assessment using catch-age data and the sampling - importance resampling algorithm. Canadian Journal of Fisheries and Aquatic Sciences **54**: 284–300. Available from <http://www.nrcresearchpress.com/doi/pdf/10.1139/f96-285> [accessed 10 March 2017].

McCoy, M.W., and Gillooly, J.F. 2008. Predicting natural mortality rates of plants and animals. Ecology Letters **11**(7): 710–716. doi: [10.1111/j.1461-0248.2008.01190.x](https://doi.org/10.1111/j.1461-0248.2008.01190.x).

Methot, R.D., and Wetzel, C.R. 2013. Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. Fisheries Research **142**: 86–99. doi: [10.1016/j.fishres.2012.10.012](https://doi.org/10.1016/j.fishres.2012.10.012).

Pikitch, E.K., Erickson, D.L., and Wallace, J.R. 1988. An evaluation of the effectiveness of trip limits as a management tool. Northwest; Alaska Fisheries Center, National Marine Fisheries Service NWAFC Processed Report. Available from <https://www.afsc.noaa.gov/Publications/ProcRpt/PR1988-27.pdf> [accessed 28 February 2017].

Ralston, S., Pearson, D.E., Field, J.C., and Key, M. 2010. Documentation of the California catch reconstruction project. US Department of Commerce, National Oceanic; Atmospheric Administration, National Marine.

Rogers, J. 2003. Species allocation of *Sebastes* and *Sebastolobus* species caught by foreign countries off Washington, Oregon, and California, U.S.A. in 1965-1976. Unpublished

document.

Rogers, J.B., and Pikitch, E.K. 1992. Numerical definition of groundfish assemblages caught off the coasts of Oregon and Washington using commercial fishing strategies. Canadian Journal of Fisheries and Aquatic Sciences **49**(12): 2648–2656. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f92-293> [accessed 9 March 2017].

Seeb, L.W., and Gunderson, D.R. 1988. Genetic variation and population structure of Pacific ocean perch (*Sebastes alutus*). Canadian Journal of Fisheries and Aquatic Sciences **45**(1): 78–88. Available from <http://www.nrcresearchpress.com/doi/abs/10.1139/f88-010> [accessed 28 February 2017].

Then, A.Y., Hoenig, J.M., Hall, N.G., and Hewitt, D.A. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. ICES Journal of Marine Science **72**(1): 82–92. doi: [10.1093/icesjms/fsu136](https://doi.org/10.1093/icesjms/fsu136).

Thorson, J.T., and Barnett, L.A.K. 2017. Comparing estimates of abundance trends and distribution shifts using single- and multispecies models of fishes and biogenic habitat. ICES Journal of Marine Science: Journal du Conseil: fsw193. doi: [10.1093/icesjms/fsw193](https://doi.org/10.1093/icesjms/fsw193).

Weinberg, J.R., Rago, P.J., Wakefield, W.W., and Keith, C. 2002. Estimation of tow distance and spatial heterogeneity using data from inclinometer sensors: An example using a clam survey dredge. Fisheries Research **55**(1–3): 49–61. doi: [10.1016/S0165-7836\(01\)00292-2](https://doi.org/10.1016/S0165-7836(01)00292-2).

Wilkins, M., and Golden, J. 1983. Condition of the Pacific ocean perch resource off Washington and Oregon during 1979: Results of a cooperative trawl survey. North American Journal of Fisheries Management **3**: 103–122.

Withler, R., Beacham, T., Schulze, A., Richards, L., and Miller, K. 2001. Co-existing populations of Pacific ocean perch, *Sebastes alutus*, in Queen Charlotte Sound, British Columbia. Marine Biology **139**(1): 1–12. doi: [10.1007/s002270100560](https://doi.org/10.1007/s002270100560).

Zimmermann, M., Wilkins, M., Weinberg, K., Lauth, R., and Shaw, F. 2003. Influence of improved performance monitoring on the consistency of a bottom trawl survey. ICES Journal of Marine Science **60**(4): 818–826. doi: [10.1016/S1054-3139\(03\)00043-2](https://doi.org/10.1016/S1054-3139(03)00043-2).