

ROMS-Tools: Reproducible Preprocessing and Analysis for Regional Ocean Modeling with ROMS

Nora Loose¹, Tom Nicholas², Scott Eilerman¹, Christopher McBride¹, Sam Maticka¹, Dafydd Stephenson¹, Scott Bachman¹, Pierre Damien³, Ulla Heede¹, Alicia Karspeck¹, Matthew C. Long¹, M. Jeroen Molemaker³, and Abigale Wyatt¹

¹ [C]Worthy LLC, Boulder, CO, United States ² Earthmover PBC ³ University of California, Los Angeles, CA, United States

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a

Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

The ocean regulates Earth's climate and sustains marine ecosystems by circulating and storing heat, carbon, oxygen, and nutrients, while exchanging heat and gases with the atmosphere. Scientists study these processes using ocean models, which simulate the ocean on a grid. **Regional ocean models** focus computational resources on a limited geographical area with fine grid spacing, and can resolve fine-scale phenomena such as mesoscale and submesoscale features, tidal dynamics, coastal currents, upwelling, and detailed biogeochemical (BGC) processes. A widely used regional ocean model is the **Regional Ocean Modeling System (ROMS)** (Shchepetkin & McWilliams, 2005). ROMS has been coupled to the Marine Biogeochemistry Library (MARBL) (Long et al., 2021; Molemaker & contributors, 2025a) to link physical and BGC processes. ROMS-MARBL supports research on environmental management, fisheries, regional climate impacts, and ocean-based carbon dioxide removal (CDR) strategies.

Configuring a regional ocean model like ROMS-MARBL is technically challenging; it requires initialization and time-dependent forcing from oceanic and atmospheric data drawn from multiple external sources in diverse formats. These global datasets can reach several petabytes and must be subsetting, processed, and mapped onto the target domain's geometry, yielding input datasets of 10–100 terabytes for large regional models. Generating these input files is time-consuming, error-prone, and hard to reproduce, creating a bottleneck for both new and experienced users. The Python package **ROMS-Tools** addresses this challenge by providing efficient, dask-backed (Dask Development Team, 2016), user-friendly tools that can be installed via Conda or PyPI and run interactively from Jupyter notebooks. The package supports creating regional grids, preprocessing all required model inputs, and postprocessing and analysis. Current capabilities are fully compatible with UCLA-ROMS (Molemaker & contributors, 2025a, 2025b), with potential support for other ROMS versions, such as Rutgers ROMS (Arango & contributors, 2024), in the future.

Input Data and Preprocessing

ROMS-Tools generates the following input files for ROMS-MARBL:

- Model Grid:** Customizable, curvilinear, and orthogonal grid designed to maintain a nearly uniform horizontal resolution across the domain. The grid is rotatable to align with coastlines and features a terrain-following vertical coordinate.
- Bathymetry:** Derived from **SRTM15** (Tozer et al., 2019).
- Land Mask:** Inferred from coastlines provided by **Natural Earth** or the Global Self-consistent, Hierarchical, High-resolution Geography (**GSHHG**) Database (Vessel &

- Smith, 1996).
4. **Physical Ocean Conditions:** Initial and open boundary conditions for sea surface height, temperature, salinity, and velocities derived from the 1/12° Global Ocean Physics Reanalysis (GLORYS) (Lellouche et al., 2021).
 5. **BGC Ocean Conditions:** Initial and open boundary conditions for dissolved inorganic carbon, alkalinity, and other biogeochemical tracers from Community Earth System Model (CESM) output (Yeager et al., 2022) or hybrid observational-model sources (Garcia et al., 2019; Huang et al., 2022; Lauvset et al., 2016; Yang et al., 2020; Yeager et al., 2022).
 6. **Meteorological forcing:** Wind, radiation, precipitation, and air temperature/humidity processed from the global 1/4° ECMWF Reanalysis v5 (ERA5) (Hersbach et al., 2020) with optional corrections for radiation bias and coastal wind.
 7. **BGC surface forcing:** Partial pressure of carbon dioxide, as well as iron, dust, and nitrogen deposition from CESM output (Yeager et al., 2022) or hybrid observational-model sources (Hamilton et al., 2022; Kok et al., 2021; Landschützer et al., 2016; Yeager et al., 2022).
 8. **Tidal Forcing:** Tidal potential, elevation, and velocities derived from TPXO (Egbert & Erofeeva, 2002) including self-attraction and loading (SAL) corrections.
 9. **River Forcing:** Freshwater runoff derived from Dai & Trenberth (Dai & Trenberth, 2002) or user-provided custom files.
 10. **CDR Forcing:** User-defined interventions that inject BGC tracers at point sources or as larger-scale Gaussian perturbations to simulate CDR interventions. The CDR forcing is prescribed as volume and tracer fluxes (e.g., alkalinity for ocean alkalinity enhancement, iron for iron fertilization, or other BGC constituents). Users can control the magnitude, spatial footprint, and temporal evolution, allowing flexible representation of CDR interventions.
- Some source datasets are accessed automatically by the package, including Natural Earth, Dai & Trenberth runoff, and ERA5 meteorology, while users must manually download SRTM15, GSHHG, GLORYS, the BGC datasets, and TPXO tidal files. Although these are the datasets currently supported, the package's modular design makes it straightforward to add new source datasets in the future.
- To generate the model inputs, ROMS-Tools automates several intermediate processing steps, including:
- **Bathymetry processing:** The bathymetry is smoothed in two stages, first across the entire model domain and then locally in areas with steep slopes, to ensure local steepness ratios do not exceed a prescribed threshold in order to reduce pressure-gradient errors. A minimum depth is enforced to prevent water levels from becoming negative during large tidal excursions.
 - **Mask definition:** The land-sea mask is generated by comparing the ROMS grid's horizontal coordinates with a coastline dataset using the regionmask package (Hauser et al., 2024). Enclosed basins are subsequently filled with land.
 - **Land value handling:** Land values are filled via an algebraic multigrid method using pyamg (Bell et al., 2023) prior to horizontal regridding. This extends ocean values into land areas to resolve discrepancies between source data and ROMS land masks, preventing land-originating values from appearing in ocean cells.
 - **Regridding:** Ocean and atmospheric fields are horizontally and vertically regridded from standard latitude-longitude-depth grids to the model's curvilinear grid with a terrain-following vertical coordinate using xarray (Hoyer & Hamman, 2017) and xgcm (Busecke & contributors, 2025). Velocities are rotated to align with the curvilinear ROMS grid.
 - **Longitude conventions:** ROMS-Tools handles differences in longitude conventions, converting between $[-180^\circ, 180^\circ]$ and $[0^\circ, 360^\circ]$ as needed.
 - **River locations:** Rivers that fall within the model domain are automatically identified and relocated to the nearest coastal grid cell. Rivers that need to be shifted manually or

span multiple cells can be configured by the user.

- **Data streaming:** ERA5 atmospheric data can be accessed directly from the cloud, removing the need for users to pre-download large datasets locally. Similar streaming capabilities may be implemented for other datasets in the future.

Users can quickly design and visualize regional grids and inspect all input fields with built-in plotting utilities. An example of surface initial conditions generated for a California Current System simulation at 5 km horizontal grid spacing is shown in [Figure 1](#).




Figure 1: Surface initial conditions for the California Current System created with ROMS-Tools from GLORYS. Left: potential temperature. Right: zonal velocity. Shown for January 1, 2000.

Postprocessing and Analysis

ROMS-Tools supports postprocessing and analysis of ROMS-MARBL output, including regridding from the native curvilinear, terrain-following grid to a standard latitude-longitude-depth grid using *xesmf* ([Zhuang et al., 2023](#)), with built-in plotting for both grid types. The analysis layer also includes specialized utilities for evaluating carbon dioxide removal (CDR) interventions, such as generating carbon uptake and efficiency curves.

Statement of Need

Setting up a regional ocean model is a major technical undertaking. Traditionally, this work has relied on a patchwork of custom scripts and lab-specific workflows, which can be time-consuming, error-prone, and difficult to reproduce. These challenges slow down science, create a steep barrier to entry for new researchers, and limit collaboration across groups.

Within the ROMS community, the preprocessing landscape has been shaped by tools like *pyroms* ([Hedstrom & contributors, 2023](#)). While providing valuable low-level utilities, *pyroms* presents challenges for new users: installation is cumbersome due to Python/Fortran dependencies, the API is inconsistent, documentation is limited, and it lacks tests, CI, and support for modern Python tools like *xarray* and *dask*. Since active development has largely ceased, its suitability for new projects, such as CDR simulations, is limited.

Tools from other modeling communities cannot simply be adopted, since each ocean model has distinct structural requirements. For example, the *regional-mom6* package ([Barnes et al.,](#)

2024), developed for the Modular Ocean Model v6 (MOM6) (Adcroft et al., 2019), cannot be used to generate ROMS inputs, because ROMS employs a terrain-following vertical coordinate system that requires a specialized vertical regridding approach, whereas MOM6 accepts inputs on arbitrary depth levels and does not require vertical regridding at all. Several other differences further prevent cross-compatibility. Together, these limitations underscored the need for a modern, maintainable, and reproducible tool designed specifically for ROMS.¹

ROMS-Tools was developed to meet this need. It draws on the legacy of the MATLAB preprocessing scripts developed at UCLA (Molemaker, 2024), which encapsulate decades of expertise in configuring regional ocean model inputs. While many of the core algorithms and design principles are retained, ROMS-Tools provides an open-source Python implementation of these MATLAB tools using an object-oriented programming paradigm. This implementation supports a modern workflow with high-level API calls, improving reproducibility, minimizing user errors, and allowing easy extension to new features, forcing datasets, and use cases. In some cases, ROMS-Tools diverges from the MATLAB implementation to take advantage of new methods or better integration with the modern Python ecosystem. By streamlining input generation and analysis, ROMS-Tools reduces technical overhead, lowers the barrier to entry, and enables scientists to focus on research rather than data preparation.

138 Software Design

139 ROMS-Tools is designed to balance **ease of use, flexibility, reproducibility, and scalability** by
140 combining high-level user interfaces with a modular, extensible architecture.

141 Lessons from MATLAB Tools

142 The legacy MATLAB preprocessing scripts were powerful but required users to edit source
143 code directly to configure simulations. This workflow led to frequent errors for new users,
144 made it difficult to track completed steps, and limited reproducibility. ROMS-Tools addresses
145 these issues with **high-level API calls**, automated error-prone steps, and explicit workflow state
146 management via YAML.

147 Design Trade-Offs

148 A central design trade-off in ROMS-Tools is between **user control** and **automation**. Rather than
149 enforcing a fixed workflow, the package exposes key choices such as physical options (e.g.,
150 corrections for radiation or wind), interpolation and fill methods, and computational backends.
151 This approach contrasts with opinionated frameworks that fix defaults and directory structures
152 to maximize automation. While users must make explicit decisions, some steps remain
153 automated to prevent errors. For example, bathymetry smoothing is applied automatically with
154 a non-tunable parameter, since overly small smoothing factors could produce rough bathymetry
155 and crash simulations due to pressure gradient errors. This design choice directly addresses
156 issues new users faced in the MATLAB scripts, and balances **flexibility** and **safety**, enabling
157 transparent experimentation without exposing users to avoidable pitfalls.

158 Another key design consideration is balancing **modular, incremental workflow steps**
159 with **reproducibility**. ROMS-Tools organizes tasks (such as creating InitialConditions,
160 BoundaryForcing, and SurfaceForcing) into small, composable components that can be
161 executed, saved, and revisited independently, rather than following a monolithic, fixed
162 workflow. All components depend on the Grid, but once it is created, the remaining objects
163 are independent. This modular approach avoids unnecessary recomputation when only some
164 inputs change but requires careful tracking of workflow state. To ensure reproducibility,
165 all configuration choices are stored in compact, text-based YAML files. These files are

¹In the future, packages like ROMS-Tools and regional-mom6 could share a common backbone, with model-specific adaptations layered on top.

version-controllable, easy to share, and eliminate the need to transfer large model input NetCDF datasets. This design directly addresses the MATLAB scripts' lack of explicit workflow tracking.

Architecture

At the user-facing level, ROMS-Tools provides high-level objects such as `Grid`, `InitialConditions`, and `BoundaryForcing`. Each object exposes a consistent interface (`.ds`, `.plot()`, `.save()`, `.to_yaml()`), allowing users to call the same methods in sequence or inspect attributes that are always present. This design reduces cognitive overhead, makes workflows predictable, and removes the need for new users to edit raw scripts or manually track intermediate files, as was required with the MATLAB tools.

Internally, ROMS-Tools follows a **layered, modular architecture**. Low-level classes (`LatLonDataset`, `ROMSDataset`) handle data ingestion and preprocessing, including common operations such as subdomain selection and lateral filling. Source-specific datasets (e.g., `ERA5Dataset`, `GLORYSDataset`, `SRTMDataset`) inherit from these base classes and encode dataset-specific conventions like variable names, coordinates, and masking. Adding support for a new data source typically requires only a small subclass to define variable mappings while reusing existing logic, minimizing changes to the core code. High-level classes (`Grid`, `InitialConditions`, `BoundaryForcing`) build on these low-level datasets to produce ready-to-use modeling inputs, performing tasks such as regridding and final assembly. This layered design enhances **extensibility and maintainability**, avoiding the pitfalls of the monolithic MATLAB scripts.

Computational and Data Model Choices

ROMS-Tools is built on `xarray`, which provides a clear, consistent interface for exploring and inspecting labeled, multi-dimensional geophysical datasets. Users can take advantage of `xarray`'s intuitive indexing, plotting, and metadata handling. Optional dask support allows workflows to scale from laptops to HPC systems, enabling parallel and out-of-core computation for very large input and output datasets. By combining modern Python tools with a user-friendly interface, ROMS-Tools addresses the usability challenges that hampered new users in the MATLAB-based workflow.

Research Impact Statement

ROMS-Tools is used by two primary research communities. First, regional ocean modelers use it to generate reproducible input datasets for ROMS simulations; external users include researchers at **PNNL**, **WHOI**, and **UCLA**. Second, researchers in the ocean-based carbon dioxide removal (CDR) community use ROMS-Tools to configure reproducible ROMS-MARBL simulations of climate intervention scenarios, with adopters including **[C]Worthy**, **Carbon to Sea**, **Ebb Carbon**, and **SCCWRP**. All of these groups have contacted the developers directly or engaged in offline discussions regarding their use of the package.

Additional evidence of community uptake comes from public usage metrics. At the time of writing, the GitHub repository shows **119 unique cloners in the past 14 days**, with stars from users at institutions including the University of Waikato, NCAR, University of Maryland, National Oceanography Centre, McGill University, UC Santa Cruz, and others. Distribution statistics indicate **over 3,100 conda-forge downloads in the past six months**, including **68 downloads of the most recent release (v3.3.0)**, and **more than 48,000 total PyPI downloads** (noting that PyPI counts include automated CI usage, whereas conda downloads do not).

ROMS-Tools is also integrated into broader workflows, including **C-Star** (Stephenson & contributors, 2025), an open-source platform to provide scientifically credible monitoring, reporting, and verification (MRV) for the emerging marine carbon market.

213 AI Usage Disclosure

214 Generative AI tools were used to assist with writing docstrings and developing tests for the
 215 ROMS-Tools software, to improve the clarity and readability of the documentation, and to
 216 shorten and edit portions of the manuscript text. All AI-assisted content was reviewed and
 217 verified by the authors for technical accuracy and correctness.

218 Acknowledgements

219 Development of ROMS-Tools has been supported by ARPA-E (DE-AR0001838) and
 220 philanthropic donations to [C]Worthy from the Grantham Foundation for the Environment,
 221 the Chan Zuckerberg Initiative, Founders Pledge, and the Ocean Resilience Climate Alliance.

222 References

- 223 Adcroft, A., Anderson, W., Balaji, V., Blanton, C., Bushuk, M., Dufour, C. O., Dunne, J.
 224 P., Griffies, S. M., Hallberg, R., Harrison, M. J., Held, I. M., Jansen, M. F., John, J. G.,
 225 Krasting, J. P., Langenhorst, A. R., Legg, S., Liang, Z., McHugh, C., Radhakrishnan,
 226 A., ... Zhang, R. (2019). The GFDL Global Ocean and Sea Ice Model OM4.0: Model
 227 Description and Simulation Features. *Journal of Advances in Modeling Earth Systems*,
 228 11(10), 3167–3211. <https://doi.org/10.1029/2019MS001726>
- 229 Arango, H., & contributors. (2024). Rutgers ROMS. In *GitHub repository*. GitHub. <https://github.com/myroms/roms>
- 231 Barnes, A. J., Constantinou, N. C., Gibson, A. H., Kiss, A. E., Chapman, C., Reilly, J.,
 232 Bhagtani, D., & Yang, L. (2024). Regional-mom6: A Python package for automatic
 233 generation of regional configurations for the Modular Ocean Model 6. *Journal of Open*
 234 *Source Software*, 9(100), 6857. <https://doi.org/10.21105/joss.06857>
- 235 Bell, N., Olson, L. N., Schroder, J., & Southworth, B. (2023). PyAMG: Algebraic multigrid
 236 solvers in python. *Journal of Open Source Software*, 8(87), 5495. <https://doi.org/10.21105/joss.05495>
- 238 Busecke, J., & contributors. (2025). Xgcm. In *GitHub repository*. GitHub. <https://github.com/xgcm/xgcm>
- 240 Dai, A., & Trenberth, K. E. (2002). *Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations*. https://journals.ametsoc.org/view/journals/hydr/3/6/1525-7541_2002_003_0660_eofdfc_2_0_co_2.xml
- 243 Dask Development Team. (2016). *Dask: Library for dynamic task scheduling*. <http://dask.pydata.org>
- 245 Egbert, G. D., & Erofeeva, S. Y. (2002). Efficient Inverse Modeling of Barotropic Ocean
 246 Tides. *Journal of Atmospheric and Oceanic Technology*, 19(2), 183–204. [https://doi.org/10.1175/1520-0426\(2002\)019%3C0183:EIMOBO%3E2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019%3C0183:EIMOBO%3E2.0.CO;2)
- 248 Garcia, H. E., Boyer, T. P., Baranova, O. K., Locarnini, R. A., Mishonov, A. V., Grodsky,
 249 A., Paver, C. R., Weathers, K. W., Smolyar, I. V., Reagan, J. R., Seidov, D., & Zweng,
 250 M. M. (2019). *World ocean atlas 2018: Product documentation* (A. Mishonov, Ed.).
 251 NOAA/NCEI.
- 252 Hamilton, D. S., Perron, M. M. G., Bond, T. C., Bowie, A. R., Buchholz, R. R., Guieu, C., Ito,
 253 A., Maenhaut, W., Myriokefalitakis, S., Olgun, N., Rathod, S. D., Schepanski, K., Tagliabue,
 254 A., Wagner, R., & Mahowald, N. M. (2022). Earth, Wind, Fire, and Pollution: Aerosol
 255 Nutrient Sources and Impacts on Ocean Biogeochemistry. *Annual Review of Marine Science*,
 256 14(Volume 14, 2022), 303–330. <https://doi.org/10.1146/annurev-marine-031921-013612>

- 257 Hauser, M., Spring, A., Busecke, J., Driel, M. van, Lorenz, R., & readthedocs-assistant.
258 (2024). *Regionmask/regionmask: Version 0.12.1*. Zenodo. <https://doi.org/10.5281/zenodo.10849860>
- 260 Hedstrom, K., & contributors. (2023). *Pyroms*. In *GitHub repository*. GitHub. <https://github.com/ESMG/pyroms>
- 262 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas,
263 J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan,
264 X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J.-N.
265 (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*,
266 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- 267 Hoyer, S., & Hamman, J. (2017). Xarray: ND labeled arrays and datasets in python. *Journal*
268 *of Open Research Software*, 5(1). <https://doi.org/10.5334/jors.148>
- 269 Huang, Y., Tagliabue, A., & Cassar, N. (2022). Data-Driven Modeling of Dissolved Iron in the
270 Global Ocean. *Frontiers in Marine Science*, 9. <https://doi.org/10.3389/fmars.2022.837183>
- 271 Kok, J. F., Adebisi, A. A., Albani, S., Balkanski, Y., Checa-Garcia, R., Chin, M., Colarco,
272 P. R., Hamilton, D. S., Huang, Y., Ito, A., Klose, M., Leung, D. M., Li, L., Mahowald,
273 N. M., Miller, R. L., Obiso, V., Pérez García-Pando, C., Rocha-Lima, A., Wan, J. S., &
274 Whicker, C. A. (2021). Improved representation of the global dust cycle using observational
275 constraints on dust properties and abundance. *Atmospheric Chemistry and Physics*, 21(10),
276 8127–8167. <https://doi.org/10.5194/acp-21-8127-2021>
- 277 Landschützer, P., Gruber, N., & Bakker, D. C. E. (2016). Decadal variations and trends
278 of the global ocean carbon sink. *Global Biogeochemical Cycles*, 30(10), 1396–1417.
279 <https://doi.org/10.1002/2015GB005359>
- 280 Lauvset, S. K., Key, R. M., Olsen, A., Heuven, S. van, Velo, A., Lin, X., Schirnick, C., Kozyr,
281 A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M.,
282 Perez, F. F., Suzuki, T., & Watelet, S. (2016). A new global interior ocean mapped
283 climatology: The 1° × 1° GLODAP version 2. *Earth System Science Data*, 8(2), 325–340.
284 <https://doi.org/10.5194/essd-8-325-2016>
- 285 Lellouche, E., Jean-Michel Greiner, Bourdallé-Badie, R., Garric, G., Melet, A., Drévillon,
286 M., Bricaud, C., Hamon, M., Le Galloudec, O., Regnier, C., Candela, T., Testut, C.-E.,
287 Gasparin, F., Ruggiero, G., Benkiran, M., Drillet, Y., & Le Traon, P.-Y. (2021). The
288 Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis. *Frontiers in Earth*
289 *Science*, 9. <https://doi.org/10.3389/feart.2021.698876>
- 290 Long, M. C., Moore, J. K., Lindsay, K., Levy, M., Doney, S. C., Luo, J. Y., Krumhardt, K.
291 M., Letscher, R. T., Grover, M., & Sylvester, Z. T. (2021). *Simulations With the Marine*
292 *Biogeochemistry Library (MARBL)*. <https://doi.org/10.1029/2021MS002647>
- 293 Molemaker, J. (2024). *UCLA MATLAB tools*. In *GitHub repository*. GitHub. <https://github.com/nmolem/ucla-tools>
- 295 Molemaker, J., & contributors. (2025a). *UCLA-ROMS*. In *GitHub repository*. GitHub.
296 <https://github.com/CESR-lab/ucla-roms>
- 297 Molemaker, J., & contributors. (2025b). *UCLA-ROMS*. In *GitHub repository*. GitHub.
298 <https://github.com/CWorthy-ocean/ucla-roms>
- 299 Shchepetkin, A. F., & McWilliams, J. C. (2005). The regional oceanic modeling system
300 (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model.
301 *Ocean Modelling*, 9(4), 347–404. <https://doi.org/10.1016/j.ocemod.2004.08.002>
- 302 Stephenson, D., & contributors. (2025). *C-star*. [C]Worthy. <https://github.com/CWorthy-ocean/C-Star>
- 303

- 304 Tozer, B., Sandwell, D. T., Smith, W. H. F., Olson, C., Beale, J. R., & Wessel, P. (2019).
305 Global Bathymetry and Topography at 15 Arc Sec: SRTM15+. *Earth and Space Science*,
306 6(10), 1847–1864. <https://doi.org/10.1029/2019EA000658>
- 307 Wessel, P., & Smith, W. H. F. (1996). A global, self-consistent, hierarchical, high-resolution
308 shoreline database. *Journal of Geophysical Research: Solid Earth*, 101(B4), 8741–8743.
309 <https://doi.org/10.1029/96JB00104>
- 310 Yang, S., Chang, B. X., Warner, M. J., Weber, T. S., Bourbonnais, A. M., Santoro, A. E.,
311 Kock, A., Sonnerup, R. E., Bullister, J. L., Wilson, S. T., & Bianchi, D. (2020). Global
312 reconstruction reduces the uncertainty of oceanic nitrous oxide emissions and reveals a
313 vigorous seasonal cycle. *Proceedings of the National Academy of Sciences of the United*
314 *States of America*, 117(22), 11954–11960. <https://doi.org/10.1073/pnas.1921914117>
- 315 Yeager, S. G., Rosenbloom, N., Glanville, A. A., Wu, X., Simpson, I., Li, H., Molina, M.
316 J., Krumhardt, K., Mogen, S., Lindsay, K., Lombardozzi, D., Wieder, W., Kim, W. M.,
317 Richter, J. H., Long, M., Danabasoglu, G., Bailey, D., Holland, M., Lovenduski, N., ...
318 King, T. (2022). The Seasonal-to-Multiyear Large Ensemble (SMYLE) prediction system
319 using the Community Earth System Model version 2. *Geoscientific Model Development*,
320 15(16), 6451–6493. <https://doi.org/10.5194/gmd-15-6451-2022>
- 321 Zhuang, J., Dussin, R., Huard, D., Bourgault, P., Banihirwe, A., Raynaud, S., Malevich,
322 B., Schupfner, M., Filipe, Levang, S., Gauthier, C., Jüling, A., Almansi, M., Scott,
323 R., RondeauG, Rasp, S., Smith, T. J., Stachelek, J., Plough, M., & Li, X. (2023).
324 xESMF: Universal regridding for geospatial data. In *GitHub repository*. Zenodo. <https://doi.org/10.5281/zenodo.4294774>
- 325

DRAFT