

Homework 3

TIAN Chenyu

October 17, 2019

-
- **Acknowledgments:** This template takes some materials from course CSE 547/Stat 548 of Washington University:
<https://courses.cs.washington.edu/courses/cse547/17sp/index.html>.
 - **Collaborators:** I finish this homework by myself.
-

2.1. The log-likelihood of the naive Bayes model can be written as

$$\begin{aligned} l(\phi_y, \phi_j(x | y)) &= \sum_{i=1}^m \log p(x^{(i)}, y^{(i)}) \\ &= \sum_{i=1}^m (\log p(x^{(i)} | y^{(i)}) + \log p(y^{(i)})) \\ &= \sum_{i=1}^m \left(\sum_{j=1}^d \log p(x_j^{(i)} | y^{(i)}) + \log p(y^{(i)}) \right) \\ &= \sum_{i=1}^m \sum_{j=1}^d \log \phi_j(x_j^{(i)} | y^{(i)}) + \sum_{i=1}^m \log \phi_{y^{(i)}} \end{aligned}$$

Also, there exist constrains for $\phi_y, \phi_j(x | y)$:

$$\begin{aligned} \sum_{y=1}^k \phi_y &= 1 \\ \sum_{x \in \{0,1\}} \phi_j(x | y) &= 1 \quad y \in \{1, \dots, K\} \end{aligned}$$

Then to find the maximum value of l with bringing these constrains into equation using lagrange multiplier

$$\begin{aligned} g(l, \lambda, \lambda_{jk}) &= \sum_{i=1}^m \sum_{j=1}^d \log \phi_j(x_j^{(i)} | y^{(i)}) + \sum_{i=1}^m \log \phi_{y^{(i)}} \\ &\quad + \lambda \left(\sum_{y=1}^k \phi_y - 1 \right) + \lambda_{jk} \left(\sum_{x \in \{0,1\}} \phi_j(x | k) - 1 \right) \end{aligned}$$

Using lagrange multiplier method to find the parameters of the maximum value, there exists:

$$\begin{aligned} \frac{dg}{d\phi_y} &= \sum_{i=1}^m \mathbb{1}(y^{(i)} = y) \frac{1}{\phi_y} + \lambda = 0 \\ \Rightarrow \sum_{i=1}^m \mathbb{1}(y^{(i)} = 1) \frac{1}{\phi_1} &= \sum_{i=1}^m \mathbb{1}(y^{(i)} = 2) \frac{1}{\phi_2} = \dots = \sum_{i=1}^m \mathbb{1}(y^{(i)} = K) \frac{1}{\phi_K} \end{aligned}$$

Then here has

$$\phi_y = \frac{\sum_{i=1}^m \mathbf{1}(y^{(i)} = y)}{m}$$

Also,

$$\begin{aligned} \frac{dg}{d\phi_j(x | y)} &= \sum_{i=1}^m \mathbf{1}(y^{(i)} = y) \mathbf{1}(x_j^{(i)} = x) \frac{1}{\phi_j(x | y)} + \lambda_j(x | y) = 0 \\ \Rightarrow \sum_{i=1}^m \mathbf{1}(y^{(i)} = y) \mathbf{1}(x_j^{(i)} = 0) \frac{1}{\phi_j(x = 0 | y)} &= \sum_{i=1}^m \mathbf{1}(y^{(i)} = y) \mathbf{1}(x_j^{(i)} = 1) \frac{1}{\phi_j(x = 1 | y)} \end{aligned}$$

Thus,

$$\phi_j(x | y) = \frac{\sum_{i=1}^m \mathbf{1}(y^{(i)} = y) \mathbf{1}(x_j^{(i)} = x)}{\sum_{i=1}^m \mathbf{1}(y^{(i)} = y)}$$