



Directional Dark Matter

searches with CYGNO



David J. G. Marques* on behalf of the CYGNO collaboration:

F. Amaro, R. Antonietti, E. Baracchini, L. Benussi, S. Bianco, R. Campagnola, L. G. M. Carvalho, C. Capoccia, M. Caponero, D. S. Cardoso, G. Cavoto, I. A. Costa, A. Croce, M. D'Astolfo, E. Di Marco, G. D'Imperio, E. Dané, G. Dho, D. Fiorina, F. Di Giambattista, F. Iacoangeli, Z. Islam, E. Kemp, H. P. Lima Júnior, G. S. P. Lopes, G. Maccarrone, R. D. P. Mano, R. R. Marcelo Gregorio, D. J. G. Marques*, G. Mazzitelli, A.G. McLean, P. Meloni, A. Messina, C. M. B. Monteiro, R. A. Nobrega, I. F. Pains, E. Paoletti, L. Passamonti, S. Pelosi, F. Petrucci, S. Piacentini, D. Piccolo, D. Pierluigi, D. Pinci, A. Prajapati, F. Renga, R. J. C. Roque, F. Rosatelli, A. Russo, J. M. F. dos Santos, G. Saviano, P. A. O. Silva, N. Spooner, R. Tesauro, S. Tomassini, S. Torelli, D. Tozzi



Part of this project has been funded by the European Union's Horizon 2020 research and innovation programme under the ERC Consolidator Grant Agreement No 818744

WIMPs - How to see them?


In the WIMP model, DM forms a halo within our galaxy

+


Solar system rotates around galaxy towards Cygnus constellation



Earth susceptible to an apparent WIMP wind from Cygnus direction!



- $\text{SM} + x \rightarrow \text{SM} + x$
- Direct detection** of nuclear recoil




WIMPs - How to see them?

In the WIMP model, DM forms a halo within our galaxy


+

Solar system rotates around galaxy towards Cygnus constellation

Earth susceptible to an apparent WIMP wind from Cygnus direction!





- $\text{SM} + x \rightarrow \text{SM} + x$
- Direct detection of nuclear recoil





DM interacting in our detectors would create different effects:

- **ENERGY** \Rightarrow Excess would result in falling exponentials.
 - The **background** has a similar spectrum, as we've seen recently in many experiments...



- **TIME** \Rightarrow Results in a few % annual modulation.



<https://doi.org/10.1155/2014/605659>



G S
S I

Directionality and *beyond the neutrino fog*


Directionality and beyond the neutrino fog

Exploring the **DIRECTION** dependency results in

a characteristic effect - anisotropy in the angular distribution of nuclear recoils



No background can mimic





Where other experiments struggle to **find striking features to prove the existence of DM**, **directional discrimination** emerges as a unique and efficient strategy to **positively identify Dark Matter!**

Directionality and beyond the neutrino fog

Exploring the **DIRECTION** dependency results in a characteristic effect - anisotropy in the angular distribution of nuclear recoils



No background can mimic



Where other experiments struggle to **find striking features to prove the existence of DM**, **directional discrimination** emerges as a unique and efficient strategy to **positively identify Dark Matter!**

The **CEvNS** produces NRs identical to the DM-induced ones.



To **search DM at smaller cross-sections**, experiments need to venture into the neutrino fog



Below 10 GeV/c² → Mostly **Solar neutrinos**



In galactic coords., the Sun and Cygnus are never superimposed!*



- Searching **beyond the neutrino fog**
- Properties of the **solar neutrino flux and DM halo**




A CYGNus TPC module
with Optical readout

CYGNO - What's the setup?


TPC

Triple GEM

Charge multiplication




Camera + PMT
Light from **gas scintillation**
during electron avalanche





Carbon tetrafluoride (CF4)

→ Significant light yield at the camera's QE peak (~600 nm)




CYGNO - What's the setup?

TPC → **Triple GEM**
Charge multiplication → **Camera + PMT**
Light from **gas scintillation**
during electron avalanche




With the high granularity of
the camera, we measure
energy + X & Y coordinates




CYGNO - What's the setup?


TPC → **Triple GEM**
Charge multiplication





→ **Camera + PMT**
Light from **gas scintillation**
during electron avalanche



With the high granularity of
the camera, we measure
energy + X & Y coordinates




1. Independent **energy** measurement.
2. Electrons **times of arrival** ⇒ **dZ coordinate** (track's tilt)



Cygnو - What's the setup?

TPC

cathode



With the
the camera,

energy + X & Y coordinates




α -particle

Track's deposited energy



topology (dE/dx)

Head-tail asymmetry
+ recoil direction

↓
Directionality



Particle
↓
BG rejection



ent
ment.

arrival ⇒
ck's tilt)

K
→









→

→

→


→

Some cool pictures



CYGNO - Dark Matter paradigm

CYGNO Dark Matter exploration region




$< 10 \text{ GeV}/c^2$

→ To observe lower WIMP masses:

- ◆ **Low thresholds** are necessary, since lower m_χ originate lower energy recoils.
- ◆ **Light nuclei** used to maximize energy transfer.

CYGNO - Dark Matter paradigm

CYGNO Dark Matter exploration region



Low Density @ atm pressure


- Allows tracks of up to **millimetres at few keV** without compromising exposure.

< 10 GeV/c²

- To observe lower WIMP masses:
 - ◆ **Low thresholds** are necessary, since lower m_{χ} originate lower energy recoils.
 - ◆ **Light nuclei** used to maximize energy transfer.

Helium (He)

- Light target for SI in low mass range.



Fluorine (F)



- Heavier target to intermediate WIMP masses.
- One of the highest sensitivity to SD coupling.

CYGNO - Roadmap

Several ongoing efforts in different fronts:


- 3D reconstruction
- Directionality
- ER vs. NR (+ML)
- Shielding optimization
- Background data vs. MC
- DM Sensitivity
- Design and Commissioning of CYGNO_04
- **Enhancement of the light yield**
- **Negative Ion drift**

Check Giorgio Dho's poster!




Where we are at...

LIME





LIME - The concept



- 50 L & 50 cm drift gaseous TPC with **Copper ring** field cage.
- Atm pressure (910 mbar), room temperature and He:CF4, 60:40
- Triple 33x33 cm² GEM stack for amplification
- Optical readout ⇒ 4 PMTs + 1 sCMOS camera (ORCA Fusion)

LIME - The concept



- 50 L & 50 cm drift gaseous TPC with **Copper ring** field cage.
- Atm pressure (910 mbar), room temperature and He:CF₄, 60:40
- Triple 33x33 cm² GEM stack for amplification
- Optical readout ⇒ 4 PMTs + 1 sCMOS camera (ORCA Fusion)

- ★ LIME was placed underground at LNGS in the beginning of 2022.
- ★ Commissioning: tests on DAQ, remote control, slow control, gas quality,etc.
- ★ Technology test in a realistic underground environment for rare event searches
- ★ Study of shielding for validation of the CYGNO Monte Carlo
- ★ Multiple radioactive source runs : ⁵⁵Fe, ¹³⁷Ba, ¹⁵²Eu, ²⁴¹Am + ²⁴¹Am⁹Be




Live monitoring of detector performance & data quality

LIME - Data vs. MC

Phase	Shielding	GEM V [V]	# pictures	Live time [s]	Rate PMTs [Hz]
Run 1	None	420	285665	175627	30
Run 2	4 cm Cu	440	297992	191382	3.5
Run 3	10 cm Cu	440	171579	191471	1.6
Run 4	+40 cm H ₂ O	Great external neutron suppression ⇒ <i>Under analysis...</i>			

LIME - Data vs. MC


Phase	Shielding	GEM V [V]	# pictures	Live time [s]	Rate PMTs [Hz]
Run 1	None	420	285665	175627	30
Run 2	4 cm Cu	440	297992	191382	3.5
Run 3	10 cm Cu	440	171579	191471	1.6
Run 4	+40 cm H ₂ O	Great external neutron suppression ⇒ Under analysis...			




- Shielding strongly **suppressed external background**
- Difference in Runs 2 & 3 attributed to **internal background**
excess (contaminations of the detector – materials, gas, etc.)
- **LIME was not meant to be radiopure**
 - ◆ Not all the components were measured
 - ◆ Not all the contaminants were taken into account (eg. Rn)

LIME - Data vs. MC

Phase	Shielding	GEM V [V]	# pictures	Live time [s]	Rate PMTs [Hz]
Run 1	None	420	285665	175627	30
Run 2	4 cm Cu	440	297992	191382	3.5
Run 3	10 cm Cu	440	171579	191471	1.6
Run 4	+40 cm H ₂ O		Great external neutron suppression \Rightarrow Under analysis...		



- Shielding strongly **suppressed external background**
- Difference in Runs 2 & 3 attributed to **internal background**
excess (contaminations of the detector – materials, gas, etc.)
- **LIME was not meant to be radiopure**
 - ◆ Not all the components were measured
 - ◆ Not all the contaminants were taken into account (eg. Rn)




LIME - Run 5 objectives

- In run 5 we want to:
 - ◆ Take data in a less saturated configuration

LIME - Run 5 objectives

→ In run 5 we want to:

- ◆ Take data in a less saturated configuration
- ◆ Longer data taking ~ Perform underground LNGS neutron flux measurement.
 - ~ 250 neutrons in 6 months.
 - Improve current knowledge, especially at low energy (< 1 MeV)
 - **Allow us to study in depth:** 3D, directionality, energy resolution on NRs and study our ability to select NR signals.



10^4 rejection @ 20 keV

Close to 10^5 rejection @ 25 keV (from simulated data)

From BG considerations, **BG-free dataset seems achievable at >20 keV**

⇒ **Constitutes a landmark for CYGNO for the study of WIMP-like events**


LIME - 3D reconstruction

- Merging **camera** with **PMTs** information allows us to get a **full description** of the ionization event.
 - ◆ With this, we will be able to improve our **particle ID**, **reject backgrounds** from known sources, and fully **characterize the 3D direction** of the incoming particles: 

LIME - 3D reconstruction

- Merging **camera** with **PMTs** information allows us to get a **full description** of the ionization event.
 - ◆ With this, we will be able to improve our **particle ID**, **reject backgrounds** from known sources, and fully **characterize the 3D direction** of the incoming particles: **Directionality**

- ★ **Alpha selection:** cluster density, waveform skewness, etc. + **ER rejection + association** between trigger and cluster
- 1. **Camera analyser class** retrieves **X-Y Angle** and the **ΔXY**
- 2. **Time over threshold** determines **ΔZ** ⇒ Together with camera X-Y angle ⇒ **Z-angle**
- 3. The **position of the Bragg** in PMT waveforms gives **Z-angle signal = head-tail** = Towards cathode or GEMs



LIME - 3D reconstruction

- Merging **camera** with **PMTs** information allows us to get a **full description** of the ionization event.
 - ◆ With this, we will be able to improve our **particle ID**, **reject backgrounds** from known sources, and fully **characterize the 3D direction** of the incoming particles: **Directionality**




- ★ **Alpha selection:** cluster density, waveform skewness, etc. + **ER rejection + association** between trigger and cluster
- 1. **Camera analyser class** retrieves **X-Y Angle** and the **ΔXY**
- 2. **Time over threshold** determines **ΔZ** ⇒ Together with camera X-Y angle ⇒ **Z-angle**
- 3. The **position of the Bragg** in PMT waveforms gives **Z-angle signal = head-tail** = Towards cathode or GEMs

Reconstructed info:

- $\Delta Z + \Delta XY$
- Phi + Theta angles
- Signal of theta
- + phi == Head-tail


↓

3D reconstruction



next step...

CYGNO-04




LNGS Hall F

CYGNO-04 - Phase I



- Test scalability on realistic scale + all ancillary systems
- Test feasibility of physics reach for directional DM searches with a radiopure large detector










CYGNO-04 - Phase I



- Test scalability on realistic scale + all ancillary systems
- Test feasibility of physics reach for directional DM searches with a radiopure large detector

- Back-to-back 0.4 m³ TPC, with central cathode.
- Projected shielding composed of **10 cm Cu + 100 cm H₂O**
- Radioactivity of all materials will be measured
 - ◆ Improve MC vs. data ⇒ Determine sensitivities
- Currently **validating** all the components:
 - ◆ **Camera:** Fusion ⇒ Quest * 6
 - ◆ **PMT:** Position under study ⇒ 8
 - ◆ **Field cage:** copper strips on insulator support
 - ◆ **GEMs:** 50x80 cm²
- Timeline: Commiss. 01/2025 ⇒ Data 08/2025 - 12/2026





R&D - Ongoing projects

- Negative ions


Reduce diffusion

during drift by adding SF₆ (thus negative ions) to the gas mixture



- Light enhancement with strong electric fields

Through strong electric fields, light is increased with constant charge and energy resolution





R&D - Ongoing projects

- Negative ions**


Reduce diffusion

during drift by adding SF₆ (thus negative ions) to the gas mixture



- Addition of H-based gases in the mixture**


Improvement of low Dark Matter Mass detection sensitivity



Secondary scintillation yield from GEM electron avalanches in He-CF₄ and He-CF₄-isobutane for CYGNO — Directional Dark Matter search with an optical TPC

- Light enhancement with strong electric fields**

Through strong electric fields, light is increased with constant charge and energy resolution



- Internal background reduction**

- Building low radioactivity camera sensor and lens together with Hamamatsu/BMI experts





R&D - Ongoing projects

- Negative ions**


Reduce diffusion

during drift by adding SF₆ (thus negative ions) to the gas mixture



- Addition of H-based gases in the mixture**


Improvement of low Dark Matter Mass detection sensitivity



Secondary scintillation yield from GEM electron avalanches in He-CF4 and He-CF4-isobutane for CYGNO — Directional Dark Matter search with an optical TPC

- Light enhancement with strong electric fields**

Through strong electric fields, light is increased with constant charge and energy resolution



Check Giorgio Dho's poster!

- internal background reduction**

- Building low radioactivity camera sensor and lens together with Hamamatsu/BMI experts



Conclusions

- The [CYGNO](#) collaboration is developing a [high-precision gaseous TPC](#) at atmospheric pressure with [optical readout](#).
- The main focus is the [directional direct search](#) of [DM WIMP-like particles](#) in the [low mass range](#) (0.5-10 GeV).
- Through [nuclear recoil direction](#), solar neutrinos can be discriminated and [unambiguous confirmation of DM](#) is possible.
- The [50L LIME prototype](#) was installed in the [underground LNGS](#) facilities.
 - ◆ [Commissioning](#) tests, [background vs. MC](#) evaluations and [measurements with sources](#) are being carried out.
- [CYGNO-04](#), will allow us to test the experiment's scalability and [physics reach](#).
- [CYGNO-30](#) is under study, with its sensitivities looking promising.
- Several [R&D projects](#) are ongoing in order to find [optimal means of TPC operation](#):
 - ◆ [Light enhancement](#) observed, and its [potentialities](#) are under study!
 - ◆ [Negative ion drift](#) observed for the first at [atmospheric pressure](#) and with [PMTs – stay tuned!](#)



...check out our white paper!
[The CYGNO Experiment - Instruments](#)



Thank you for
your attention!

The CYGNO Project counts with the collaboration
of several international researchers, coming from:







G S
S I

Backup

& more

details


LIME - Light yield stability



- This corresponds to about **50 days** of "total" data taking.
- The **light yield stability** is evaluated through the mean value of LY for high energy tracks.
- Several efforts have been made throughout the initial runs to **stabilize** the light yield inside LIME, and, in run 4 (latest), we can see a **constant light yield at 3% sigma**.

LIME - AmBe run details

AmBe run



→ AmBe + background simulation matches

data @ $E < 40$ keV

- ◆ External background ✓
- ◆ Source simulation ✓
- ◆ Internal background - Shape ✓
- ◆ Internal background - Absolute 🧑‍🔧





Figure 6.27: Comparison between the data acquired with the AmBe source during Run 3 (blue) and the simulated background (red) and AmBe-induced events (green), below 100 keV. The AmBe-induced events dominate the spectrum below ~ 50 keV.

CYGNO-30 - Prospects

- **Low mass (0.5 - 10 GeV) directional DM searches**
- > 2027
- **30 - 100 m³ detector**
- **0.5 - 1 keV_{ee} energy threshold**
- **30° angular resolution**



Expected **SI** and **SD** (90% CL)



interaction cross-section exclusion

Quenching factor simulated

with **SRIM** → Direct



measurement incoming!

He / (eventually H) allows us to explore very low DM masses!



The CYGNUS project

CYGNUS is part of a proto-collaboration, CYGNUS, focused on establishing a **Galactic Directional Recoil Observatory** that could test and study DM hypothesis beyond the neutrino floor.



Within the CYGNUS collaboration, several approaches are being studied.

The Italian group, CYGNUS, is developing a **gaseous TPC** based on the setup:

GEMs + sCMOS + PMT to test Optical Readout