

Lecture Notes - MATH 455

Charles Zitella

January 15, 2026

Contents

<i>Abstract Metric and Topological Spaces</i>	1
<i>Metric Spaces Review</i>	1
<i>Compactness, Separability</i>	2
<i>Arzelà-Ascoli</i>	4
<i>Baire Category Theorem</i>	7
<i>Topological Spaces</i>	10
<i>Separation, Countability, Separability</i>	10
<i>Continuity and Compactness</i>	10

Abstract Metric and Topological Spaces

Metric Spaces Review

Throughout, assume X is a non empty set.

Definition 1. (Metric): $p : X \times X \rightarrow \mathbb{R}$ is called a *metric*, and thus (X, p) a metric space, if for all $x, y, z \in X$

- $p(x, y) \geq 0$,
- $p(x, y) = 0 \iff x = y$,
- $p(x, y) = p(y, x)$,
- $p(x, y) \leq p(x, z) + p(z, y)$ (Triangle Inequality).

Definition 2. (Norm): Let X be a vector space.¹ A function $\|\cdot\| : X \rightarrow [0, \infty)$ is called a *norm*, and thus $(X, \|\cdot\|)$ a *normed vector space*, if for all $u, v \in X$ and $\alpha \in \mathbb{R}$

- $\|u\| = 0 \iff u = 0$,
- $\|u + v\| \leq \|u\| + \|v\|$,
- $\|\alpha u\| = |\alpha| \|u\|$.

Remark 3. A norm induces a metric by $p(x, y) := \|x - y\|$.

Example 4. Examples of normed vector spaces:

1. $(\mathbb{R}^n, |\cdot|)$ where $|x| = (x_1^2 + \dots x_n^2)$
2. $L^p(E)$ for $E \subseteq \mathbb{R}^n, 1 \leq p \leq \infty$ where $\|f\|_{L^p(E)} = (\int_E |f(x)|^p dx)^{\frac{1}{p}}$
3. Discrete metric: if X is a non empty set, then $p(x, y) = \begin{cases} 0 & x = y \\ 1 & x \neq y \end{cases}$
4. $C([a, b]) = \{f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is continuous on } [a, b]\}$ for $a, b \subseteq \mathbb{R}$. Then, $\|f\|_{\infty} := \sup_{x \in [a, b]} |f(x)| = \max_{x \in [a, b]} |f(x)|$, $p(f, g) = \|f - g\|_{\infty}$

Definition 5. Given two metrics p, σ on X , we say they are *equivalent* if \exists a $C > 0$ such that $\frac{1}{C}\sigma(x, y) \leq p(x, y) \leq C\sigma(x, y)$ for every $x, y \in X$. A similar definition follows for equivalence of norms.

Given a metric space (X, p) , then, we have the notion of

- open balls $B(x, r) = \{y \in X : p(x, y) \leq r\}$
- open sets (subsets of X with the property that for every $x \in X$, there is a constant $r > 0$ such that $B(x, r) \subseteq X$), closed sets, closures, and
- *convergence*

Definition 6. (Convergence): $\{x_n\}_{n=1}^{\infty} \subseteq X$ converges to x in (X, p) if $\lim_{n \rightarrow \infty} p(x_n, x) = 0$

We have several (equivalent) notions, then, of continuity; via sequences, $\epsilon - \delta$ definition, and by pullbacks (inverse images of open sets are open).

Definition 7. (Uniform Continuity): $f : (X, p) \rightarrow (\mathbb{R}, |\cdot|)$ uniformly continuous if f has a "modulus of continuity", i.e. there is a continuous function $\omega : [0, \infty) \rightarrow [0, \infty)$ such that $\lim_{t \rightarrow 0^+} \omega(t) = 0$, and

$$|f(x) - f(y)| \leq \omega(p(x, y))$$

for every $x, y \in X$

Remark 8. For instance, we say f Lipschitz continuous if there is a constant $C > 0$ such that $\omega(\cdot) = C(\cdot)$. Let $\alpha \in (0, 1)$. We say f α -Holder continuous if $\omega(\cdot) = C(\cdot)^{\alpha}$ for some constant C .

Definition 9. (Completeness): We say (X, p) *complete* if every Cauchy sequence in (X, p) converges to a point in X .

Remark 10. let $E \subseteq X$ and (X, p) complete metric space. Then (E, p) is complete iff $E \subseteq X$ is closed (so limits belong to E)

Compactness, Separability

Definition 11. (Open Cover, Compactness): $\{X_\lambda\}_{\lambda \in \Lambda} \subseteq 2^X$, where X_λ open in X and Λ an arbitrary index set, an *open cover* of X if for every $x \in X$, $\exists \lambda \in \Lambda$ such that $x \in X_\lambda$. ² X is *compact* if every open cover of X admits a finite subcover. We say $E \subseteq X$ compact if (E, p) compact.

2^X denotes the power set of X , i.e. the set of all subsets of X .

² A cover is finite if $|\Lambda| < \infty$

Remark 12. for $E \subseteq X$, $X_\lambda \subseteq E$ is open in (E, p) iff X_λ is open in (X, p) Therefore, $E \subseteq X$ is compact iff every open cover of E (in X) has a finite subcover.

Remark 13. This definition leads to another definition of compactness based on the finite intersection property.

Useful consequence: if (X, p) is compact metric space, and $\{E_k\}_{k=1}^\infty \subseteq X$ closed, and $E_{k+1} \subseteq E_k \forall k$, $\cap_{k=1}^\infty E_k \neq \emptyset$

Definition 14. (Totally Bounded, ϵ -nets): (X, p) is *totally bounded* if $\forall \epsilon > 0$, there is a finite cover of X of balls with radius $\epsilon > 0$. ³ If $E \subseteq X$, an ϵ -net of E is a collection $\{B(x_i, \epsilon)\}_{i=1}^N$ such that $E \subseteq \bigcup_{i=1}^N B(x_i, \epsilon)$ and $x_i \in X$ (note that x_i need not be in E).

³ Totally bounded implies (X, p) is bounded

Definition 15. (Sequentially Compact): (X, p) *sequentially compact* if every sequence in X has a convergent subsequence whose limit is in X .

Definition 16. (Relatively/Pre-Compact): $E \subseteq X$ *precompact* if \bar{E} compact.

Theorem 17. TFAE:

1. X complete and totally bounded;
2. X compact;
3. X sequentially compact.

Remark 18. TFAE:

1. E is totally bdd and Cauchy Seq. converge
2. E is precompact
3. $\forall \{x_k\}_{k=1}^\infty \subseteq E$, \exists a convergent subsequence

Let $f : (X, p) \rightarrow (\mathbb{R}, |\cdot|)$ continuous with (X, p) compact. Then,

- $f(X)$ compact in $(\mathbb{R}, |\cdot|)$;
- The max and min of f over X are attained;
- f is uniformly continuous.

Lemma 19. Any cauchy sequence ⁴ converges iff it has a convergent subsequence.

⁴ $\forall \epsilon > 0, \exists N > 0$ s.t. $\forall m, n > N$, $\|x_n - x_m\| < \epsilon$

Proof.

\Rightarrow If $\{f_n\}_{n=1}^\infty$ converges, then $\exists f : X \rightarrow \mathbb{R}$ s.t. $\|f_n - f\|_\infty \rightarrow 0$, so all subsequences also converge to f .

\Leftarrow Now assume \exists a subsequence $\{f_{n_k}\}_{k=1}^\infty \subseteq C(X)$ s.t. $\lim_{k \rightarrow \infty} f_{n_k} = f$ in $C(X) \iff \|f_{n_k} - f\|_\infty \rightarrow 0$. Suppose for the purpose of contradiction that $f_n \not\rightarrow f$. Thus, $\exists \epsilon > 0$, and a subsequence $\{f_{n_j}\}_{j=1}^\infty \subseteq C(X)$ s.t. $\|f_{n_j} - f\|_\infty > \epsilon$ for every $j \geq 1$. Then,

$$\|f_{n_k} - f_{n_j}\|_\infty \geq \|f_{n_j} - f\|_\infty - \|f - f_{n_k}\|_\infty > \epsilon - \frac{\epsilon}{2} = \frac{\epsilon}{2}$$

for k sufficiently large and for n_k, n_j large enough. But this violates $\{f_n\}_{n=1}^\infty$ being Cauchy. (Contradiction), so we must have $f_n \rightarrow f$ in $C(X)$. \square

Let $C(X) := \{f : X \in \mathbb{R} \mid f \text{ continuous}\}$ and $\|f\|_\infty := \max_{x \in X} |f(x)|$ the sup norm. Then,

Proposition 20. *Let (X, p) compact. Then $(C(X), \|\cdot\|_\infty)$ is complete.*

Proof. let $\{f_n\}_{n=1}^\infty \subseteq C(X)$ be Cauchy. Fix $k \in \mathbb{N}$. By Cauchy defn, let $\epsilon = 2^{-k}$, so $\exists N_k$ sufficiently large s.t. $\|f_{N_k} - f_{N_k+1}\|_\infty < 2^{-k}$. We can then choose $\{n_k\}_{k=1}^\infty$ s.t. $n_k \rightarrow \infty$ and $\|f_{n_k} - f_{n_{k+1}}\| < 2^{-k} \quad \forall k \in \mathbb{N}$. Let $j \in \mathbb{N}$. Then

$$\|f_{n_{k+j}} - f_{n_k}\|_\infty \leq \sum_{\ell=k}^{k+j-1} \|f_{n_{\ell+1}} - f_{n_\ell}\|_\infty \leq \sum_{\ell=k}^{k+j-1} 2^{-\ell} \leq \sum_{\ell=k}^{\infty} 2^{-\ell} \xrightarrow{k \rightarrow \infty} 0$$

In particular, $\forall x \in X$ fixed, let $c_k := f_{n_k}(x)$. Then $|c_{k+j} - c_k| \leq \|f_{n_{k+j}} - f_{n_k}\|_\infty \rightarrow 0 \quad \forall j \in \mathbb{N}$. Thus $\{c_k\}_{k=1}^\infty \subseteq \mathbb{R}$ is Cauchy, so by completeness of \mathbb{R} , $\exists \bar{c} \in \mathbb{R}$ s.t. $\lim_{k \rightarrow \infty} c_k = \bar{c} =: f(x)$ Doing this $\forall x \in X$, we have

$$\begin{aligned} |f_{n_k}(x) - f(x)| &= \lim_{j \rightarrow \infty} |f_{n_k}(x) - f_{n_{k+j}}(x)| \\ &\leq \lim_{j \rightarrow \infty} \|f_{n_k} - f_{n_{k+j}}\|_\infty \\ &\leq \sum_{\ell=k}^{\infty} 2^{-\ell} \xrightarrow{k \rightarrow \infty} 0 \end{aligned}$$

$\Rightarrow \|f_{n_k} - f\|_\infty = \sup_{x \in X} |f_{n_k}(x) - f(x)| \xrightarrow{k \rightarrow \infty} 0$, so $f_{n_k} \rightarrow f$ in $C(X)$. Finally, by the lemma this implies $f_n \rightarrow f$ in $C(X)$, so $(C(X), \|\cdot\|_\infty)$ is complete. \square

Definition 21. (Density/Separability): A set $D \subseteq X$ is called *dense* in (X, p) if for every nonempty open subset $A \subseteq X$, $D \cap A \neq \emptyset$. We say that X is *separable* if there is a countable dense subset $D \subseteq X$.

If A dense in X , then \bar{A} dense in X

Proposition 22. *If X compact, then X is separable*

Proof. Since X is compact, it is totally bounded. Therefore, for $n \in \mathbb{N}$, there is some K_n and $\{x_i^n\} \subseteq X$ such that $X \subseteq \bigcup_{i=1}^{K_n} B(x_i^n, \frac{1}{n})$. Then, $D = \bigcup_{n=1}^\infty \bigcup_{i=1}^{K_n} \{x_i^n\}$ countable and dense in X \square

Arzelà-Ascoli

Goal: Find suitable conditions for a sequence to have a convergent subsequence in $(C(X), \|\cdot\|_\infty)$.

Definition 23. (Equicontinuous): A family $\mathcal{F} \subseteq C(X)$ is called *equicontinuous* at $x \in X$ if $\forall \epsilon > 0$ there exists a $\delta_x > 0$ such that if $p(x, x') < \delta_x$ then $|f(x) - f(x')| < \epsilon$ for every $f \in \mathcal{F}$. \mathcal{F} is pointwise equicontinuous on X if \mathcal{F} is equicontinuous at every point $x \in X$.⁵

Example 24. Fix $M > 0, [a, b] \subseteq \mathbb{R}$. $\mathcal{F} := \{f \in C([a, b]) \cap C'((a, b)) \mid |f'| \leq M\}$. By Mean Value Theorem, $|f(x) - f(y)| \leq |f'(x^*)| |x - y| \leq M |x - y|$ for some $x^* \in [x, y]$, so $\forall x \in [a, b]$ if $|x - y| < \frac{\epsilon}{M}$ then $|f(x) - f(y)| < \epsilon$, $\forall f \in \mathcal{F}$, therefore \mathcal{F} is pointwise equicontinuous on $[a, b]$.

Example 25. Consider $f_n(x) := x^n$ on $[0, 1]$. Then $\{f_n\}_{n=1}^\infty$ is non equicontinuous at $x = 1$. $f_n(1) = 1 \forall n$, but the threshold to be close to $f_n(1)$ is not uniform on n .

Definition 26. (Pointwise, Uniform Boundedness): $\{f_n\}$ pointwise bounded if $\forall x \in X, \exists M(x) > 0$ such that $|f_n(x)| \leq M(x) \forall n$, and uniformly bounded if such an M exists independent of X .

Definition 27. (Uniform Equicontinuous): $\mathcal{F} \subseteq C(X)$ is uniformly equicontinuous on X if $\forall \epsilon > 0 \exists \delta > 0$ s.t. $\forall x, y \in X$ if $p(x, y) < \delta$, then $|f(x) - f(y)| < \epsilon$, $\forall f \in \mathcal{F}$.

Remark 28. \mathcal{F} equicontinuous at $x \iff$ all $f \in \mathcal{F}$ share the same modulus of continuity at x , i.e. $\exists \omega_x$ s.t. $|f(x) - f(y)| \leq \omega_x |x - y|$, $\forall f \in \mathcal{F}$.

Proposition 29. (Sufficient Conditions for Uniform Equicontinuity):

1. $\mathcal{F} \subseteq C(X)$ is uniformly Lipschitz continuous, i.e. $\exists M > 0$ s.t. $|f(x) - f(y)| \leq Mp(x, y) \forall f \in \mathcal{F}$;
2. $\mathcal{F} \subseteq C(X) \cap C^1(X)$ has a uniform L^∞ bound on the 1st derivative (same as earlier example, by MVT);
3. If (X, p) is compact and $\mathcal{F} \subseteq C(X)$ is pointwise equicontinuous on $X \Rightarrow \mathcal{F}$ is uniformly equicontinuous (Homework).

Lemma 30. (Arzelà-Ascoli Lemma): Let X be separable and let $\{f_n\}_{n=1}^\infty \subseteq C(X)$ be pointwise bounded and equicontinuous. Then, there is a function $f \in C(X)$ and a subsequence $\{f_{n_k}\}_{k=1}^\infty$ which converges pointwise to f on all of X .

Proof. Let $D = \{x_j\}_{j=1}^\infty \subseteq X$ be a countable dense subset of X . Since $\{f_n\}$ is pointwise bounded, $\{f_n(x_1)\}$ as a sequence of real numbers is bounded and so by Bolzano-Weierstrass, there is a convergent subsequence $\{f_{n(1,k)}(x_1)\}_k$ that converges to some $a_1 \in \mathbb{R}$. Consider now $\{f_{n(1,k)}(x_2)\}_k$, which is again a bounded sequence of \mathbb{R} and so has a convergent subsequence, call it $\{f_{n(2,k)}(x_2)\}_k$, which converges to some $a_2 \in \mathbb{R}$. Note that $\{f_{n(2,k)}\} \subseteq \{f_{n(1,k)}\}$, so also $f_{n(2,k)}(x_1) \rightarrow a_1$ as $k \rightarrow \infty$. We can repeat this procedure, producing a sequence of real numbers $\{a_\ell\}$, and for each $j \in \mathbb{N}$ a subsequence $\{f_{n(j,k)}\}_k \subseteq \{f_n\}$ such that $f_{n(j,k)}(x_\ell) \rightarrow a_\ell$ for each $1 \leq \ell \leq j$. Define then

$$f : D \rightarrow \mathbb{R}, \quad f(x_j) := a_j$$

⁵ if $|\mathcal{F}| < \infty$, then \mathcal{F} is pointwise equicontinuous on X .




Figure 1: The sequence $f_n(x) = x^n$ is not equicontinuous.

Consider now

$$f_{n_k} := f_{n(k,k)}, \quad k \geq 1$$

the "diagonal sequence", and remark that $f_{n_k}(x_j) \rightarrow a_j = f(x_j)$ as $k \rightarrow \infty$ for every $j \geq 1$. Hence, $\{f_{n_k}\}_k$ converges to f on D , pointwise.

We claim now that $\{f_{n_k}\}_k$ converges on all of X to some function $f : X \rightarrow \mathbb{R}$, pointwise. Put $g_k := f_{n_k}$ for notational convenience. Fix $x_0 \in X, \epsilon > 0$, and let $\delta_{x_0} > 0$ be such that if $x \in X$ such that $p(x, x_0) < \delta_{x_0}$, $|g_k(x) - g_k(x_0)| < \frac{\epsilon}{3}$. Since D is dense in X , $\exists x_j \in D$ s.t. $p(x_j, x_0) < \delta_{x_0}$. Since $\{g_k(x_j)\}_k$ converges, it is thus Cauchy, and hence for every $k, \ell \geq K$, $|g_k(x_j) - g_\ell(x_j)| < \frac{\epsilon}{3}$. Therefore,

$$|g_k(x_0) - g_\ell(x_0)| \leq |g_k(x_0) - g_k(x_j)| + |g_k(x_j) - g_\ell(x_j)| + |g_\ell(x_j) - g_\ell(x_0)| < \epsilon$$

And thus $\{g_k(x_0)\}_k$ Cauchy as a sequence in \mathbb{R} . Since \mathbb{R} is complete, then $\{g_k(x_0)\}_k$ also converges, to, say, $f(x_0) \in \mathbb{R}$. Since x_0 was arbitrary, this means there is some function $f : X \rightarrow \mathbb{R}$ such that $g_k \rightarrow f$ pointwise on X as we aimed to show. \square

Theorem 31. (Arzelà-Ascoli Theorem): Let X be compact and let $\{f_n\}_{n=1}^\infty \subseteq C(X)$ be uniformly bounded and uniformly equicontinuous. Then, \exists subseq $\{f_{n_k}\}_{k=1}^\infty$ and $f \in C(X)$ s.t. $f_{n_k} \xrightarrow[k \rightarrow \infty]{} f$ in $C(X)$ (i.e. uniformly)

Proof. Since (X, p) is compact, it is thus separable. Also, uniform bounded/equicontinuous implies pointwise bounded/equicontinuous. Therefore, by Arzelà-Ascoli lemma, $\exists f : X \rightarrow \mathbb{R}$ and $\{f_{n_k}\}_{k=1}^\infty$ s.t. $f_{n_k} \rightarrow f$ pointwise in X .

Now let $g_k := f_{n_k}$.

Claim: $\{g_k\}_{k=1}^\infty$ is uniformly Cauchy.⁶

⁶ Cauchy sequence in $(C(X), \|\cdot\|_\infty)$

Fix $\epsilon > 0$. By uniform equicontinuity, $\exists \delta > 0$ s.t.

$$p(x, y) < \delta \implies |f_n(x) - f_n(y)| < \frac{\epsilon}{3} \quad \forall n \in \mathbb{N}.$$

Letting $n = n_k$,

$$p(x, y) < \delta \implies |g_k(x) - g_k(y)| < \epsilon \quad \forall k \in \mathbb{N}.$$

Since X is compact, it is totally bounded, so $\exists \{x_i\}_{i=1}^N$ s.t. $X \subseteq \bigcup_{i=1}^N B^p(x_i, \delta)$.

Moreover, $\forall 1 \leq i \leq N$ fixed, we know $\{g_k(x_i)\}_{k=1}^\infty \subseteq \mathbb{R}$ converges because $\{g_k\}_{k=1}^\infty$ converges pointwise, so $\{g_k(x_i)\}_{k=1}^\infty$ is a Cauchy sequence. So $\exists K_i > 0$ s.t. $\forall k, \ell \geq K_i$,

$$|g_k(x_i) - g_\ell(x_i)| \leq \frac{\epsilon}{3}.$$

Let $K := \max_{1 \leq i \leq N} K_i$. Then, $\forall k, \ell \geq K$, we have

$$|g_k(x_i) - g_\ell(x_i)| \leq \frac{\epsilon}{3} \quad \forall 1 \leq i \leq N.$$

So $\forall x \in X \subseteq \bigcup_{i=1}^N B^p(x_i, \delta)$, $\exists x_i$ s.t. $p(x, x_i) < \delta$, and $\forall k, \ell \geq K$,

$$|g_k(x) - g_\ell(x)| \leq |g_k(x) - g_k(x_i)| + |g_k(x_i) - g_\ell(x_i)| + |g_\ell(x_i) - g_\ell(x)| < \epsilon.$$

This implies $\forall \epsilon > 0, \exists K > 0$ s.t. $\forall k, \ell > K$,

$$\|g_k - g_\ell\|_\infty = \sup_{x \in X} |g_k(x) - g_\ell(x)| < \epsilon,$$

so $\{g_k\}_{k=1}^\infty$ is uniformly Cauchy. Since (X, p) is compact, $C(X)$ is complete, so $\{g_k\}_{k=1}^\infty = \{f_{n_k}\}_{k=1}^\infty$ converges uniformly. Since $f_{n_k} \rightarrow f$ pointwise in X , it must be that $f_{n_k} \rightarrow f$ uniformly, and thus $f \in C(X)$. \square

Remark 32. How do we use the AA theorem? To extract convergent subsequence, which may give us convergence of the original sequence.

Fact: Let $\{f_n\}_{n=1}^\infty \subseteq C(X)$. If $\exists! f$ s.t. for every subsequence, \exists a further subsequence $\{f_{n_{k_j}}\}_{j=1}^\infty$ s.t. $f_{n_{k_j}} \xrightarrow{j \rightarrow \infty} f$ uniformly, then $f_n \xrightarrow{n \rightarrow \infty} f$ uniformly.

Typical Application

- Verify $\{f_n\}$ satisfies hypothesis of AA;
- For every subseq $\{f_{n_k}\}$ also satisfies hypothesis of AA;
- Use AA to extract $\{f_{n_{k_j}}\}_{j=1}^\infty$ s.t. $f_{n_{k_j}} \rightarrow f$ uniformly on X .
- If you can show f is unique, then $f_n \rightarrow f$ in $C(X)$.

Corollary 33. Let (X, p) be a compact metric space. Let $\mathcal{F} \subseteq C(X)$ be uniformly bounded and uniformly equicontinuous. Then, \mathcal{F} is precompact in $(C(X), \|\cdot\|_\infty)$.

Proof. If f is uniformly bounded and uniformly equicontinuous, then by the AA theorem, \forall sequence $\{f_n\}_{n=1}^\infty \subseteq \mathcal{F}$, there is a subseq. $\{f_{n_j}\}_{j=1}^\infty$ and $f \in C(X)$ s.t. $f_{n_j} \rightarrow f$ in $C(X)$. Note, f may not be in \mathcal{F} . So \mathcal{F} is precompact. \square

Example 34. Let $M > 0$, $\mathcal{F} = \{f \in C([a, b]) \cap C^1([a, b]) : \|f\|_\infty + \|f'\|_\infty < M\}$. \mathcal{F} is uniformly bounded and uniformly equicontinuous. So by AA then, for $\{f_n\} \subseteq \mathcal{F}$, $\exists \{f_{n_k}\}_{k=1}^\infty$ s.t. $f_{n_k} \rightarrow f$ uniformly. But, f may not be in $C^1([a, b])$. (So, f may not be in \mathcal{F})

Extra stuff left in the assignment, go back and look at it.

Baire Category Theorem

Definition 35. (Hollow/Nowhere Dense): We say a set E is *hollow* if $\text{Int}(E) = \emptyset$.⁷ We say $E \subseteq X$ *nowhere dense* if its closure is hollow, i.e. $\text{Int}(\overline{E}) = \emptyset$.

⁷ i.e. E contains no nontrivial open sets

Remark 36. E hollow $\iff E^c$ dense in X , since $\text{Int}(E) = \emptyset \iff (\text{Int}(E))^c = \overline{E^c} = X$.

Goal: When can we guarantee that

- a union of hollow sets is hollow?
- an intersection of dense sets is dense?

Theorem 37 (Baire Category Theorem). Let (X, p) be a complete metric space.

1. Let $\{F_n\}_{n=1}^\infty \subseteq X$ be a collection of closed hollow sets. Then $\bigcup_{n=1}^\infty F_n$ is hollow.
2. Let $\{\mathcal{O}_n\}_{n=1}^\infty \subseteq X$ be a collection of open dense sets. Then $\bigcap_{n=1}^\infty \mathcal{O}_n$ is dense.

Proof. (2) \Rightarrow (1) by taking complements and using the previous remark, so we prove only (2).

Claim: Let $G = \bigcap_{n=1}^\infty \mathcal{O}_n$. Then G is dense in X .

Fix $x \in X, r > 0$. $\forall n \in \mathbb{N}, \mathcal{O}_n$ is open and dense, so $\exists y \in \mathcal{O}_n$ and $s > 0$ s.t.

$$B(x, r) \cap \mathcal{O}_n \supseteq B(y, 2s) \supseteq \overline{B(y, s)}.$$

Now we use this fact inductively in n . Let $x_1 \in X, r_1 < \frac{1}{2}$ s.t. $\overline{B(x_1, r_1)} \subseteq B(x, r) \cap \mathcal{O}_1$. Let $x_2 \in X, r_2 < 2^{-2}$ s.t. $\overline{B(x_2, r_2)} \subseteq B(x_1, r_1) \cap \mathcal{O}_2$. Repeating this process, take $x_n \in X, r_n < 2^{-n}$ s.t. $\overline{B(x_n, r_n)} \subseteq B(x_{n-1}, r_{n-1}) \cap \mathcal{O}_n$.

$$\Rightarrow \overline{B(x_1, r_1)} \supseteq \overline{B(x_2, r_2)} \supseteq \cdots \supseteq \overline{B(x_n, r_n)} \supseteq \cdots,$$

and $r_n \rightarrow 0$. Therefore $\{x_n\}_{n=1}^\infty$ is Cauchy, and (X, p) is complete, so $\exists x_0 \in X$ s.t. $x_n \rightarrow x_0$. Thus,

$$x_0 = \bigcap_{n=1}^\infty \overline{B(x_n, r_n)}.$$

Since $x_0 \in \overline{B(x_n, r_n)} \subseteq \mathcal{O}_n \forall n$, and $x_0 \in \overline{B(x_1, r_1)} \subseteq B(x, r) \Rightarrow x_0 \in G \cap B(x, r)$.

$$\Rightarrow G \cap B(x, r) \neq \emptyset \forall x \in X, \forall r > 0.$$

$\Rightarrow G$ is dense in X . □

Another restatement of the Baire Category Theorem is as follows: If (X, p) is complete, the countable union of nowhere dense sets is hollow.

Proof. Let $\{E_n\}_{n=1}^\infty$ be nowhere dense sets. Then by BCT, $\bigcup_{n=1}^\infty \overline{E_n}$ is hollow. It follows that $\bigcup_{n=1}^\infty E_n \subseteq \bigcup_{n=1}^\infty \overline{E_n}$ so $\bigcup_{n=1}^\infty E_n$ is also hollow. □

The main way we will use the Baire Category Theorem is the following:

Corollary 38. Let (X, p) be complete. Suppose $\{F_n\}_{n=1}^\infty$ is a collection of closed sets. If $X = \bigcup_{n=1}^\infty F_n$, then $\exists n_0$ s.t. $\text{Int}(F_{n_0}) \neq \emptyset$.

Proof. If $\nexists n_0$, then F_n is hollow $\forall n$, so by BCT $X = \bigcup_{n=1}^\infty F_n$ is hollow, but this is a contradiction because $X \subseteq X$ is open and nontrivial. □

Baire Category Theorem Application:

Theorem 39. Let $X \subseteq C(X)$ where (X, p) is complete. Suppose \mathcal{F} is pointwise bounded. Then, \exists non-empty open set $\mathcal{O} \subseteq X$ s.t. \mathcal{F} is uniformly bounded on \mathcal{O} , i.e. $\exists M > 0$ s.t.

$$\sup_{f \in \mathcal{F}} \sup_{x \in \mathcal{O}} |f(x)| \leq M$$

Proof. Let $E_n = \{x \in X : |f(x)| \leq n \forall f \in \mathcal{F}\} = \bigcap_{f \in \mathcal{F}} \{x \in X : |f(x)| \leq n\} \Rightarrow E_n$ is closed $\forall n$. Since \mathcal{F} is pointwise bounded, $\forall x \in X, \exists M_x > 0$ s.t. $\sup_{f \in \mathcal{F}} |f(x)| \leq M_x$. Thus, $\forall n$ s.t. $M_x \leq n$, then $x \in E_n$ ($|f(x)| \leq M_x \leq n$). So, $X = \bigcup_{n=1}^{\infty} E_n$ and E_n is closed. By corollary, $\exists n_0$ s.t. $\text{Int}(E_{n_0}) \neq \emptyset$. So $\exists x_0 \in X, r$ s.t. $B^p(x_0, r) \subseteq E_{n_0}$. Letting $\mathcal{O} = B^p(x_0, r)$, we have $\sup_{x \in \mathcal{O}} |f(x)| \leq n_0 \quad \forall f \in \mathcal{F}$. \square

Corollary 40. Let (X, p) be a complete metric space. Suppose $\{F_n\}_{n=1}^{\infty}$ is a collection of closed sets. Then $\bigcup_{n=1}^{\infty} \partial F_n$ is hollow.

Proof. Claim: ∂F_n is hollow $\forall n$. Suppose for contradiction that $\exists n$ s.t. $\text{Int}(\partial F_n) \neq \emptyset$. Then $\exists x_0 \in \partial F_n, r > 0$ s.t. $B^p(x_0, r) \subseteq \partial F_n$. But then,

$$B^p(x_0, r) \cap F_n^c = B^p(x_0, r) \cap \overline{F_n^c} = B^p(x_0, r) \cap (F_n \cup \partial F_n)^c = B^p(x_0, r) \cap \partial F_n^c \cap F_n^c = \emptyset$$

and this contradicts $x_0 \in \partial F_n$ by defn. $\Rightarrow \partial F_n$ is hollow $\forall n$. Furthermore, ∂F_n is closed, since it contains all of its limit points by definition. Thus, by BCT, $\bigcup_{n=1}^{\infty} \partial F_n$ is hollow. \square

Now recall that in general, $\{f_n\}_{n=1}^{\infty} \subseteq C(X)$ and $f_n \rightarrow f$ pointwise, then f is not necessarily continuous.

Theorem 41. Let (X, p) be complete. Let $\{f_n\}_{n=1}^{\infty} \subseteq C(X)$ s.t. $f_n \rightarrow f$ pointwise in X . Then there is a dense subset $D \subseteq X$ where $\{f_n\}_{n=1}^{\infty}$ is pointwise equicontinuous on D and $\forall x_0 \in D, f$ is continuous at x_0 .

Proof. Let $m, n \in \mathbb{N}$. Define

$$E(m, n) = \left\{ x \in X : |f_j(x) - f_k(x)| \leq \frac{1}{m} \quad \forall j, k \geq n \right\} = \bigcap_{j, k \geq n} \underbrace{\left\{ x \in X : |f_j(x) - f_k(x)| \leq \frac{1}{m} \right\}}_{\text{closed, since } f_k, f_j \in C(X)}.$$

So $E(m, n)$ is closed $\forall m, n$. Thus, by the corollary, $\bigcup_{m, n \in \mathbb{N}} \partial E(m, n)$ is hollow. This implies that

$$D := \left(\bigcup_{m, n \in \mathbb{N}} \partial E(m, n) \right)^c = \bigcap_{m, n \in \mathbb{N}} \partial E(m, n)^c \text{ is dense.}$$

Claim 1: If $\exists x \in X, m, n \in \mathbb{N}$ s.t. $x \in D \cap E(m, n)$, then $x \in \text{Int}(E(m, n))$.

If $x \in D$, then

$$x \in \underbrace{\partial E(m, n)^c}_{\text{open}} = \text{Int}(E(m, n)) \cup \text{Ext}(E(m, n)).$$

For the exterior term:

$$\begin{aligned} \text{Ext}(E(m, n)) &= X \setminus (\text{Int}(E(m, n)) \cup \partial E(m, n)) \\ &= X \setminus E(m, n) = E(m, n)^c. \end{aligned}$$

Since we also have $x \in E(m, n)$, this means $x \in \text{Int}(E(m, n))$.

Claim 2: $\{f_n\}_{n=1}^{\infty}$ is equicontinuous on D .

Let $x_0 \in D$ and $\epsilon > 0$. Choose m s.t. $\frac{1}{m} < \frac{\epsilon}{4}$. Since $\{f_n\}_{n=1}^\infty$ converges, $\{f_n(x_0)\}_{n=1}^\infty \subseteq \mathbb{R}$ is a Cauchy sequence. So $\exists N$ s.t. $\forall j, k \geq N$,

$$|f_j(x_0) - f_k(x_0)| \leq \frac{1}{m}.$$

This means $x_0 \in E(m, n) \cap D$, so by Claim 1, $x_0 \in \text{Int}(E(m, n))$. Let $B^p(x_0, r) \subseteq E(m, N)$, so $\forall j, k \geq N, \forall x \in B(x_0, r)$,

$$|f_j(x) - f_k(x)| \leq \frac{1}{m}.$$

Since f_N is continuous at x_0 , $\exists \delta_{x_0} > 0$ (which WLOG we can choose $< r$), s.t. $\forall x \in B^p(x_0, \delta_{x_0})$,

$$|f_N(x) - f_N(x_0)| \leq \frac{1}{m}.$$

So $\forall j \geq N, \forall x \in B^p(x_0, \delta_{x_0})$,

$$|f_j(x) - f_j(x_0)| \leq |f_j(x) - f_N(x)| + |f_N(x) - f_N(x_0)| + |f_N(x_0) - f_j(x_0)| \leq \frac{3}{m} \leq \frac{3\epsilon}{4}.$$

Since this holds $\forall j \geq N$, this implies that $\{f_n\}_{n=1}^\infty$ is equicontinuous at x_0 . Furthermore, $\forall x \in B^p(x_0, \delta_{x_0})$, sending $j \rightarrow \infty$, we obtain that $\forall x \in B^p(x_0, \delta_{x_0})$,

$|f(x) - f(x_0)| \leq \frac{3\epsilon}{4}$, so f is continuous at $x_0 \in D$. \square

Topological Spaces

Separation, Countability, Separability

Continuity and Compactness