



An Introduction to Hadron Spectroscopy from Lattice QCD

Fernando Romero-López

Uni Bern

fernando.romero-lopez@unibe.ch

Lattice Practices 2024
The Cyprus Institute



Goals:

- How does the QCD spectrum look like?
- How can we investigate the QCD spectrum with lattice QCD?
- What are the finite-volume effects in QCD stable particles?
- How to obtain resonance properties from lattice QCD?
- What are the finite-volume effects in scattering states?

Outline

1. The Hadron Spectrum
2. Lattice QCD spectroscopy
3. Finite-volume effects: stable particles
4. Scattering processes and resonances
5. Finite-volume effects: multi-hadron states
6. Selection of recent results

The Hadron Spectrum

Quantum Chromodynamics

Quantum chromodynamics is conceptually simple. Its realization in nature, however, is usually very complex.

Frank Wilczek

Quantum Chromodynamics

Quantum chromodynamics is conceptually simple. Its realization in nature, however, is usually very complex.

Frank Wilczek

$$\mathcal{L}_{QCD} = \sum_i^{N_f} \bar{q}_i \left(D_\mu \gamma^\mu + \textcolor{red}{m}_i \right) q_i + \frac{1}{4g_s^2} G_{\mu\nu}^a G_a^{\mu\nu}$$

Quantum Chromodynamics


Quantum chromodynamics is conceptually simple. Its realization in nature, however, is usually very complex.

Frank Wilczek


$$\mathcal{L}_{QCD} = \sum_i^{N_f} \bar{q}_i \left(D_\mu \gamma^\mu + \textcolor{red}{m_i} \right) q_i + \frac{1}{4g_s^2} G_{\mu\nu}^a G_a^{\mu\nu}$$



The QCD coupling



The QCD coupling




Perturbation theory:

$$\mathcal{O} = \mathcal{O}_0 + \alpha_s \mathcal{O}_1 + \alpha_s^2 \mathcal{O}_2 + \dots$$



Asymptotic freedom

[Gross, Politzer, Wilczek,
Nobel 2004]

The QCD coupling



The QCD coupling



Perturbation theory:

$$\mathcal{O} = \mathcal{O}_0 + \alpha_s \mathcal{O}_1 + \alpha_s^2 \mathcal{O}_2 + \dots$$


Asymptotic freedom

[Gross, Politzer, Wilczek, Nobel 2004]

Hadrons ≠ Confinement

Our understanding of the SM is limited by the complexity of the strong force


- Quarks and gluons carry the strong charge: the so-called “color”



Hadrons ≠ Confinement


Our understanding of the SM is limited by the complexity of the strong force

- Quarks and gluons carry the strong charge: the so-called “color”



Confinement

Quarks and gluons can only be found within colorless composite states. These are called “hadrons”.





source: IFIC

Hadrons ≠ Confinement

Our understanding of the SM is limited by the complexity of the strong force



- Quarks and gluons carry the strong charge: the so-called “color”




Hadrons ≠ Confinement

Our understanding of the SM is limited by the complexity of the strong force

- Quarks and gluons carry the strong charge: the so-called “color”



source: IFIC




The QCD Fock space

- The only stable hadron in Nature is the **proton**
- In Lattice QCD, we (typically) treat QCD in isolation. Several hadrons become stable: π, K, D, p, n, Σ

The QCD Fock space

- The only stable hadron in Nature is the **proton**
 - In Lattice QCD, we (typically) treat QCD in isolation. Several hadrons become stable: π, K, D, p, n, Σ
- QCD stable hadrons, e.g. pseudo-Goldstone bosons




$|\pi\rangle, |K\rangle \in \text{QCD Fock}$

The QCD Fock space


- The only stable hadron in Nature is the **proton**
- In Lattice QCD, we (typically) treat QCD in isolation. Several hadrons become stable: π, K, D, p, n, Σ

► QCD stable hadrons, e.g. pseudo-Goldstone bosons



$|\pi\rangle, |K\rangle \in \text{QCD Fock}$

► Resonances show up in scattering processes



$|\rho\rangle \notin \text{QCD Fock}$

The quark model picture

- Crude classification of the hadron spectrum but still useful for intuition

[Gell-Mann '64 & Zweig '64]

- Construct color singlets combining quarks and antiquarks

$$q \rightarrow \mathbf{3} \text{ irrep of } \mathrm{SU}(3)_c$$

$$\bar{q} \rightarrow \bar{\mathbf{3}} \text{ irrep of } \mathrm{SU}(3)_c$$

Meson:

$$\mathbf{3} \otimes \bar{\mathbf{3}} = \boxed{\mathbf{1}} \oplus \mathbf{8}$$

Baryon:

$$\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \boxed{\mathbf{1}} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$$

color singlet

The quark model picture

- Crude classification of the hadron spectrum but still useful for intuition

[Gell-Mann '64 & Zweig '64]

- Construct color singlets combining quarks and antiquarks

Meson: $3 \otimes \bar{3} = \boxed{1} \oplus 8$

$$q \rightarrow \mathbf{3} \text{ irrep of } \mathrm{SU}(3)_c$$

Baryon: $3 \otimes 3 \otimes 3 = \boxed{1} \oplus 8 \oplus 8 \oplus 10$
 color singlet


$$\bar{q} \rightarrow \bar{\mathbf{3}} \text{ irrep of } \mathrm{SU}(3)_c$$

- Use approximate flavor symmetries to classify multiplets

$$m_u \simeq m_d \simeq m_s \quad \mathrm{SU}(3)_f$$

$$3_f \otimes \bar{3}_f = 1_f \oplus 8_f$$

singlet + octet



The quark model picture

- Crude classification of the hadron spectrum but still useful for intuition
[Gell-Mann '64 & Zweig '64]
- Construct color singlets combining quarks and antiquarks

Meson: $3 \otimes \bar{3} = \boxed{1} \oplus 8$

Baryon: $3 \otimes 3 \otimes 3 = \boxed{1} \oplus 8 \oplus 8 \oplus 10$
color singlet

- Use approximate flavor symmetries to classify multiplets

$$m_u \simeq m_d \simeq m_s \quad 3_f \otimes \bar{3}_f = 1_f \oplus 8_f$$


singlet + octet

- Many hadrons do not fit in this picture: “exotics”

► They are not mesons/baryons (e.g. tetraquarks) or cannot be explained through multi-quark objects

$$q \rightarrow \mathbf{3} \text{ irrep of } \mathrm{SU}(3)_c$$


$$\bar{q} \rightarrow \bar{\mathbf{3}} \text{ irrep of } \mathrm{SU}(3)_c$$



Low-lying hadron spectrum

Light meson octet
Pseudo-Nambu-Goldstone
bosons from spontaneous chiral
symmetry breaking

$$M_m^2 \propto m_q$$



Low-lying hadron spectrum

heavy singlet meson

Mass generated through the chiral anomaly


$$M_{\eta'}^2 \propto \chi_{\text{topo}}$$

[Witten, Veneziano]

Light meson octet

Pseudo-Nambu-Goldstone bosons from spontaneous chiral symmetry breaking

$$M_m^2 \propto m_q$$



Low-lying hadron spectrum

heavy singlet meson

Mass generated through the chiral anomaly


$$M_{\eta'}^2 \propto \chi_{\text{topo}}$$

[Witten, Veneziano]

Light meson octet


Pseudo-Nambu-Goldstone bosons from spontaneous chiral symmetry breaking

$$M_m^2 \propto m_q$$





An experimental conundrum

A growing hadron spectrum still requires first principles understanding





An experimental conundrum

A growing hadron spectrum still requires first principles understanding



An experimental conundrum

A growing hadron spectrum still requires first principles understanding



Lattice QCD spectroscopy

Energies from correlations

- We measure **energy levels** and **matrix elements**: "Spectral decomposition"

$$\begin{aligned} C(t) &= \langle \mathcal{O}^\dagger(t) \mathcal{O}(0) \rangle = \sum_n \langle 0 | \mathcal{O}^\dagger(t) | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \langle 0 | e^{Ht} \mathcal{O}^\dagger(0) e^{-Ht} | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \left| \langle 0 | \mathcal{O}(0) | n \rangle \right|^2 e^{-E_n t} \end{aligned}$$

Energies from correlations

- We measure **energy levels** and **matrix elements**: "Spectral decomposition"

$$\begin{aligned} C(t) &= \langle \mathcal{O}^\dagger(t) \mathcal{O}(0) \rangle = \sum_n \langle 0 | \mathcal{O}^\dagger(t) | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \langle 0 | e^{Ht} \mathcal{O}^\dagger(0) e^{-Ht} | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \left| \langle 0 | \mathcal{O}(0) | n \rangle \right|^2 e^{-E_n t} \end{aligned}$$

Ground state


$$\lim_{t \rightarrow \infty} C(t) = A_0 e^{-E_0 t}$$

Energies from correlations

- We measure **energy levels** and **matrix elements**: "Spectral decomposition"


$$\begin{aligned} C(t) &= \langle \mathcal{O}^\dagger(t) \mathcal{O}(0) \rangle = \sum_n \langle 0 | \mathcal{O}^\dagger(t) | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \langle 0 | e^{Ht} \mathcal{O}^\dagger(0) e^{-Ht} | n \rangle \langle n | \mathcal{O}(0) | 0 \rangle \\ &= \sum_n \left| \langle 0 | \mathcal{O}(0) | n \rangle \right|^2 e^{-E_n t} \end{aligned}$$

Ground state

$$\lim_{t \rightarrow \infty} C(t) = A_0 e^{-E_0 t}$$


QCD spectrum from LQCD

- Consider isospin-symmetric QCD ($m_u = m_d$)
- Neglect QED effects in hadrons
- Need a 3 inputs to fix quark masses
 - ▶ Fix light and strange quark mass
 - ▶ Fix lattice spacing
- Reproduce the lowest-lying hadrons!




[<https://www.science.org/doi/full/10.1126/science.1163233>]

[BMW collaboration, 2008]

QCD spectrum from LQCD

- More recently: add QED and $m_u \neq m_d$
- Can reproduce neutron/proton mass difference
- More precise than some experimental results






[BMW collaboration (2015)]

[<https://www.science.org/doi/full/10.1126/science.1257050>]

Towards Nuclear Physics

- Lattice QCD can in principle compute properties of nuclei. In practice, it is still very preliminary.
- Signal-to-noise problem grows rapidly with the baryon number: hard to control excited states.



[Beane et al (NPLQCD), 2012]

- Computational cost of Wick contractions grows (naively) factorially
contractions $\sim (3A)!$

Beyond the ground state

- ▶ Compute matrix of Euclidean correlation functions using operators with the same quantum numbers


$$C_{ij}(t) = \langle \mathcal{O}_i(t)\mathcal{O}_j^\dagger(0) \rangle$$

$$C_{ij}(t) = \langle 0 | \mathcal{O}_i | n \rangle \langle n | \mathcal{O}_j | 0 \rangle^* e^{-E_n t}$$

- ▶ Variational techniques
(Generalized EigenValue Problem, GEVP)




Nuclear Physics B
Volume 339, Issue 1, 23 July 1990, Pages 222-252




How to calculate the elastic scattering matrix in two-dimensional quantum field theories by numerical simulation

Martin Lüscher, Ulli Wolff^{a,b}




Hadron Spectrum



[HadSpec Collaboration, arXiv:1104.5152]

Hadron Spectrum



[HadSpec Collaboration, arXiv:1104.5152]

Finite-volume effects: stable particles

Stable particle in a box

- We want to compute the mass of a particle in lattice QCD.


- ▶ Lattice QCD calculations are performed in a finite volume.
- ▶ What are the finite-volume effects?

- Qualitative picture in Quantum Mechanics:

$$\psi(x) = \frac{1}{\sqrt{\mathcal{N}}} \exp(-m|x|) \quad m = \langle \psi | H | \psi \rangle$$

$$m(L) = \langle \psi_L | H | \psi_L \rangle \sim m \times \frac{1}{\mathcal{N}} \int_{-L/2}^{L/2} dx \exp(-2m x^2) = m + O(\exp(-mL))$$

- ▶ Finite volume is like a infinite potential well



Stable massive particle only “sees” boundary exponentially

Finite volume QFT

- In a QFT is formulated in finite-volume, loop integrals become sums:

$$\int \frac{d^4 k}{(2\pi)^4} \rightarrow \int \frac{dk^0}{2\pi} \frac{1}{L^3} \sum_{\vec{k} \in (2\pi/L)\vec{n}}$$

- Mass of a particle given by the self-energy:

$$m^2 = m_0^2 + \Sigma(m^2) \xrightarrow{\text{Finite Volume}} m^2(L) = m_0^2 + \Sigma_L(m^2(L))$$


- Mass of a particle given by the self-energy:

$$m(L) - m = \frac{1}{2m} [\Sigma_L(m^2) - \Sigma(m^2)]$$

Example: ϕ^4 theory

- Self-energy given by tad-pole diagram:
- Finite-volume effects given as:

$$\mathcal{L}_I = -\frac{\lambda}{4!} \phi^4$$



$$\Sigma_L - \Sigma = \int \frac{2k^0}{2\pi} \left(\frac{1}{L^3} \sum_{\vec{n}} - \int \frac{d^3 k}{(2\pi)^3} \right) \frac{-\lambda}{k^2 - m^2} \simeq \text{const} \times m \frac{e^{-mL}}{L}$$

EXERCISE Demonstrate that the finite-volume corrections are: $\Delta m(L) \propto \frac{K_1(mL)}{mL} \approx \frac{e^{-mL}}{mL}$

1. Perform the temporal integral in the self energy

2. Use the first term in the Poisson summation formula: $\left(\frac{1}{L^3} \sum_{\vec{n}} - \int \frac{d^3 p}{(2\pi)^3} \right) f(p^2) = \sum_{\vec{n} \neq \vec{0}} \int \frac{d^3 p}{(2\pi)^3} f(p^2) e^{i(\vec{n} \cdot \vec{p})L}$

Volume dependence

Non-perturbative volume dependence of stable particles

Volume Dependence of the Energy Spectrum in Massive Quantum Field Theories

I. Stable Particle States

M. Lüscher


Theory Division, Deutsches Elektronen-Synchrotron DESY, D-2000 Hamburg 52,
Federal Republic of Germany

$$\Delta m = -\frac{3}{16\pi m^2 L} \left\{ \lambda_3^2 e^{-\frac{\sqrt{3}}{2} mL} + \frac{m}{\pi} \int_{-\infty}^{\infty} dy e^{-\sqrt{m^2+y^2}L} F(iy) + O\left(e^{-\bar{m}L}\right) \right\}$$



Summary (1)

- Spectroscopy of stable particles leads only to exponentially suppressed effects
- Form of the FV effects is calculable in QFT
- Motivates choice: $M_\pi L > 4$, $\exp(-4) \sim 2\%$ error
- Example: Volume dependence in phi^4 theory
[FRL, Rusetsky, Urbach, 1806.02367]
- Related ideas on the volume dependence of stable particles:
 - ▶ Volume dependence in Chiral Perturbation Theory
[Colangelo et al, 0311023 & 0503014]
 - ▶ Volume dependence of bound states.
[Hansen, Sharpe arXiv:1609.04317]
 - ▶ Finite-volume effects in g-2
[Hansen, Patella arXiv:2004.03935]



Scattering processes and resonances

Scattering basics

- The “Scattering Matrix” is a unitary operator that connects asymptotic states

$$S_{ab}(E) \equiv \langle \text{out} | \hat{S} | \text{in} \rangle$$

- It is related to the scattering amplitude as:

$$\langle \text{out} | (\hat{S} - 1) | \text{in} \rangle = (2\pi)^4 \delta^{(4)}(P_{\text{in}} - P_{\text{out}}) i\mathcal{M}(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2)$$

Scattering basics

- The “Scattering Matrix” is a unitary operator that connects asymptotic states

$$S_{ab}(E) \equiv \langle \text{out} | \hat{S} | \text{in} \rangle$$

- It is related to the scattering amplitude as:

$$\langle \text{out} | (\hat{S} - 1) | \text{in} \rangle = (2\pi)^4 \delta^{(4)}(P_{\text{in}} - P_{\text{out}}) i \mathcal{M}(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2)$$

- Unitarity imposes important constraints: $\hat{S} \hat{S}^\dagger = 1$


$$\mathcal{M}_2(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2) - \mathcal{M}_2^*(\mathbf{p}_1, \mathbf{p}_2; \mathbf{k}_1, \mathbf{k}_2) =$$

$$\frac{i}{2} \int \frac{d^3 q_1 d^3 q_2}{(2\pi)^6 4\omega(q_1)\omega(q_2)} \mathcal{M}_2(\mathbf{k}_1, \mathbf{k}_2; \mathbf{q}_1, \mathbf{q}_2) \mathcal{M}_2^*(\mathbf{p}_1, \mathbf{p}_2; \mathbf{q}_1, \mathbf{q}_2) \times (2\pi)^4 \delta^{(4)}(k_1 + k_2 - q_1 - q_2)$$

Two-particle scattering

- Consider two-hadron scattering in their CM frame
 - ▶ Amplitude depends on two kinematic variables:

$$\mathcal{M}(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2) \equiv \mathcal{M}(s, \cos \theta)$$




Two-particle scattering

- Consider two-hadron scattering in their CM frame

► Amplitude depends on two kinematic variables:

$$\mathcal{M}(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2) \equiv \mathcal{M}(s, \cos \theta)$$



- We typically work in partial waves:


$$\mathcal{M}(s, \cos \theta) = \sum_{\ell} (2\ell + 1) P_{\ell}(\cos \theta) \mathcal{M}_{\ell}(s)$$

Two-particle scattering

- Consider two-hadron scattering in their CM frame

► Amplitude depends on two kinematic variables:

$$\mathcal{M}(\mathbf{k}_1, \mathbf{k}_2; \mathbf{p}_1, \mathbf{p}_2) \equiv \mathcal{M}(s, \cos \theta)$$



- We typically work in partial waves:

$$\mathcal{M}(s, \cos \theta) = \sum_{\ell} (2\ell + 1) P_{\ell}(\cos \theta) \mathcal{M}_{\ell}(s)$$

- Unitarity implies that:

$$\mathcal{M}_{\ell} = \frac{16\pi\sqrt{s}}{k \cot \delta_{\ell} - ik}$$

- Where $k^{2\ell+1} \cot \delta_{\ell}$ is a **meromorphic** function, e.g. simple polynomial or rational function
(holomorphic up to isolated points)



Scattering ≠ Resonances

- Resonances typically show up as enhancements in the cross-section

$$\sigma_\ell \propto |\mathcal{M}_\ell|^2 = (16\pi\sqrt{s})^2 \frac{\sin^2 \delta_\ell}{k}$$

► Maximum when $\delta_\ell = 90^\circ$


► Unitarity bound is $\sin \delta_\ell = 1$



Poles in the complex plane

- The rigorous definition of a hadronic resonance is a pole in the complex plane

pole residue:
a.k.a coupling

$$\mathcal{M}_\ell \sim -\frac{g^2}{s - s_R}$$


mass of
the resonance


$$\sqrt{s_R} = M_R - i \frac{\Gamma}{2}$$

width of
the resonance

Poles in the complex plane

- The rigorous definition of a hadronic resonance is a pole in the complex plane

pole residue:
a.k.a coupling

$$\mathcal{M}_\ell \sim -\frac{g^2}{s - s_R}$$


width of
the resonance

$$\sqrt{s_R} = M_R - i \frac{\Gamma}{2}$$

mass of
the resonance

- Based on the location of the poles, they receive different names

- ▶ Bound states: stable particles, e.g. the deuteron is an NN bound state
- ▶ Resonances: unstable hadrons, e.g. the rho resonance
- ▶ Virtual states: “non-renormalizable QM states”, e.g. “dineutron”




Fig. 1 Naming convention for the poles in the k -plane. The thick red line for positive real valued k marks the physical momenta in the scattering regime

Pole calculation

EXERCISE Compute the pole positions in the complex plane for the following cases

1. $k \cot \delta_0 = -\frac{1}{a_0}$
2. $k \cot \delta_0 = A(k_R^2 - k^2), \quad A \gg 1/k_R$

What kind of poles do these parametrizations lead to?

What is the residue of the pole? What sign does the residue have?

Summary (2)

The S-Matrix contains the physical information of the theory:

$$S_{ab}(E) \equiv \langle \text{out} | \mathcal{S} | \text{in} \rangle$$

Lattice QCD \rightarrow QCD S-matrix


Summary (2)

The S-Matrix contains the physical information of the theory:

$$S_{ab}(E) \equiv \langle \text{out} | \mathcal{S} | \text{in} \rangle$$

Lattice QCD \rightarrow QCD S-matrix

- Resonances as poles in the S-matrix (or scattering amplitude)



$$\sim \frac{g}{E^2 - E_R^2}$$
$$E_R = M_R - i\Gamma/2$$


Summary (2)

The S-Matrix contains the physical information of the theory:

$$S_{ab}(E) \equiv \langle \text{out} | \mathcal{S} | \text{in} \rangle$$


Lattice QCD \rightarrow QCD S-matrix

► Resonances as poles in the S-matrix (or scattering amplitude)



$$\sim \frac{g}{E^2 - E_R^2}$$

$E_R = M_R - i\Gamma/2$




Summary (2)

The S-Matrix contains the physical information of the theory:

$$S_{ab}(E) \equiv \langle \text{out} | \mathcal{S} | \text{in} \rangle$$


Lattice QCD \rightarrow QCD S-matrix

► Resonances as poles in the S-matrix (or scattering amplitude)



$$\sim \frac{g}{E^2 - E_R^2}$$

$E_R = M_R - i\Gamma/2$




Finite-volume effects:
scattering \neq resonances

Scattering amplitudes in LQCD

Experiments

- Asymptotic states
- Direct access to cross sections




[Protopopescu et al, PRD7 1973]

Scattering amplitudes in LQCD

Experiments

- Asymptotic states
- Direct access to cross sections




[Protopopescu et al, PRD7 1973]

Lattice QCD

- Euclidean time
- Stationary states in a box


$$C(t) = \langle \mathcal{O}(t)\mathcal{O}^\dagger(0) \rangle = \sum_n \left| \langle 0 | \mathcal{O}(0) | n \rangle \right|^2 e^{-E_n t}$$



Scattering amplitudes in LQCD

Experiments

- Asymptotic states
- Direct access to cross sections




[Protopopescu et al, PRD7 1973]

Lattice QCD

- Euclidean time
- Stationary states in a box

$$C(t) = \langle \mathcal{O}(t)\mathcal{O}^\dagger(0) \rangle = \sum_n \left| \langle 0 | \mathcal{O}(0) | n \rangle \right|^2 e^{-E_n t}$$




Finite-volume formalism
[Lüscher, 89']

Finite-volume systems

Free scalar particles in finite volume
with periodic boundaries

$$\psi(\vec{x}) = \psi(\vec{x} + \vec{n}L)$$




$$\vec{p} = \frac{2\pi}{L}(n_x, n_y, n_z)$$

Two particles: $E = 2\sqrt{m^2 + \frac{4\pi^2}{L^2}\vec{n}^2}$


Finite-volume systems

Free scalar particles in finite volume
with periodic boundaries



Interactions change the spectrum:
it can be treated as a perturbation

$$\psi(\vec{x}) = \psi(\vec{x} + \vec{n}L)$$



Ground state to leading order

$$E_2 - 2m = \langle \phi(\vec{0})\phi(\vec{0}) | \mathbf{H}_I | \phi(\vec{0})\phi(\vec{0}) \rangle$$

$$\Delta E_2 = \frac{\mathcal{M}_2(E=2m)}{8m^2L^3} + O(L^{-4})$$


[Huang, Yang, 1958]

$$\vec{p} = \frac{2\pi}{L}(n_x, n_y, n_z)$$

$$\text{Two particles: } E = 2\sqrt{m^2 + \frac{4\pi^2}{L^2}\vec{n}^2}$$


Finite-volume systems

Free scalar particles in finite volume
with periodic boundaries



Interactions change the spectrum:
it can be treated as a perturbation

$$\psi(\vec{x}) = \psi(\vec{x} + \vec{n}L)$$



$$\vec{p} = \frac{2\pi}{L}(n_x, n_y, n_z)$$

Two particles: $E = 2\sqrt{m^2 + \frac{4\pi^2}{L^2}\vec{n}^2}$

Ground state to leading order

$$E_2 - 2m = \langle \phi(\vec{0})\phi(\vec{0}) | H_I | \phi(\vec{0})\phi(\vec{0}) \rangle$$

$$\Delta E_2 = \frac{\mathcal{M}_2(E=2m)}{8m^2L^3} + O(L^{-4})$$

[Huang, Yang, 1958]

The energy shift of the two-particle ground state
is related to the $2 \rightarrow 2$ scattering amplitude


Finite-volume systems

Free scalar particles in finite volume
with periodic boundaries



Interactions change the spectrum:
it can be treated as a perturbation

$$\psi(\vec{x}) = \psi(\vec{x} + \vec{n}L)$$



In general a problem of
Quantum Field Theory
in finite volume

$$\vec{p} = \frac{2\pi}{L}(n_x, n_y, n_z)$$

Two particles: $E = 2\sqrt{m^2 + \frac{4\pi^2}{L^2}\vec{n}^2}$

and state to leading order

$$-2m = \langle \phi(\vec{0})\phi(\vec{0}) | H_I | \phi(\vec{0})\phi(\vec{0}) \rangle$$


$$\Delta E_2 = \frac{\mathcal{M}_2(E = 2m)}{8m^2 L^3} + O(L^{-4})$$

[Huang, Yang, 1958]

The energy shift of the two-particle ground state
is related to the $2 \rightarrow 2$ scattering amplitude

Key insight

Volume dependence of finite-volume energy states contains scattering information




1/L expansions

- Perturbative expansions are useful for
[Lüscher, 89']
- ▶ Weakly interacting systems (no nearby bound states or resonances)
- ▶ Just obtaining the scattering length

$$\Delta E_2 = E_2 - 2m = \frac{4\pi a_0}{m L^3} \left\{ 1 + c_1 \left(\frac{a_0}{L} \right) + c_2 \left(\frac{a_0}{L} \right)^2 \right\} + \mathcal{O}(L^{-6})$$

$(c_1 = 2.837, c_2 = 6.375)$



General approach

- In order to derive the full relation, consider the finite-volume correlator:


$$C_L(E, \vec{P}) = \int e^{iPx} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle =$$

[à la Kim, Sachrajda, Sharpe]

General approach

- In order to derive the full relation, consider the finite-volume correlator:

Skeleton expansion


$$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{---} + \text{---} + \text{---} + \dots$$


[à la Kim, Sachrajda, Sharpe]

General approach

- In order to derive the full relation, consider the finite-volume correlator:

$$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion} + \sum_{\vec{k}} \text{Finite-volume sums}$$



[à la Kim, Sachrajda, Sharpe]

General approach

- In order to derive the full relation, consider the finite-volume correlator:

$$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion} + \dots$$

[à la Kim, Sachrajda, Sharpe]

$\sum_{\vec{k}}$
Finite-volume sums

$$\mathcal{B}_2 = \text{Feynman diagrams} + \dots$$

General approach

- In order to derive the full relation, consider the finite-volume correlator:

$$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion} + \dots$$

[à la Kim, Sachrajda, Sharpe]

Only exponentially small effects in L

$\sum_{\vec{k}}$

Bethe-Salpeter Kernels

Finite-volume sums

$\mathcal{B}_2 = \dots$

$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion} + \dots$

[à la Kim, Sachrajda, Sharpe]

Only exponentially small effects in L

$\sum_{\vec{k}}$

Bethe-Salpeter Kernels

Finite-volume sums

$\mathcal{B}_2 = \dots$

General approach

- In order to derive the full relation, consider the finite-volume correlator:

$$C_L(E, \vec{P}) = \int e^{iP_x} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion}$$

[à la Kim, Sachrajda, Sharpe]

$$\sum_{\vec{k}} \rightarrow \int d^3k + \left[\sum_{\vec{k}} - \int d^3k \right]$$

Only exponentially small effects in L

$$\text{Bethe-Salpeter Kernels}$$
$$B_2 = \text{Diagram} + \text{Diagram} + \text{Diagram} + \dots$$

Finite-volume sums

General approach

- In order to derive the full relation, consider the finite-volume correlator:

$$C_L(E, \vec{P}) = \int e^{iPx} \langle \mathcal{O}(x) | \mathcal{O}(0) \rangle = \text{Skeleton expansion}$$


[à la Kim, Sachrajda, Sharpe]

$$\begin{aligned} &= \text{Diagram with two O's} + \text{Diagram with one O and one B}_2 + \text{Diagram with two B}_2's + \dots \\ &\quad \downarrow \\ &\quad \text{Bethe-Salpeter Kernels} \\ &\quad \text{Only exponentially small effects in } L \quad \text{B} = \text{Diagram} + \dots \\ &\quad \sum_{\vec{k}} \rightarrow \int d^3k + \left[\sum_{\vec{k}} - \int d^3k \right] \\ &\quad \text{Finite-volume sums} \end{aligned}$$

1. Separation of finite-volume effects
2. Resummation of diagrams

General approach

- In order to derive the full relation, consider the finite-volume correlator:



General approach


$$C_L(E, \vec{P}) = \text{some algebra ...} = C_\infty(E, \vec{P}) + A^\dagger \frac{1}{\mathcal{K}_2 + F^{-1}} A + O(e^{-mL})$$

General approach

$$C_L(E, \vec{P}) = \text{some algebra ...} = C_\infty(E, \vec{P}) + A^\dagger \frac{1}{\mathcal{K}_2 + F^{-1}} A + O(e^{-mL})$$

K-matrix parametrized
in terms of phase shift

$$\mathcal{K}_2^\ell = \frac{16\pi\sqrt{s}}{q^{2\ell+1} \cot \delta_\ell}$$



General approach

$$C_L(E, \vec{P}) = \text{some algebra ...} = C_\infty(E, \vec{P}) + A^\dagger - \frac{1}{\mathcal{K}_2 + F^{-1}} A + O(e^{-mL})$$

K-matrix parametrized
in terms of phase shift

$$\mathcal{K}_2^\ell = \frac{16\pi\sqrt{s}}{q^{2\ell+1} \cot \delta_\ell}$$

Finite-volume states appear
when the correlation function
has a pole

General approach

$$C_L(E, \vec{P}) = \text{some algebra ...} = C_\infty(E, \vec{P}) + A^\dagger - \frac{1}{\mathcal{K}_2 + F^{-1}} A + O(e^{-mL})$$

K-matrix parametrized
in terms of phase shift

$$\mathcal{K}_2^\ell = \frac{16\pi\sqrt{s}}{q^{2\ell+1} \cot \delta_\ell}$$

Finite-volume states appear
when the correlation function
has a pole

! It holds below $E_{cm} < 4m$

Two-particle Quantization Condition

$$\det \left[\mathcal{K}_2(E_n) + F^{-1}(E_n, \vec{P}, L) \right] = 0$$

Scattering
K-Matrix

Known kinematic
function

"QC2"

The quantization condition

- Indirect connection between the spectrum and the two-particle scattering amplitude [Lüscher 89']

Two-particle Quantization Condition

$$\det_{\ell m} \left[\mathcal{K}_2(E_n) + F^{-1}(E_n, \vec{P}, L) \right] \Big|_{E=E_n} = 0$$

Scattering K-Matrix Known kinematic function

"QC2"

The quantization condition

- Indirect connection between the spectrum and the two-particle scattering amplitude [Lüscher 89']

K-matrix parametrized in terms of phase shift

$$\mathcal{K}_2^\ell = \frac{16\pi\sqrt{s}}{q^{2\ell+1} \cot \delta_\ell}$$

Two-particle Quantization Condition


$$\det_{\ell m} \left[\mathcal{K}_2(E_n) + F^{-1}(E_n, \vec{P}, L) \right] \Big|_{E=E_n} = 0$$

Scattering K-Matrix Known kinematic function

"QC2"

The quantization condition

- Indirect connection between the spectrum and the two-particle scattering amplitude [Lüscher 89']



Example application

Two pions in s-wave

$$\mathcal{K}_2^{s\text{-wave}}(E_n) = \frac{-1}{F_{00}(E_n, \vec{P}, L)}$$

one
energy
level



a phase
shift
point



Example application

Two pions in s-wave

$$\mathcal{K}_2^{s\text{-wave}}(E_n) = \frac{-1}{F_{00}(E_n, \vec{P}, L)}$$

one
energy
level

a phase
shift
point





Example application

Two pions in s-wave

$$\mathcal{K}_2^{s-wave}(E_n) = \frac{-1}{F_{00}(E_n, \vec{P}, L)}$$

one
energy
level

a phase shift point





Example application

Two pions in s-wave

$$\mathcal{K}_2^{s\text{-wave}}(E_n) = \frac{-1}{F_{00}(E_n, \vec{P}, L)}$$

one
energy
level

a phase
shift
point





Example application

Two pions in s-wave

$$\mathcal{K}_2^{s-wave}(E_n) = \frac{-1}{F_{00}(E_n, \vec{P}, L)}$$

one
energy
level

a phase shift point



Exercise

EXERCISE Code up the quantization condition and solve it for a system with:

$$mL = 4, \quad \frac{k}{m} \cot \delta_0 = -\frac{1}{ma_0}, \quad ma_0 = 0.1$$

You can use an easy implementation of the zeta function, and this form of the QC2:

$$\mathcal{Z}_0(k^2) = \lim_{\Lambda \rightarrow \infty} \sum_{\vec{n} \in \mathbb{Z}}^{| \vec{n} | < \Lambda} \frac{1}{\vec{n}^2 - (\frac{kL}{2\pi})^2} - 4\pi\Lambda$$

$$k \cot \delta_0(k^2) = \frac{1}{\pi L} \mathcal{Z}_0(k^2, L)$$

What is the ground-state energy of the system in the frame $\vec{P} = 0$?


How does it compare to the 1/L expansion?

Solving the QC2

Leading-partial wave approximation:

$$\text{QC2: } \frac{1}{\mathcal{K}_2^{\ell=0}(k^2)} = -F_0(k^2, L)$$

$$\mathcal{K}_2^{\ell=0} = \frac{16\pi\sqrt{s}}{k \cot \delta_0}$$



Summary (3)

- Volume dependence of multi-hadron states contains information about scattering amplitude
- Energy shift to the non-interaction theory can be related to scattering parameters

$$\Delta E_2 = E_2 - 2m = \frac{4\pi a_0}{mL^3} \left\{ 1 + c_1 \left(\frac{a_0}{L} \right) + c_2 \left(\frac{a_0}{L} \right)^2 \right\} + \mathcal{O}(L^{-6})$$

- Two-body quantization condition relates the spectrum to amplitude in generic systems

$$k \cot \delta_0(k^2) = \frac{1}{\pi L} \mathcal{Z}_0(k^2, L)$$

Selection of recent results

The $\Lambda(1405)$

$$\begin{pmatrix} \pi\Sigma \rightarrow \pi\Sigma & \pi\Sigma \rightarrow Kp \\ Kp \rightarrow \pi\Sigma & Kp \rightarrow Kp \end{pmatrix}$$

$$\det_{\ell m} \left[\begin{pmatrix} \tilde{K}_{\pi\Sigma \rightarrow \pi\Sigma} & \tilde{K}_{\pi\Sigma \rightarrow KN} \\ \tilde{K}_{KN \rightarrow \pi\Sigma} & \tilde{K}_{KN \rightarrow KN} \end{pmatrix} + \begin{pmatrix} F_{\pi\Sigma}^{-1}(E_n, \vec{P}, L) & 0 \\ 0 & F_{KN}^{-1}(E_n, \vec{P}, L) \end{pmatrix} \right] = 0$$

Multi-channel K-Matrix

Zeta function

$\Lambda(1405) 1/2^-$

$I(J^P) = 0(\frac{1}{2}^-)$ Status: ****




$\Lambda(1380) 1/2^-$


$J^P = \frac{1}{2}^-$ Status: **



- **** Existence is certain.
- *** Existence is very likely.
- ** Evidence of existence is fair.
- * Evidence of existence is poor.



The $\Lambda(1405)$



▶ Scattering amplitudes for “preferred” fit
i.e. with lowest AIC = $\chi^2 - 2 \text{ dof}$

▶ Scattering amplitudes for different parametrizations


▶ Pole positions for “preferred” fit

▶ Pole positions for different parametrization
All find two poles!

▶ Lattice QCD energies used in fits


Continuum Limit of H dibaryon

- H dibaryon is a $\Lambda\Lambda$ bound state (uuddss quarks)




Three-hadron amplitudes

Qualitatively more complicated than the two-particle case!



Three-hadron amplitudes


Qualitatively more complicated than the two-particle case!



- Three-particle scattering amplitudes can be divergent for specific kinematics.

Three-hadron amplitudes

Qualitatively more complicated than the two-particle case!



- Three-particle scattering amplitudes can be divergent for specific kinematics.
- They depend also on two-to-two interactions.
 - But any separation between “two-particle” and “three-particle” effects is not well-defined

Quantization Condition

Skeleton expansion

$$C_L = \text{Diagram} + \text{Diagram } B_3 + \text{Diagram } B_3 B_3 + \dots + \text{Diagram } B_2 + \text{Diagram } B_2 B_2 + \dots + \text{Diagram } B_2 B_2 + \text{Diagram } B_2 B_2 B_2 + \dots + \dots + \text{Diagram } B_2 B_3 + \text{Diagram } B_2 B_2 B_3 + \dots$$

Quantization Condition

Skeleton expansion

$$C_L = \begin{aligned} & \text{Diagram 1: } \text{Diamond shape with red vertical lines and dashed boxes.} \\ & + \text{Diagram 2: } \text{Diamond shape with circle } B_3 \text{ and red vertical lines.} \\ & + \text{Diagram 3: } \text{Diamond shape with circles } B_3, B_3, B_3 \text{ and red vertical lines.} \\ & + \dots \\ & + \text{Diagram 4: } \text{Diamond shape with circle } B_2 \text{ and red vertical lines.} \\ & + \text{Diagram 5: } \text{Diamond shape with circles } B_2, B_2 \text{ and red vertical lines.} \\ & + \dots \\ & + \text{Diagram 6: } \text{Diamond shape with circles } B_2, B_2 \text{ and blue vertical lines.} \\ & + \text{Diagram 7: } \text{Diamond shape with circles } B_2, B_2, B_2 \text{ and blue vertical lines.} \\ & + \dots \\ & + \text{Diagram 8: } \text{Diamond shape with circles } B_2, B_2, B_2 \text{ and red vertical lines.} \\ & + \dots \\ & + \text{Diagram 9: } \text{Diamond shape with circles } B_2, B_3 \text{ and red vertical lines.} \\ & + \text{Diagram 10: } \text{Diamond shape with circles } B_2, B_3, B_2 \text{ and red vertical lines.} \\ & + \dots \end{aligned}$$

Separation of finite and infinite volume terms:

$$= C_\infty(P) + A_3 \frac{1}{\mathcal{K}_{df,3} + F_3^{-1}} A'_3 + O(e^{-mL})$$

Easier derivation: Blanton, Sharpe [2007.16188]

Quantization Condition

Skeleton expansion

$$C_L = \text{Diagram} + \text{Diagram } B_3 + \text{Diagram } B_3 B_3 + \dots + \text{Diagram } B_2 + \text{Diagram } B_2 B_2 + \dots + \text{Diagram } B_2 B_2 + \text{Diagram } B_2 B_2 B_2 + \dots + \text{Diagram } B_2 B_2 + \text{Diagram } B_2 B_2 B_2 B_2 + \dots + \dots + \text{Diagram } B_2 B_3 + \text{Diagram } B_2 B_3 B_3 + \dots$$

Separation of finite and infinite volume terms:

$$= C_\infty(P) + A_3 \frac{1}{\mathcal{K}_{df,3} + F_3^{-1}} A'_3 + O(e^{-mL})$$



Three-particle Quantization Condition
for identical scalars with G-parity

$$\det [\mathcal{K}_3(E) + F_3^{-1}(E, \vec{P}, L)] = 0$$

"QC3"

Easier derivation: Blanton, Sharpe [2007.16188]

Quantization Condition

Skeleton expansion

$$C_L = \text{Diagram with one loop} + \text{Diagram with two loops (B}_3\text{)} + \text{Diagram with three loops (B}_3\text{)} + \dots \\ + \text{Diagram with one loop (B}_2\text{)} + \text{Diagram with two loops (B}_2\text{)} + \dots \\ + \text{Diagram with one loop (B}_2\text{), one blue vertical line} + \text{Diagram with two loops (B}_2\text{), one blue vertical line} + \dots \\ + \text{Diagram with one loop (B}_2\text{), two blue vertical lines} + \text{Diagram with two loops (B}_2\text{), two blue vertical lines} + \dots \\ + \dots \\ + \text{Diagram with one loop (B}_2\text{), one red vertical line} + \text{Diagram with two loops (B}_2\text{), one red vertical line} + \dots$$

Separation of finite and infinite volume terms:

$$= C_\infty(P) + A_3 \frac{1}{\mathcal{K}_{df,3} + F_3^{-1}} A'_3 + O(e^{-mL})$$



Three-particle Quantization Condition
for identical scalars with G-parity

$$\det \left[\mathcal{K}_3(E) + F_3^{-1}(E, \vec{P}, L) \right] = 0$$

“QC3”

Easier derivation: Blanton, Sharpe [2007.16188]

“Formally” similar to the two-particle case

Quantization Condition

Skeleton expansion

$$C_L = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{n! m! k!} \frac{1}{(B_2)^n (B_3)^m} \frac{1}{(B_2)^k} + \dots$$

Easier derivation: Blanton, Sharpe [2007.16188]

Separation of finite and infinite volume terms:

$$= C_\infty(P) + A_3 \frac{1}{\mathcal{K}_{df,3} + F_3^{-1}} A'_3 + O(e^{-mL})$$



Three-particle Quantization Condition
for identical scalars with G-parity


$$\det \left[\mathcal{K}_3(E) + F_3^{-1}(E, \vec{P}, L) \right] = 0$$

“QC3”

“Formally” similar to the two-particle case

Three-hadron formalism

2π and 3π
Spectrum




1. Determine \mathcal{K}_2 and $\mathcal{K}_{df,3}$ from the two and three-pion spectrum

Hansen, Sharpe [arXiv:1408.5933]

Three-hadron formalism

2π and 3π
Spectrum




$$\det [\mathcal{K}_2 + F_2^{-1}] = 0$$

1. Determine \mathcal{K}_2 and $\mathcal{K}_{df,3}$ from the two and three-pion spectrum

Hansen, Sharpe [arXiv:1408.5933]


Three-hadron formalism



1. Determine \mathcal{K}_2 and $\mathcal{K}_{df,3}$ from the two and three-pion spectrum

Hansen, Sharpe [arXiv:1408.5933]

Three-hadron formalism



1. Determine \mathcal{K}_2 and $\mathcal{K}_{df,3}$ from the two and three-pion spectrum

Hansen, Sharpe [arXiv:1408.5933]

2. Solve integral equations to obtain The physical three-to-three amplitude

Hansen, Sharpe [arXiv:1504.04248]

