

# Algebraic geometry 1

## Exercise sheet 12

Solutions by: Eric Rudolph and David Čadež

22. Januar 2024

### Exercise 3.

1. We want to argue that all local rings  $\mathcal{O}_{X,x}$  are regular by showing that their maximal is (non-zero) and principal.

This basically follows immediately from writing down explicitly how the primes in the local rings look like and of course we use that their Krull dimension is one. This shows smoothness, because we have for all the local rings  $A$  and their maximal ideals we have

$$\dim_k(m/m^2) = 1 = \dim(A)$$

Since  $\dim(Y) = 1$ , a theorem from class implies that  $Y = \text{Spec}(A)$  is normal and hence  $A$  is normal.

Since  $X$  is smooth, all local rings are UFDs. So in this exercise we will show that locally factorial is not affine-local, i.e. that  $\mathcal{O}_{X,x}$  UFD does not imply  $\mathcal{O}_X(\text{Spec}(A))$  for an affine  $\text{Spec}(A)$ .

2. We can write every  $a \in A$  as

$$a = c_0 + c_1 a_1(x) + c_2 y a_2(x),$$

so as a polynomial of degree 1 over  $R$ . Since the sum of two integral elements is integral again (StacksProject) it is enough to show that  $c_2 y a_2(x)$  is integral over  $R$ . This is clearly true since  $(c_2 y a_2(x))^2 \in R$ .

3. We need to check that zero gets mapped to zero. This follows from

$$(-y)^2 - x^3 + x = y^2 - x^3 + x$$

4. Write an element  $a(x, y) \in A$  as

$$a(x, y) = c + a_1(x) + y a_2(x).$$

We compute

$$N(a) = a(x, y)a(x, -y) = (c + a_1(x))^2 - (y a_2(x))^2 = (c + a_1(x))^2 - (x^3 - x)a_2(x)^2 \in R.$$

Multiplicativity of  $N$  follows from the properties of ring maps.

5. Take  $a \in A^*$ . Then  $N(a) \in k[x]^* = k^*$  and since  $\deg(a\sigma(a)) \leq 1$  implies  $\deg(a) \leq 1$  this implies  $a \in k^*$ .

We can check irreducibility using the map  $N$ .

Assume that  $x = fg$  for  $f, g \in A$ . Then

$$x^2 = N(x) = N(fg).$$

Write  $fg = a_1(x)$ . Then by the calculations from above,

$$a_1(x)^2 = x^2 \in R.$$

Therefore  $a_1(x) = \pm x$ , but  $x$  is irreducible in  $R$ , so w.l.o.g.  $f \in A^* = k^*$ .

I guess a similar argument works to show that  $y$  is irreducible.

This shows that  $y^2 = x(x^2 - 1)$  are two decompositions into irreducible elements, showing that  $A$  is not factorial (to me it is not clear that  $x^2 - 1$  is irreducible, but we could just compose it further into its irreducible factors and on the left side there is no  $x$ , but on the right side there is).

First,  $A$  cannot be isomorphic to  $\mathbb{A}_k^1$ , because  $A$  is not a UFD, but  $k[x]$  is.

Let us now try to show that  $A$  can also not be isomorphic to a nontrivial open subset of  $\mathbb{A}_k^1$ . Remember that all opens of  $\mathbb{A}_k^1$  are principal, because  $k[x]$  is principal. So it suffices to show that  $A$  is not isomorphic to a principal open  $D(f) \subset \mathbb{A}_k^1$ . We do this by showing that their corresponding rings are not isomorphic.

This can be seen by comparing units, namely by observing that  $(k[x]_f)^* \subsetneq A^* = k^*$ .