

# Algebraic geometry 1

## Exercise sheet 3

Solutions by: Eric Rudolph and David Čadež

27. Oktober 2023

### Exercise 1.

1. Define

$$\begin{aligned}\pi^{-1} : U &\longrightarrow \pi^{-1}(U) \\ (x_1, \dots, x_n) &\mapsto (x_1, \dots, x_n)[x_1 : \dots : x_n].\end{aligned}$$

This is well-defined, because by definition of  $U$  not all  $x_i$  can be zero at the same time, so  $[x_1 : \dots : x_n]$  is actually a point in projective space. Of course we also have  $(x_1, \dots, x_n)[x_1 : \dots : x_n] \in Z$  for  $(x_1, \dots, x_n) \in U$ , because  $x_i x_j = x_j x_i$  for all  $1 \leq i, j \leq n$ . To see injectivity of  $\pi^{-1}$ , let  $(x_1, \dots, x_n) \in U$  with  $x_j \neq 0$ . Then we have  $y_j \neq 0$ , because if we assume  $x_j \neq 0$  and  $y_j = 0$ , then for some  $y_i \neq 0$  (which exists since  $[y_1 : \dots : y_n]$  is a point in projective space) we have  $0 \neq x_j y_i = x_i y_j = 0$ . Therefore, we can just set  $y_j = 1$ . Then

$$x_i y_j = x_j y_i \implies y_i = \frac{x_i y_j}{x_j} = \frac{x_i}{x_j},$$

showing that all the  $y_i$  are fixed up to a scalar after fixing all the  $x_i$ .