

# Algebraic geometry 1

## Exercise sheet 3

Solutions by: Eric Rudolph and David Čadež

2. November 2023

### Exercise 1.

1. Define

$$\begin{aligned}\pi^{-1} : U &\longrightarrow \pi^{-1}(U) \\ (x_1, \dots, x_n) &\mapsto (x_1, \dots, x_n)[x_1 : \dots : x_n].\end{aligned}$$

This is well-defined, because by definition of  $U$ , not all  $x_i$  can be zero at the same time, so  $[x_1 : \dots : x_n]$  is actually a point in projective space. We also have  $(x_1, \dots, x_n)[x_1 : \dots : x_n] \in Z$  for  $(x_1, \dots, x_n) \in U$ , because  $x_i x_j = x_j x_i$  for all  $1 \leq i, j \leq n$ . To see injectivity of  $\pi^{-1}$ , let  $(x_1, \dots, x_n) \in U$  with  $x_j \neq 0$ . Then we have  $y_j \neq 0$ , because if we assume  $x_j \neq 0$  and  $y_j = 0$ , then for some  $y_i \neq 0$  (which exists since  $[y_1 : \dots : y_n]$  is a point in projective space) we have  $0 \neq x_j y_i = x_i y_j = 0$ . Therefore, we can just set  $y_j = 1$ . Then

$$x_i y_j = x_j y_i \implies y_i = \frac{x_1 y_j}{x_j} = \frac{x_1}{x_j},$$

showing that all the  $y_i$  are fixed up to a scalar after fixing all the  $x_i$ .

2. Define

$$\begin{aligned}\phi : V_i &\rightarrow \mathbb{A}_n^k \\ (x, y) &\mapsto \left(\frac{x_1}{y_i}, \dots, x_i, \dots, \frac{x_n}{y_i}\right),\end{aligned}$$

where the inverse map is given by

$$\begin{aligned}\phi^{-1} : \mathbb{A}_n^k &\rightarrow V_i \\ (x_1, \dots, x_n) &\mapsto (x_1 x_i, \dots, x_i, \dots, x_n x_i)[x_1 : \dots : x_{i-1} : 1 : \dots : x_n].\end{aligned}$$

**Exercise 2.** The part which to me seemed the hardest was to calculate the closure of  $\pi^{-1}(Y \setminus \{t\})$  in  $Z$ . So to for that we look at  $\pi^{-1}(Y)$  and decompose

it into irreducible components. Or actually we first cover it  $V_i$  and then look at them inside each  $V_i$ . In these two cases it decomposed into nice irreducible components, one of which is the blow-up and the other whole  $\pi^{-1}(t)$ .

1. Let  $Y = V(x_1^2 - x_2^3) \subseteq \mathbb{A}^2(k)$ . We look at

$$\pi^{-1}(Y) = \{((x_1, x_2), [y_1 : y_2]) \mid x_1 y_2 = x_2 y_1, x_1^2 - x_2^3 = 0\}. \quad (1)$$

We can cover it with  $V_i$  ( $i = 1, 2$ ). Lets first look inside  $V_1 \cong \mathbb{A}^2(k) \times \mathbb{A}^1(k)$ , where  $y_1 = 1$ . Equations then become  $x_1 y_2 = x_2$  and  $x_1^2(1 - x_1 y_2^3) = 0$ . The latter equation can be decomposed, so we get two closed subsets:

- $\{x_1 = x_2 = 0\} \subseteq V_1 \subseteq \mathbb{A}^2(k) \times \mathbb{P}^1(k)$
- $\{x_1 y_2^3 - 1 = 0, x_2 y_2^2 - 1 = 0\} \subseteq V_1 \subseteq \mathbb{A}^2(k) \times \mathbb{P}^1(k)$

First one lies in  $\pi^{-1}(t)$ , so  $\{x_1 = x_2 = 0\} \cap \pi^{-1}(Y \setminus \{t\}) = \emptyset$ . Therefore

$$\pi^{-1}(Y \setminus \{t\}) \cap V_1 = \{x_1 y_2^3 - 1 = 0, x_2 y_2^2 - 1 = 0, x_1 \neq 0, x_2 \neq 0\} \cap V_1 \quad (2)$$

Therefore taking the closure inside  $V_1$ :

$$\text{BL}_t(Y) \cap V_1 = \{x_1 y_2^3 - 1 = 0, x_2 y_2^2 - 1 = 0\} \cap V_1. \quad (3)$$

Before definition a morphism  $\text{BL}_t(Y) \rightarrow \mathbb{A}^1(k)$ , lets look at  $\text{BL}_t(Y) \cap V_2$ . Similar as before we set  $y_2 = 1$  and get that  $\pi^{-1}(Y) \cap V_2$  is made up of two closed subsets

- $\{x_1 = x_2 = 0\} \subseteq V_2 \subseteq \mathbb{A}^2(k) \times \mathbb{P}^1(k)$
- $\{x_1 = y_1^3, x_2 = y_1^2\} \subseteq V_2 \subseteq \mathbb{A}^2(k) \times \mathbb{P}^1(k)$

This intersection contains more information, since it also contains  $\text{BL}_t(Y) \cap \pi^{-1}(t)$ .

Define a morphism  $\phi: \text{BL}_t(Y) \rightarrow \mathbb{A}^2(k)$

- 2.

#### Exercise 4.

1. Lets first prove that  $V_U$  are stable under intersections:

**Claim.** Take  $U, W \subseteq X$  open subsets. Then  $V_{U \cap W} = V_U \cap V_W$ .

**Proof of claim.** Inclusion  $V_{U \cap W} \subseteq V_U \cap V_W$  is clear.

For the other inclusion take  $Z \in V_U \cap V_W$ . By definition  $Z \cap U \neq \emptyset$  and  $Z \cap V \neq \emptyset$ . Suppose  $Z \cap (U \cap V) = \emptyset$ . Then  $(Z \cap U)^c \cup (Z \cap V)^c = X$ . But since  $Z$  is irreducible, and is covered by  $U^c \cup V^c$ , we must have (WLOG)  $Z \subseteq U^c$ . That is in contradiction with  $Z \cap U \neq \emptyset$ .  $\square$ (of claim)

It also behaves well under unions:

$$\begin{aligned}
V_{U \cup W} &= \{Z \text{ cl. irred.} \mid Z \cap (U \cup W) \neq \emptyset\} \\
&= \{Z \text{ cl. irred.} \mid (Z \cap U) \neq \emptyset \text{ or } (Z \cap W) \neq \emptyset\} \\
&= \{Z \text{ cl. irred.} \mid (Z \cap U) \neq \emptyset\} \cup \{Z \text{ cl. irred.} \mid (Z \cap W) \neq \emptyset\} \\
&= V_U \cup V_W
\end{aligned}$$

and practically same argument applies to infinite unions.

Therefore every open subset of  $X^{\text{sob}}$  can be written as  $V_U$  for some open  $U \subseteq X$  (in general it could've been just a base of topology, but this shows it is the whole topology).

**Claim.** If  $V_{U_1} = V_{U_2}$  then  $U_1 = U_2$ .

**Proof of claim.** Suppose  $x \in U_1 \setminus U_2$ , then  $\overline{\{x\}} \in V_{U_1} \setminus V_{U_2}$ . This proves the claim.  $\square$ (of claim) Next claim is a direct consequence of one above.

**Claim.** Closed irreducible subsets of  $X^{\text{sob}}$  are exactly  $V_U^c$  for open  $U \subseteq X$  such that (closed) subset  $U^c \subseteq X$  is irreducible.

**Proof of claim.** Let  $V_U^c$  be irreducible and  $U^c = U_1^c \cup U_2^c \subseteq X$ . Then  $V_U = V_{U_1 \cap U_2} = V_{U_1} \cap V_{U_2}$  and thus  $V_U^c = V_{U_1}^c \cup V_{U_2}^c$ . Since  $V_U^c$  is irreducible, we must have  $V_U^c = V_{U_1}^c$  and thus  $U = U_1$ , which proves irreducibility of  $U^c$ .

For the other implication, let  $U^c$  be irreducible and  $V_U^c = V_{U_1}^c \cup V_{U_2}^c$ . Then  $U_1 \cap U_2 = U$ . Since  $U^c$  is irreducible, we must have (WLOG)  $U_1 = U$  and therefore  $V_U = V_{U_1}$ .  $\square$ (of claim)

Let us show  $X^{\text{sob}}$  is sober. Let  $V_U^c$  be closed irreducible. Then by last claim  $U^c$  is closed and irreducible. The set  $U^c$  is the generic point with  $\overline{\{U^c\}} = V_U^c$ . The inclusion  $\overline{\{U^c\}} \subseteq V_U^c$  is obvious, because  $V_U^c$  contains the point  $U^c$  and is a closed set. For the other inclusion take a closed set that  $V_W^c$  that contains  $U^c$ . That means  $U^c \cap W = \emptyset$  and thus  $W \subseteq U$ . Then we have  $V_W \subseteq V_U$  and  $V_U^c \subseteq V_W^c$ . This proves that  $V_U^c$  is the closure of the point  $U^c$ .

2. Define

$$\begin{aligned}
h: X^{\text{sob}} &\rightarrow Z \\
W &\mapsto \text{unique generic point of } \overline{g(W)}.
\end{aligned}$$

Note that: a continuous image of an irreducible set is irreducible and the closure of an irreducible set is irreducible. So  $\overline{g(W)} \subseteq Z$  is a closed irreducible subset and thus has a unique generic point in  $Z$ .

Let's now prove  $g = h \circ f$ . Take  $x \in X$ . We have to prove  $g(x)$  is the unique generic point of  $\overline{g(\{x\})}$ . Clearly  $g(x) \in \overline{g(\{x\})}$ . Take any closed  $W \subseteq Z$  with  $g(x) \in W$ . Then, by definition,  $x \in g^{-1}(W)$ . Because  $g^{-1}(W)$

is closed, also  $\overline{\{x\}} \subseteq g^{-1}(W)$ . So  $g(\overline{\{x\}}) \subseteq W$ . But since  $W$  is closed we have  $\overline{g(\overline{\{x\}})} \subseteq W$ . This proves that  $g(x)$  is indeed a generic point of  $\overline{g(\overline{\{x\}})}$ . So we have  $g = h \circ f$ .

To prove  $h$  is continuous we take an open set  $U \subseteq Z$ . we want to see that  $h^{-1}(U)$  is open. Since  $g^{-1}(U) = f^{-1}(h^{-1}(U))$  is open and  $f^{-1}$  induces a bijection of open sets, the set  $h^{-1}(U)$  is open as well. So  $h$  is continuous.

We should also argue why  $h$  is unique. Take  $h, h': X^{\text{sob}} \rightarrow Z$  both continuous and satisfying  $g = h \circ f = h' \circ f$ . Pick any closed irreducible  $W \subseteq X$ . Suppose  $h(W) \neq h'(W)$ . WLOG there exists open  $U \subseteq Z$  such that  $h(W) \in U$  and  $h'(W) \notin U$  (because requiring *unique* generic point implies  $T_0$  property). Open sets  $h^{-1}(U)$  and  $h'^{-1}(U)$  therefore differ. Using one of the claims above, they are of the form  $V_{U_1} = h^{-1}(U)$  and  $V_{U_2} = h'^{-1}(U)$ . So we have  $W \in V_{U_1}$  and  $W \notin V_{U_2}$ . Then there exists  $w \in W \cap U_1$ , for which  $\overline{\{w\}} \in V_{U_1}$  and  $\overline{\{w\}} \notin V_{U_2}$ . By definition  $\overline{\{w\}} \in h^{-1}(U)$  and  $\overline{\{w\}} \notin h'^{-1}(U)$  which means that  $h(\overline{\{w\}}) \in U$  and  $h'(\overline{\{w\}}) \notin U$ . But that is a contradiction with assumption  $g = h \circ f = h' \circ f$ .

3. We can define  $h: V \rightarrow \text{MaxSpec}(A)$  with  $h(x_1, \dots, x_n) = (X_1 - x_1, \dots, X_n - x_n)$ . Due to Hilberts Nullstellensatz this is a bijection (because  $k$  alg. closed). Take a closed subset  $C \subseteq V$ . Again by Hilberts Nullstellensatz we get a radical ideal  $I$  such that  $V = V(I)$ . So  $h(C) = \{m \in \text{MaxSpec}(A) \mid I \subseteq m\}$ , which is a closed set. And if we take a closed set  $V(I) \subseteq \text{MaxSpec}(A)$  for some  $I \subseteq A$  we have  $h^{-1}(V(I)) = \{x \in V \mid \forall f \in I: f(x) = 0\}$ . So  $h$  is a homeomorphism.

Observe that if we take  $X = \text{MaxSpec}(A)$ , then  $X^{\text{sob}} = \text{Spec}(A)$ . As sets that is clear, because all irreducible closed sets of  $V$  are exactly vanishing sets of prime ideals in  $A$ . And topology on  $V \cong \text{MaxSpec}(A)$  is defined by the basis of closed sets being the vanishing sets of prime ideals, so  $X^{\text{sob}} = \text{Spec}(A)$ .

Then we have a diagram

$$\begin{array}{ccc} V & \xrightarrow{h} & \text{MaxSpec}(A) \\ \downarrow i & & \downarrow j \\ V^{\text{sob}} & \longrightarrow & \text{Spec}(A) \end{array}$$

where  $i: x \mapsto \overline{\{x\}}$  and  $j: m \mapsto \{m\}$  (since maximal ideals are already closed points).

By 2. part we know there exists  $g: V^{\text{sob}} \rightarrow \text{Spec}(A)$  with  $j \circ h = g \circ i$ . And similar there exists  $f: \text{Spec}(A) \rightarrow V^{\text{sob}}$  with  $i \circ h^{-1} = f \circ j$ . Combining these two equations gives us  $j = g \circ f \circ j$  and  $i = f \circ g \circ i$ .

Observe that the map  $g \circ f$  satisfies the “universal condition” from 2. part for the map  $i: V \rightarrow V^{\text{sob}}$ . So it is the unique map  $\pi: V \rightarrow V^{\text{sob}}$  that makes

$$\begin{array}{ccc}
 V & \xrightarrow{i} & V^{\text{sob}} \\
 \downarrow i & \nearrow & \\
 V^{\text{sob}} & & 
 \end{array}$$

And identity would also satisfy that condition, so  $g \circ f$  must be the identity on  $V^{\text{sob}}$ .

We argue exactly the same that  $f \circ g$  is the identity on  $\text{Spec}(A)$ .

This proves that  $V^{\text{sob}} \cong \text{Spec}(A)$ .