GTU, Fall 2020, MATH 101

* Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$ near x = 1.

* Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$ near x = 1.

Note that f(x) is defined for all real numbers x except x = 1.

* Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$ near x = 1.

Note that f(x) is defined for all real numbers x except x = 1.

Even though f(1) is not defined, f(x) approaches arbitrarily close to 2 as x approaches 1.

GTU, Fall 2020, MATH 101

* Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$ near x = 1.

Note that f(x) is defined for all real numbers x except x = 1.

Even though f(1) is not defined, f(x) approaches arbitrarily close to 2 as x approaches 1.We write this as

$$\lim_{x \to 1} f(x) = 2 \text{ or } \lim_{x \to 1} \frac{x^2 - 1}{x - 1} = 2.$$


* Consider the function $f(x) = \frac{x^2 - 1}{x - 1}$ near x = 1.

Note that f(x) is defined for all real numbers x except x = 1.

Even though f(1) is not defined, f(x) approaches arbitrarily close to 2 as x approaches 1.We write this as

$$\lim_{x \to 1} f(x) = 2 \text{ or } \lim_{x \to 1} \frac{x^2 - 1}{x - 1} = 2.$$

We say f(x) approaches the limit 2 as x approaches 1.



An informal definition of limit:

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself).

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself). If we can ensure that f(x) is as close as we want to L by taking x close enough to a, but not equal to a, we say that the limit of f(x) as x approaches a is L.

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself). If we can ensure that f(x) is as close as we want to L by taking x close enough to a, but not equal to a, we say that the limit of f(x) as x approaches a is L and we write

$$\lim_{x\to a} f(x) = L.$$

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself). If we can ensure that f(x) is as close as we want to L by taking x close enough to a, but not equal to a, we say that the limit of f(x) as x approaches a is L, and we write

$$\lim_{x\to a} f(x) = L.$$

The Alternative notation:

$$f(x) \to L$$
 as $x \to a$

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself). If we can ensure that f(x) is as close as we want to L by taking x close enough to a, but not equal to a, we say that the limit of f(x) as x approaches a is L, and we write

$$\lim_{x\to a} f(x) = L.$$

The Alternative notation:

$$f(x) \rightarrow L$$
 as $x \rightarrow a$

* Limits are unique;

An informal definition of limit:

Let f(x) be a function defined near the point a (maybe not a itself). If we can ensure that f(x) is as close as we want to L by taking x close enough to a, but not equal to a, we say that the limit of f(x) as x approaches a is L, and we write

$$\lim_{x\to a}f(x)=L.$$

The Alternative notation:

$$f(x) \rightarrow L$$
 as $x \rightarrow a$

* Limits are unique; if $\lim_{x\to a} f(x) = L$ and $\lim_{x\to a} f(x) = M$, then L = M.

* Example; $\lim_{x\to a} x = a$ and $\lim_{x\to a} c = c$.

* Example; $\lim_{x\to a} x = a$ and $\lim_{x\to a} c = c$.

You can get an answer just by plugging in the limiting value.

* Example; $\lim_{x\to a} x = a$ and $\lim_{x\to a} c = c$.

You can get an answer just by plugging in the limiting value. We will see that if f is an elementary function defined at a then

$$\lim_{x \to a} f(x) = f(a).$$

* A function may be defined on both sides of x = a but still not have a limit at x = a.

* A function may be defined on both sides of x = a but still not have a limit at x = a.

For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.

- * A function may be defined on both sides of x=a but still not have a limit at x=a.
 - For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.
- * The limit of f at x = a may be different from f(a).

- * A function may be defined on both sides of x = a but still not have a limit at x = a.
 - For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.
- * The limit of f at x = a may be different from f(a). For example,

- * A function may be defined on both sides of x = a but still not have a limit at x = a.
 - For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.
- * The limit of f at x = a may be different from f(a). For example,

$$g(x) = \begin{cases} x & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$$

* A function may be defined on both sides of x = a but still not have a limit at x = a.

For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.

* The limit of f at x = a may be different from f(a). For example,

$$g(x) = \begin{cases} x & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$$

then $\lim_{x \to 1} g(x) = 1 \neq g(1)$.

GTU, Fall 2020, MATH 101


* A function may be defined on both sides of x=a but still not have a limit at x=a.

For example, $f(x) = \frac{1}{x}$ has no limit as $x \to 0$.

* The limit of f at x = a may be different from f(a). For example,

$$g(x) = \begin{cases} x & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$$

then $\lim_{x \to 1} g(x) = 1 \neq g(1)$.



GTU, Fall 2020, MATH 101

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

* $x \rightarrow a+$ means x approaches a from the right.

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

* $x \rightarrow a+$ means x approaches a from the right.

Left-limit: $\lim_{x\to a^-} f(x) = L$ is found by looking at values of f(x) when x is less than a (to the left of a on the real line).

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

* $x \rightarrow a+$ means x approaches a from the right.

Left-limit: $\lim_{x\to a^-} f(x) = L$ is found by looking at values of f(x) when x is less than a (to the left of a on the real line).

* $x \rightarrow a-$ means x approaches a from the left.

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

* $x \rightarrow a+$ means x approaches a from the right.

Left-limit: $\lim_{x\to a^-} f(x) = L$ is found by looking at values of f(x) when x is less than a (to the left of a on the real line).

* $x \rightarrow a-$ means x approaches a from the left.

Theorem

Relationship between one-sided and two-sided limits:

One-Sided Limits

Right-limit: $\lim_{x\to a+} f(x) = L$ is known as the right-limit and means that you should use values of x that are greater than a (to the right of a on the real line) to compute the limit.

* $x \rightarrow a+$ means x approaches a from the right.

Left-limit: $\lim_{x\to a^-} f(x) = L$ is found by looking at values of f(x) when x is less than a (to the left of a on the real line).

* $x \rightarrow a-$ means x approaches a from the left.

Theorem

Relationship between one-sided and two-sided limits:

$$\lim_{x \to a} f(x) = L \Longleftrightarrow \lim_{x \to a-} f(x) = \lim_{x \to a+} f(x) = L.$$



GTU, Fall 2020, MATH 101

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

$$\lim_{x\to 0+}\frac{x}{|x|}=\lim_{x\to 0+}\frac{x}{x}=1$$

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

$$\lim_{x\to 0+}\frac{x}{|x|}=\lim_{x\to 0+}\frac{x}{x}=1$$

and

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

$$\lim_{x\to 0+}\frac{x}{|x|}=\lim_{x\to 0+}\frac{x}{x}=1$$

and

$$\lim_{x \to 0-} \frac{x}{|x|} = \lim_{x \to 0+} \frac{x}{-x} = -1.$$

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

$$\lim_{x\to 0+}\frac{x}{|x|}=\lim_{x\to 0+}\frac{x}{x}=1$$

and

$$\lim_{x \to 0-} \frac{x}{|x|} = \lim_{x \to 0+} \frac{x}{-x} = -1.$$

$$\implies \lim_{x \to 0+} \frac{x}{|x|} \neq \lim_{x \to 0-} \frac{x}{|x|}$$

* Let's try to find the $\lim_{x\to 0} \frac{x}{|x|}$.

$$\lim_{x\to 0+}\frac{x}{|x|}=\lim_{x\to 0+}\frac{x}{x}=1$$

and

$$\lim_{x \to 0-} \frac{x}{|x|} = \lim_{x \to 0+} \frac{x}{-x} = -1.$$

$$\implies \lim_{x \to 0+} \frac{x}{|x|} \neq \lim_{x \to 0-} \frac{x}{|x|}$$

 $\implies \lim_{x\to 0} \frac{x^{|x|}}{|x|} does not exist.$

GTU, Fall 2020, MATH 101

Rules for Limits of Functions at a Point

Rules for Limits of Functions at a Point

Rules for Limits of Functions at a Point

If $\lim_{x\to a} f(x) = L$, $\lim_{x\to a} g(x) = M$, and c is a constant, then $*\lim_{x\to a} [f(x) \mp g(x)] = L \mp M$

Rules for Limits of Functions at a Point

- * $\lim_{x\to a}[f(x)\mp g(x)]=L\mp M$
- $* \lim_{x \to a} f(x)g(x) = LM$

Rules for Limits of Functions at a Point

- * $\lim_{x\to a}[f(x)\mp g(x)]=L\mp M$
- * $\lim_{x\to a} f(x)g(x) = LM$
- $* \lim_{x \to a} cf(x) = cL$

Rules for Limits of Functions at a Point

- * $\lim_{x\to a}[f(x)\mp g(x)]=L\mp M$
- * $\lim_{x\to a} f(x)g(x) = LM$
- $* \lim_{x \to a} cf(x) = cL$
- * $\lim_{x\to a} \frac{f(x)}{g(x)} = \frac{L}{M}$, if $M \neq 0$.

Rules for Limits of Functions at a Point

If $\lim_{x\to a} f(x) = L$, $\lim_{x\to a} g(x) = M$, and c is a constant, then

- * $\lim_{x\to a} [f(x) \mp g(x)] = L \mp M$
- $* \lim_{x \to a} f(x)g(x) = LM$
- $* \lim_{x \to a} cf(x) = cL$
- * $\lim_{x\to a} \frac{f(x)}{g(x)} = \frac{L}{M}$, if $M \neq 0$.
- * If m is an integer and n is a positive integer , then

$$\lim_{x\to a} [f(x)]^{m/n} = L^{m/n}$$

provided L > 0 if n is even, and $L \neq 0$ if m < 0.

Rules for Limits of Functions at a Point

If $\lim_{x\to a} f(x) = L$, $\lim_{x\to a} g(x) = M$, and c is a constant, then

- * $\lim_{x\to a}[f(x)\mp g(x)]=L\mp M$
- * $\lim_{x\to a} f(x)g(x) = LM$
- * $\lim_{x\to a} cf(x) = cL$
- * $\lim_{x\to a} \frac{f(x)}{g(x)} = \frac{L}{M}$, if $M \neq 0$.
- * If m is an integer and n is a positive integer, then

$$\lim_{x \to a} [f(x)]^{m/n} = L^{m/n}$$

provided L > 0 if n is even, and $L \neq 0$ if m < 0.

* If $f(x) \leq g(x)$ on an interval containing a in its interior , then

$$L \leq M$$
.



Rules for Limits of Functions at a Point

If $\lim_{x\to a} f(x) = L$, $\lim_{x\to a} g(x) = M$, and c is a constant, then

- * $\lim_{x\to a}[f(x)\mp g(x)]=L\mp M$
- * $\lim_{x\to a} f(x)g(x) = LM$
- * $\lim_{x\to a} cf(x) = cL$
- * $\lim_{x\to a} \frac{f(x)}{g(x)} = \frac{L}{M}$, if $M \neq 0$.
- * If m is an integer and n is a positive integer, then

$$\lim_{x\to a} [f(x)]^{m/n} = L^{m/n}$$

provided L > 0 if n is even, and $L \neq 0$ if m < 0.

* If $f(x) \leq g(x)$ on an interval containing a in its interior , then

$$L \leq M$$
.

Remark: These rules are also valid for right limits and left limits.

We can easily obtain the following important results by using the rules for limits of functions at a point.

We can easily obtain the following important results by using the rules for limits of functions at a point.

Limits of Polynomials and Rational Functions:

We can easily obtain the following important results by using the rules for limits of functions at a point.

Limits of Polynomials and Rational Functions:

1. If P(x) is a polynomial and a is any real number, then

$$\lim_{x \to a} P(x) = P(a)$$

We can easily obtain the following important results by using the rules for limits of functions at a point.

Limits of Polynomials and Rational Functions:

1. If P(x) is a polynomial and a is any real number, then

$$\lim_{x\to a}P(x)=P(a)$$

2. If P(x) and Q(x) are polynomials and $Q(a) \neq 0$, then

$$\lim_{x \to a} \frac{P(x)}{Q(x)} = \frac{P(a)}{Q(a)}$$

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Solution: By using the limit rules

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Solution: By using the limit rules a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$

a)
$$\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$$

Example

Find a)
$$\lim_{x\to 1} \frac{x^2 + x + 4}{x^3 - 2x^2 + 7}$$
 b) $\lim_{x\to 3} \sqrt{2x + 3}$.

Solution: By using the limit rules a)
$$\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7} = \lim_{x\to 1} \frac{1^2+1+4}{1^3-21^2+7} = 1$$

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Solution: By using the limit rules a)
$$\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7} = \lim_{x\to 1} \frac{1^2+1+4}{1^3-21^2+7} = 1$$

b)
$$\lim_{x\to 3} \sqrt{2x+3}$$

Example

Find a) $\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7}$ b) $\lim_{x\to 3} \sqrt{2x+3}$.

Solution: By using the limit rules a)
$$\lim_{x\to 1} \frac{x^2+x+4}{x^3-2x^2+7} = \lim_{x\to 1} \frac{1^2+1+4}{1^3-21^2+7} = 1$$

b)
$$\lim_{x\to 3} \sqrt{2x+3} = \sqrt{2.3+3} = 3$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

a)
$$\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Solution:

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

b)
$$\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Solution:

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

b)
$$\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1} = \lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1} \frac{\sqrt{x}+1}{\sqrt{x}+1}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Solution:

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

b)
$$\lim_{x \to 1} \frac{\sqrt{x} - 1}{x^2 - 1} = \lim_{x \to 1} \frac{\sqrt{x} - 1}{x^2 - 1} \frac{\sqrt{x} + 1}{\sqrt{x} + 1} = \lim_{x \to 1} \frac{x - 1}{(x - 1)(x + 1)(\sqrt{x} + 1)}$$



Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Solution:

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

b)
$$\lim_{x \to 1} \frac{\sqrt{x} - 1}{x^2 - 1} = \lim_{x \to 1} \frac{\sqrt{x} - 1}{x^2 - 1} \frac{\sqrt{x} + 1}{\sqrt{x} + 1} = \lim_{x \to 1} \frac{x - 1}{(x - 1)(x + 1)(\sqrt{x} + 1)} = \lim_{x \to 1} \frac{1}{(x + 1)(\sqrt{x} + 1)}$$

Example

Evaluate a) $\lim_{x\to -3} \frac{x^2+2x-3}{x^2+5x+6}$ b) $\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1}$.

Solution:

a)
$$\lim_{x \to -3} \frac{x^2 + 2x - 3}{x^2 + 5x + 6} = \lim_{x \to -3} \frac{(x+3)(x-1)}{(x+3)(x+2)} = \lim_{x \to -3} \frac{x-1}{x+2} = \frac{-3-1}{-3+2} = 4.$$

b)
$$\lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1} = \lim_{x\to 1} \frac{\sqrt{x}-1}{x^2-1} \frac{\sqrt{x}+1}{\sqrt{x}+1} = \lim_{x\to 1} \frac{x-1}{(x-1)(x+1)(\sqrt{x}+1)} = \lim_{x\to 1} \frac{1}{(x+1)(\sqrt{x}+1)} = \frac{1}{4}.$$



Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Example

If
$$f(x) = \frac{|x-3|}{x^2-x-6}$$
 find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x\to 3+} f(x)$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2-x-6}$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)}$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2-x-6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)}$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{5}.$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2-x-6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{\epsilon}.$$

$$\lim_{x\to 3-} f(x)$$



Example

If
$$f(x) = \frac{|x-3|}{x^2-x-6}$$
 find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3\\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{5}.$$

$$\lim_{x\to 3-} f(x) = \lim_{x\to 3-} \frac{-(x-3)}{x^2-x-6}$$

Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{\overline{\epsilon}}.$$

$$\lim_{x\to 3-} f(x) = \lim_{x\to 3-} \frac{-(x-3)}{x^2-x-6} = \lim_{x\to 3-} \frac{3-x}{(x-3)(x+2)}$$



Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{5}.$$

$$\lim_{x \to 3-} f(x) = \lim_{x \to 3-} \frac{-(x-3)}{x^2 - x - 6} = \lim_{x \to 3-} \frac{3-x}{(x-3)(x+2)} = \lim_{x \to 3-} -\frac{1}{(x+2)}$$



Example

If $f(x) = \frac{|x-3|}{x^2-x-6}$ find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{5}.$$

$$\lim_{x\to 3-} f(x) = \lim_{x\to 3-} \frac{-(x-3)}{x^2 - x - 6} = \lim_{x\to 3-} \frac{3-x}{(x-3)(x+2)} = \lim_{x\to 3-} -\frac{1}{(x+2)} = -\frac{1}{5}.$$



Example

If
$$f(x) = \frac{|x-3|}{x^2-x-6}$$
 find $\lim_{x\to 3+} f(x)$, $\lim_{x\to 3-} f(x)$ and $\lim_{x\to 3} f(x)$.

Solution:

Observe that

$$|x-3| = \begin{cases} x-3 & \text{if } x > 3 \\ -(x-3) & \text{if } x < 3 \end{cases}$$

Therefore

$$\lim_{x \to 3+} f(x) = \lim_{x \to 3+} \frac{x-3}{x^2 - x - 6} = \lim_{x \to 3+} \frac{x-3}{(x-3)(x+2)} = \lim_{x \to 3+} \frac{1}{(x+2)} = \frac{1}{2}$$

$$\lim_{x \to 3^{-}} f(x) = \lim_{x \to 3^{-}} \frac{-(x-3)}{x^{2}-x-6} = \lim_{x \to 3^{-}} \frac{3-x}{(x-3)(x+2)} = \lim_{x \to 3^{-}} -\frac{1}{(x+2)} = -\frac{1}{5}.$$

The limit does not exist since $\lim_{x\to 3+} f(x) \neq \lim_{x\to 3-} f(x)$.



Evaluate $\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$.

Evaluate
$$\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$$
.

Solution:

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}} \cdot \frac{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}$$

Evaluate
$$\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$$
.

Solution:

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}} \cdot \frac{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}$$
$$= \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{x^3 - 5x^2 + 8x - 4} \cdot (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

Evaluate
$$\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$$
.

Solution:

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}} \cdot \frac{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}$$

$$= \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{x^3 - 5x^2 + 8x - 4} \cdot (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

$$= \lim_{x \to 2} \frac{(x - 2)^2(x + 1)}{(x - 2)^2(x - 1)} \cdot \lim_{x \to 2} (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$



Evaluate
$$\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$$
.

Solution:

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}} \cdot \frac{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}$$

$$= \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{x^3 - 5x^2 + 8x - 4} \cdot (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

$$= \lim_{x \to 2} \frac{(x - 2)^2(x + 1)}{(x - 2)^2(x - 1)} \cdot \lim_{x \to 2} (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

$$= \lim_{x \to 2} \frac{x + 1}{x - 1} \cdot 4\sqrt{6}$$



Evaluate
$$\lim_{x\to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}}$$
.

Solution:

$$\lim_{x \to 2} f(x) = \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{\sqrt{x^3 + 8x} - \sqrt{5x^2 + 4}} \cdot \frac{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}{\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4}}$$

$$= \lim_{x \to 2} \frac{x^3 - 3x^2 + 4}{x^3 - 5x^2 + 8x - 4} \cdot (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

$$= \lim_{x \to 2} \frac{(x - 2)^2(x + 1)}{(x - 2)^2(x - 1)} \cdot \lim_{x \to 2} (\sqrt{x^3 + 8x} + \sqrt{5x^2 + 4})$$

$$= \lim_{x \to 2} \frac{x + 1}{x - 1} \cdot 4\sqrt{6} = 12\sqrt{6}$$



The Squeeze Theorem

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that f satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that f satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

Is it possible to determine the value of $\lim_{x\to 2} f(x)$ using only this information?

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that f satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

Is it possible to determine the value of $\lim_{x\to 2} f(x)$ using only this information? Yes

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that *f* satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

Is it possible to determine the value of $\lim_{x\to 2} f(x)$ using only this information? Yes

$$\lim_{x\to 2} (x^3 - 3x - 3) = 2^3 - 3.2 - 3 = -1$$
 and



The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that *f* satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

Is it possible to determine the value of $\lim_{x\to 2} f(x)$ using only this information? Yes

$$\lim_{x\to 2} (x^3 - 3x - 3) = 2^3 - 3.2 - 3 = -1$$
 and $\lim_{x\to 2} (9x^2 - 27x + 17) = 9.2^2 - 27.2 + 17 = -1$

◆ロト ◆個ト ◆園ト ◆園ト ■ めので

The Squeeze Theorem

Suppose that $f(x) \le g(x) \le h(x)$ holds for all x in some open interval containing a, except possibly at x = a itself. If

$$\lim_{x\to a} f(x) = \lim_{x\to a} h(x) = L,$$

the limit of g(x) is also L as $x \to a$. Similar statements hold for left and right limits.

* For example, suppose that *f* satisfied the condition

$$x^3 - 3x - 3 \le f(x) \le 9x^2 - 27x + 17$$
 for $|x - 2| < 1$.

Is it possible to determine the value of $\lim_{x\to 2} f(x)$ using only this information? Yes

$$\lim_{x\to 2} (x^3 - 3x - 3) = 2^3 - 3.2 - 3 = -1$$
 and $\lim_{x\to 2} (9x^2 - 27x + 17) = 9.2^2 - 27.2 + 17 = -1$

 \implies By Squeeze Theorem, $\lim_{x\to 2} f(x) = -1$.

Show that $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

Show that $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

Solution:

$$-1 \le \sin \frac{1}{x} \le 1 \ (\forall x \ne 0) \implies -x \le x \sin \frac{1}{x} \le x \ (\forall x > 0)$$

Show that $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

Solution:

$$-1 \le \sin \frac{1}{x} \le 1 \ (\forall x \ne 0) \implies -x \le x \sin \frac{1}{x} \le x \ (\forall x > 0)$$

 $\lim_{x\to 0+} x = \lim_{x\to 0+} -x = 0$ then $\lim_{x\to 0+} x \sin\frac{1}{x} = 0$ by Squeeze Theorem.

Show that $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

Solution:

$$-1 \le \sin \frac{1}{x} \le 1 \ (\forall x \ne 0) \implies -x \le x \sin \frac{1}{x} \le x \ (\forall x > 0)$$

 $\lim_{x\to 0+} x = \lim_{x\to 0+} -x = 0$ then $\lim_{x\to 0+} x \sin\frac{1}{x} = 0$ by Squeeze Theorem.

Similarly $\lim_{x\to 0-} x \sin\frac{1}{x} = 0$.



Show that $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

Solution:

$$-1 \le \sin \frac{1}{x} \le 1 \ (\forall x \ne 0) \implies -x \le x \sin \frac{1}{x} \le x \ (\forall x > 0)$$

 $\lim_{x\to 0+} x = \lim_{x\to 0+} -x = 0$ then $\lim_{x\to 0+} x \sin\frac{1}{x} = 0$ by Squeeze Theorem.

Similarly $\lim_{x\to 0-} x \sin \frac{1}{x} = 0$.

Thus $\lim_{x\to 0} x \sin \frac{1}{x} = 0$.

