

GTU, Spring 2021, MATH 101

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .
- * Keywords in this definition: 'to each element x in D ' and 'a unique element in E '.

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .
- * Keywords in this definition: 'to each element x in D ' and 'a unique element in E '.
- * We simply write $f : D \rightarrow E$. The set D is called the domain of f .

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .
- * Keywords in this definition: 'to each element x in D ' and 'a unique element in E '.
- * We simply write $f : D \rightarrow E$. The set D is called the domain of f .
- * In calculus we will not see strange functions, all functions will have an explicit formula.

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .
- * Keywords in this definition: 'to each element x in D ' and 'a unique element in E '.
- * We simply write $f : D \rightarrow E$. The set D is called the domain of f .
- * In calculus we will not see strange functions, all functions will have an explicit formula.
- * For example; $f(x) = x^2$ or $y = f(x) = x^2$.

Preliminaries, Functions

- * We are all familiar with the notion of functions. Traditionally let's recall the definition of a function.
- * Let D and E be two sets. A function f on D into E is a rule that assigns a unique element denoted by $f(x)$ in E to each element x in D .
- * Keywords in this definition: 'to each element x in D ' and 'a unique element in E '.
- * We simply write $f : D \rightarrow E$. The set D is called the domain of f .
- * In calculus we will not see strange functions, all functions will have an explicit formula.
- * For example; $f(x) = x^2$ or $y = f(x) = x^2$.
- * x : independent variable, y : dependent variable

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .
- * Sometimes we may specify (or restrict) the domain.

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .
- * Sometimes we may specify (or restrict) the domain.
- * For example, let $f : [0, 1] \rightarrow \mathbb{R}$, $f(x) = \sqrt{x}$. Even the square root function is defined at any positive real number, here we only deal with the interval $[0, 1]$.

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .
- * Sometimes we may specify (or restrict) the domain.
- * For example, let $f : [0, 1] \rightarrow \mathbb{R}$, $f(x) = \sqrt{x}$. Even the square root function is defined at any positive real number, here we only deal with the interval $[0, 1]$.
- * (!) As we will see in this course, many properties of functions depend on the domain as well as the formula of the function.

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .
- * Sometimes we may specify (or restrict) the domain.
- * For example, let $f : [0, 1] \rightarrow \mathbb{R}$, $f(x) = \sqrt{x}$. Even the square root function is defined at any positive real number, here we only deal with the interval $[0, 1]$.
- * (!) As we will see in this course, many properties of functions depend on the domain as well as the formula of the function.
- * So we always need to be careful about the domain of the function (!)

Preliminaries, Functions

- * Domain convention: Whenever a function is given by a formula without any specified domain, we take the largest subset of real numbers on which f makes sense.
- * For example, let $f(x) = \sqrt{x}$, so we take $[0, \infty)$ as the domain of f .
- * Sometimes we may specify (or restrict) the domain.
- * For example, let $f : [0, 1] \rightarrow \mathbb{R}$, $f(x) = \sqrt{x}$. Even the square root function is defined at any positive real number, here we only deal with the interval $[0, 1]$.
- * (!) As we will see in this course, many properties of functions depend on the domain as well as the formula of the function.
- * So we always need to be careful about the domain of the function (!)
- * We may denote the domain of a function by $\mathcal{D}(f)$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.
The only restriction on f is that the denominator can not be zero. So we need to exclude x for which $x^2 = 1$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.
The only restriction on f is that the denominator can not be zero. So we need to exclude x for which $x^2 = 1$.
So $\mathcal{D}(f) = \mathbb{R} - \{1, -1\} = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.
The only restriction on f is that the denominator can not be zero. So we need to exclude x for which $x^2 = 1$.
So $\mathcal{D}(f) = \mathbb{R} - \{1, -1\} = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$.
- * Example: Find the domain of $f(x) = \sqrt{x} + \sqrt{1 - x^2}$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.
The only restriction on f is that the denominator can not be zero. So we need to exclude x for which $x^2 = 1$.
So $\mathcal{D}(f) = \mathbb{R} - \{1, -1\} = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$.
- * Example: Find the domain of $f(x) = \sqrt{x} + \sqrt{1 - x^2}$.
- * The term inside the square root can not be negative, so $x \geq 0$ and $1 - x^2 \geq 0$.

Preliminaries, Functions

- * Example: Find the domain of $f(x) = \frac{x}{x^2 - 1}$.
- * Solution: We need to find the largest subset of real numbers on which f make sense.
The only restriction on f is that the denominator can not be zero. So we need to exclude x for which $x^2 = 1$.
So $\mathcal{D}(f) = \mathbb{R} - \{1, -1\} = (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$.
- * Example: Find the domain of $f(x) = \sqrt{x} + \sqrt{1 - x^2}$.
- * The term inside the square root can not be negative, so $x \geq 0$ and $1 - x^2 \geq 0$.
- * Exercise: Solve these inequalities to find $\mathcal{D}(f)$ (Answer: $\mathcal{D}(f) = [0, 1]$).

Preliminaries, Graphs of Functions

- * By the graph of a function $y = f(x)$ we mean all points of the form $(x, f(x))$ on the Cartesian system.


Preliminaries, Graphs of Functions

- * By the graph of a function $y = f(x)$ we mean all points of the form $(x, f(x))$ on the Cartesian system.
- * (!) Sketching the graphs of functions is difficult in general, and we will see advanced tools to sketch graphs of functions later.


Preliminaries, Graphs of Functions

- * By the graph of a function $y = f(x)$ we mean all points of the form $(x, f(x))$ on the Cartesian system.
- * (!) Sketching the graphs of functions is difficult in general, and we will see advanced tools to sketch graphs of functions later.
- * For now let's see graphs of some basic functions (probably you know these from high school).


Preliminaries, Graphs of Functions




$$y = f(x) = c \quad (\text{a constant function})$$



$$y = f(x) = x \quad (\text{the identity function})$$




$$y = f(x) = x^2$$



$$y = f(x) = \sqrt{x}$$

Preliminaries, Graphs of Functions



Preliminaries, Graphs of Functions


* Let's sketch $y = f(x) = \sqrt{1 - x^2}$.

Preliminaries, Graphs of Functions

- * Let's sketch $y = f(x) = \sqrt{1 - x^2}$.
- * Observe that $\mathcal{D}(f) = [-1, 1]$. Also $y \geq 0$, and that $y^2 = 1 - x^2$, or $x^2 + y^2 = 1$.


Preliminaries, Graphs of Functions

- * Let's sketch $y = f(x) = \sqrt{1 - x^2}$.
- * Observe that $\mathcal{D}(f) = [-1, 1]$. Also $y \geq 0$, and that $y^2 = 1 - x^2$, or $x^2 + y^2 = 1$.
- * So we see that the graph is the part of the unit circle (i.e. $x^2 + y^2 = 1$) for which $y \geq 0$, so that it is the upper semi-circle.



Preliminaries, Graphs of Functions

- * Let's sketch $y = f(x) = \sqrt{1 - x^2}$.
- * Observe that $\mathcal{D}(f) = [-1, 1]$. Also $y \geq 0$, and that $y^2 = 1 - x^2$, or $x^2 + y^2 = 1$.
- * So we see that the graph is the part of the unit circle (i.e. $x^2 + y^2 = 1$) for which $y \geq 0$, so that it is the upper semi-circle.



- * Exercise: How do the graphs of $y = x^4, x^5, x^6, \dots$ look like?

Preliminaries, Graphs of Functions


- * We can also use shifting to sketch the graphs of some functions.

Preliminaries, Graphs of Functions

- * We can also use shifting to sketch the graphs of some functions.
- * Example: Sketch $y = f(x) = 1 + \sqrt{x - 4}$.

Preliminaries, Graphs of Functions

- * We can also use shifting to sketch the graphs of some functions.
- * Example: Sketch $y = f(x) = 1 + \sqrt{x - 4}$.
- * $f(x)$ is similar to \sqrt{x} .



Preliminaries, Graphs of Functions

- * Symmetry is also an important property of functions.

Preliminaries, Graphs of Functions

- * Symmetry is also an important property of functions.
- * A function f is called even if $f(-x) = f(x)$ for any $x, -x \in D(f)$,
e.g; $f(x) = c, x^2, x^4, \dots$

Preliminaries, Graphs of Functions

- * Symmetry is also an important property of functions.
- * A function f is called *even* if $f(-x) = f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = c, x^2, x^4, \dots$
- * A function f is called *odd* if $f(-x) = -f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = x, x^3, x^5, \dots$


Preliminaries, Graphs of Functions

- * Symmetry is also an important property of functions.
- * A function f is called *even* if $f(-x) = f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = c, x^2, x^4, \dots$
- * A function f is called *odd* if $f(-x) = -f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = x, x^3, x^5, \dots$
- * If $f(x)$ is even, and $a \in \mathcal{D}(f)$ then both (a, b) and $(-a, b)$ are on the graph of f .

Preliminaries, Graphs of Functions


- * Symmetry is also an important property of functions.
- * A function f is called *even* if $f(-x) = f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = c, x^2, x^4, \dots$
- * A function f is called *odd* if $f(-x) = -f(x)$ for any $x, -x \in \mathcal{D}(f)$,
e.g; $f(x) = x, x^3, x^5, \dots$
- * If $f(x)$ is even, and $a \in \mathcal{D}(f)$ then both (a, b) and $(-a, b)$ are on the graph of f .
- * If $f(x)$ is odd, and $a \in \mathcal{D}(f)$ then both (a, b) and $(-a, -b)$ are on the graph of f .

Preliminaries, Graphs of Functions



- * So the graph of an even function is symmetric about the y-axis.
- * Similarly the graph of an odd function is symmetric about the origin.

Preliminaries, Graphs of Functions



- * So the graph of an even function is symmetric about the y-axis.
- * Similarly the graph of an odd function is symmetric about the origin.
- * For example compare the graphs of $y = x^2$ and $y = x^3$.

Preliminaries, Graphs of Functions

- * We can reflect the graphs of functions about the x-axis or y-axis.

Preliminaries, Graphs of Functions


- * We can reflect the graphs of functions about the x -axis or y -axis.
- * If we reflect $y = f(x)$ about the x -axis we obtain the graph of $y = -f(x)$.

Preliminaries, Graphs of Functions

- * We can reflect the graphs of functions about the x -axis or y -axis.
- * If we reflect $y = f(x)$ about the x -axis we obtain the graph of $y = -f(x)$.
- * If we reflect $y = f(x)$ about the y -axis we obtain the graph of $y = f(-x)$.


Preliminaries, Graphs of Functions

- * We can reflect the graphs of functions about the x -axis or y -axis.
- * If we reflect $y = f(x)$ about the x -axis we obtain the graph of $y = -f(x)$.
- * If we reflect $y = f(x)$ about the y -axis we obtain the graph of $y = f(-x)$.
- * For example,



Preliminaries, Graphs of Functions

- * We can reflect the graphs of functions about the x -axis or y -axis.
- * If we reflect $y = f(x)$ about the x -axis we obtain the graph of $y = -f(x)$.
- * If we reflect $y = f(x)$ about the y -axis we obtain the graph of $y = f(-x)$.
- * For example,




- * Be careful about the domain (!)

Preliminaries, Graphs of Functions

- * We also have seen the meaning of reflecting about the line $y = x$.


Preliminaries, Graphs of Functions

- * We also have seen the meaning of reflecting about the line $y = x$.
- * This means interchanging x and y . For example,



Preliminaries, Graphs of Functions

- * We also have seen the meaning of reflecting about the line $y = x$.
- * This means interchanging x and y . For example,



- * Exercise: Sketch the graphs of $y = (x - 1)^3 - 5$ and $y = \frac{x - 2}{x + 1}$.