

1. Solutions: 1)a) Since $\sin x$ is a continuous function we have

$$\lim_{n \rightarrow \infty} \sin\left(\frac{\pi n}{2n+1}\right) = \sin\left(\lim_{n \rightarrow \infty} \frac{\pi n}{2n+1}\right) = \sin\left(\frac{\pi}{2}\right) = 1 \neq 0.$$

By using n^{th} term test, the series diverges.

b) Let $a_n = \frac{n}{n^3-4n+1}$ and $b_n = \frac{1}{n^2}$. Then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{n^3}{n^3-4n+1} = 1 \neq 0, \infty$. Since $\sum_{n=2}^{\infty} \frac{1}{n^2}$ convergent p-series $\sum_{n=2}^{\infty} \frac{n}{n^3-4n+1}$ is also convergent by the limit comparison test.

2) Let $u_n = (-1)^n \frac{(x-4)^n}{n^{1/2}}$. By the ratio test

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = |x-4| \lim_{n \rightarrow \infty} \left(\frac{n}{n+1}\right)^{\frac{1}{2}} < 1.$$

So $|x-4| < 1$, i.e., $3 < x < 5$. We have to check end points. For $x = 3$, we have $\sum_{n=1}^{\infty} \frac{1}{n^{\frac{1}{2}}}$ divergent p-series. For $x = 5$, $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^{\frac{1}{2}}}$ does not converge absolutely but $0 < \frac{1}{(n+1)^{1/2}} < \frac{1}{(n)^{1/2}}$ and $\lim_{n \rightarrow \infty} \frac{1}{(n)^{1/2}} = 0$. Therefore, using Alternating series test we get series converge conditionally at 5. Then, the interval of convergence is $3 < x \leq 5$. Also the radius of convergence is $R = 1$.

b) Along $x = 0$ $\lim_{y \rightarrow 1} \frac{0}{(y-1)^2} = 0$ and along $x = y - 1$ $\lim_{x \rightarrow 0} \frac{x^2}{2x^2} = \frac{1}{2}$. As $0 \neq \frac{1}{2}$, limit does not exist.

3)

a) Since $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$ for $|x| < 1$. Using this series we obtain $\frac{1}{1-(-x^2)} = \sum_{n=0}^{\infty} (-1)^n x^{2n}$ for $|x| < 1$.

b) We can integrate above series in this interval and we get

$$\int_0^x \frac{dt}{1+t^2} = \sum_{n=0}^{\infty} (-1)^n \int_0^x t^{2n} dt \implies \arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \text{ for } -1 < x < 1.$$

We have to check the end points: For $x = 1$, we get $\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1}$ alternating series and

$$\sum_{n=0}^{\infty} \left| \frac{(-1)^n}{2n+1} \right| = \sum_{n=0}^{\infty} \frac{1}{2n+1}$$

does not converge absolutely but it converges conditionally from the Alternating series test because $0 < a_{n+1} = \frac{1}{2n+3} < a_n = \frac{1}{2n+1}$ for all n and $\lim_{n \rightarrow \infty} \frac{1}{2n+1} = 0$. For $x = -1$, similarly above case, we get $\sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{2n+1}$ is also convergent conditionally. Therefore, this representation valid for $-1 \leq x \leq 1$.

c) Using b) we obtain

$$\arctan x^2 = \sum_{n=0}^{\infty} (-1)^n \frac{x^{4n+2}}{2n+1} \text{ for } -1 \leq x \leq 1.$$

d) From b) we know that $\arctan x \approx P_3 = x - \frac{x^3}{3}$. Therefore,
 $\arctan \frac{1}{2} = \frac{1}{2} - \frac{1}{3} \left(\frac{1}{2}\right)^3 = \frac{13}{24}$.

4)a)

$$\frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v} = (3v^2 u) f_x(x, y) + u f_y(x, y).$$

Then,

$$\frac{\partial z}{\partial v} \bigg|_{(1,1)} = 3f_x(2,1) + f_y(2,1) = 3 \cdot (1) - 2 = 1.$$

b) Let $F(x, y, z) = x^2 - y^2 - z = 0$. Then the normal of the tangent plane at $(-2, 1, 3)$ is $\nabla F(-2, 1, 3) = (-4, -2, -1)$. So the tangent plane at this point is $-4x - 2y - z = d$ and $d = -4(-2) - 2(1) - 3 = 3$. Then the tangent plane is $-4x - 2y - z = 3$.

5)a) These two planes are perpendicular to each other if their normals are perpendicular to each other. Therefore the dot product of $n_1 = (-1, 3, -5)$ and $n_2 = (2, -1, -1)$ must be zero. If we product these normals, we get $n_1 \cdot n_2 = -2 - 3 + 5 = 0$.

b) The plane which is perpendicular to D_1 and D_2 has the normal n such that its normal is perpendicular to normals of the D_1 and D_2 , that is, $n = n_1 \times n_2 = -8i - 11j - 5k$. As a result of this we have $-8x - 11y - 5z = d$. The point $(0, 1, 1)$ holds the equation of the plane therefore we have $d = -16$ and so the equation of the plane is $8x + 11y + 5z = 16$.

c) $d = \frac{|2(1) - 1 - 1 - 4|}{\sqrt{2^2 + (-1)^2 + (-1)^2}} = \frac{4}{\sqrt{6}}$.