

GTU, Fall 2020, MATH 101

The Derivative

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists.

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$
(If $x \in \mathcal{D}(f) - \mathcal{D}(f')$ and x is not endpoint of $\mathcal{D}(f)$, then x is called singular point of f .)

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$
(If $x \in \mathcal{D}(f) - \mathcal{D}(f')$ and x is not endpoint of $\mathcal{D}(f)$, then x is called singular point of f .)
- * At particular point x_0 :

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$
(If $x \in \mathcal{D}(f) - \mathcal{D}(f')$ and x is not endpoint of $\mathcal{D}(f)$, then x is called singular point of f .)
- * At particular point x_0 :

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$$

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$
(If $x \in \mathcal{D}(f) - \mathcal{D}(f')$ and x is not endpoint of $\mathcal{D}(f)$, then x is called singular point of f .)
- * At particular point x_0 :

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{take } x = x_0 + h)$$

The Derivative

- * The **derivative** of f is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

at all points x for which the limit exists. If f' exist, we say that f is **differentiable** at x .

- * $\mathcal{D}(f') \subseteq \mathcal{D}(f)$
(If $x \in \mathcal{D}(f) - \mathcal{D}(f')$ and x is not endpoint of $\mathcal{D}(f)$, then x is called singular point of f .)
- * At particular point x_0 :

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h} = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{take } x = x_0 + h)$$

- * If $f'(x_0)$ exist then the eq. of the tangent line to $y = f(x)$ at $(x_0, f(x_0))$:

$$y = f(x_0) + f'(x_0)(x - x_0).$$

The Derivative

* If the limit exist

$$f'_+(x_0) = \lim_{x \rightarrow x_0+} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{right derivative at } x = x_0)$$

The Derivative

* If the limit exist

$$f'_+(x_0) = \lim_{x \rightarrow x_0+} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{right derivative at } x = x_0)$$

$$f'_-(x_0) = \lim_{x \rightarrow x_0-} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{left derivative at } x = x_0).$$

The Derivative

- * If the limit exist

$$f'_+(x_0) = \lim_{x \rightarrow x_0+} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{right derivative at } x = x_0)$$

$$f'_-(x_0) = \lim_{x \rightarrow x_0-} \frac{f(x) - f(x_0)}{x - x_0} \quad (\text{left derivative at } x = x_0).$$

- * f is differentiable on $[a, b]$ if $f'(x)$ exists for all x in (a, b) and $f'_+(a)$ and $f'_-(b)$ exist.

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h) - x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h) - x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

* If $f(x) = \sqrt{x}$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h}$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h) - x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

* If $f(x) = \sqrt{x}$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} = \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{x+h} + \sqrt{x})}$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h) - x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

* If $f(x) = \sqrt{x}$, then

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} = \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{(\sqrt{x+h} + \sqrt{x})} \end{aligned}$$

The Derivative

Some Examples:

* If $f(x) = c$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

* If $f(x) = x$, then

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h) - x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1$$

* If $f(x) = \sqrt{x}$, then

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} = \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{(\sqrt{x+h} + \sqrt{x})} = \frac{1}{2\sqrt{x}}. \end{aligned}$$

The Derivative

General Power Rule: If $f(x) = x^r$, then

$$f'(x) = rx^{r-1}$$

for all $r \in \mathbb{R}$ and x for which x^{r-1} makes sense as a real number.

The Derivative

General Power Rule: If $f(x) = x^r$, then

$$f'(x) = rx^{r-1}$$

for all $r \in \mathbb{R}$ and x for which x^{r-1} makes sense as a real number.

For example

* $f(x) = x^{3/2}$

The Derivative

General Power Rule: If $f(x) = x^r$, then

$$f'(x) = rx^{r-1}$$

for all $r \in \mathbb{R}$ and x for which x^{r-1} makes sense as a real number.

For example

$$* f(x) = x^{3/2} \implies f'(x) = \frac{3}{2}x^{(3/2)-1} = \frac{3}{2}\sqrt{x} \text{ for all } x \geq 0.$$

The Derivative

General Power Rule: If $f(x) = x^r$, then

$$f'(x) = rx^{r-1}$$

for all $r \in \mathbb{R}$ and x for which x^{r-1} makes sense as a real number.
For example

$$* f(x) = x^{3/2} \implies f'(x) = \frac{3}{2}x^{(3/2)-1} = \frac{3}{2}\sqrt{x} \text{ for all } x \geq 0.$$

Exercise: $f(x) = x^n$ where $n = 1, 2, 3, \dots$

Show that $f'(x) = nx^{n-1}$.

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

$$f'_-(0) = \lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

$$f'_-(0) = \lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^-} \frac{|x|}{x}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

$$f'_-(0) = \lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x}$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

$$f'_-(0) = \lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x} = -1.$$

The Derivative

Example

Find the derivative function of $f(x) = |x|$.

Solution: By definition of absolute value function

$$f'(x) = 1 \text{ if } x > 0 \text{ and } f'(x) = -1 \text{ if } x < 0.$$

f is not differentiable at $x = 0$ since the left and right derivatives both exist but are not equal:

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = 1$$

$$f'_-(0) = \lim_{x \rightarrow 0^-} \frac{|x| - |0|}{x - 0} = \lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x} = -1.$$

$$\text{Then } f'(x) = \begin{cases} 1 & \text{if } x > 0 \\ -1 & \text{if } x < 0 \end{cases} = \frac{x}{|x|} = \text{sgn}(x).$$

Notations:

- * To show derivative of $y = f(x)$ we can use:

$$\frac{dy}{dx} = \frac{d}{dx}f(x) \text{ (Leibniz' notation)}$$

Notations:

- * To show derivative of $y = f(x)$ we can use:

$$\frac{dy}{dx} = \frac{d}{dx}f(x) \text{ (Leibniz' notation)}$$

$$f'(x) \text{ (Newton's notation)}$$

Notations:

- * To show derivative of $y = f(x)$ we can use:

$$\frac{dy}{dx} = \frac{d}{dx}f(x) \text{ (Leibniz' notation)}$$

$$f'(x) \text{ (Newton's notation)}$$

or

$$D_x y.$$

- * To express the value of the derivative of $f(x)$ at x_0 :

$$\left. \frac{dy}{dx} \right|_{x=x_0} = \left. \frac{d}{dx}f(x) \right|_{x=x_0} = f'(x_0) = D_x f(x_0).$$

Notations:

- * To show derivative of $y = f(x)$ we can use:

$$\frac{dy}{dx} = \frac{d}{dx}f(x) \text{ (Leibniz' notation)}$$

$$f'(x) \text{ (Newton's notation)}$$

or

$$D_x y.$$

- * To express the value of the derivative of $f(x)$ at x_0 :

$$\left. \frac{dy}{dx} \right|_{x=x_0} = \left. \frac{d}{dx}f(x) \right|_{x=x_0} = f'(x_0) = D_x f(x_0).$$

- * the symbol $\frac{d}{dx}$ is a *differential operator* and denoting "the derivative with respect to x of ...".

Notations:

- * To show derivative of $y = f(x)$ we can use:

$$\frac{dy}{dx} = \frac{d}{dx}f(x) \text{ (Leibniz' notation)}$$

$$f'(x) \text{ (Newton's notation)}$$

or

$$D_x y.$$

- * To express the value of the derivative of $f(x)$ at x_0 :

$$\left. \frac{dy}{dx} \right|_{x=x_0} = \left. \frac{d}{dx}f(x) \right|_{x=x_0} = f'(x_0) = D_x f(x_0).$$

- * the symbol $\frac{d}{dx}$ is a *differential operator* and denoting "the derivative with respect to x of ...".

- * $\frac{d}{dx}x^2 = 2x$ (the derivative with respect to x of x^2 is $2x$)

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution:

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

The Derivative

Example

Use the definition of the derivative to calculate $\frac{d}{dx} \left(\frac{x}{x^2+4} \right) \Big|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\frac{d}{dx} \left(\frac{x}{x^2+4} \right) \Big|_{x=0}$$

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0} = \lim_{h \rightarrow 0} \frac{\frac{0+h}{(0+h)^2+4} - \frac{0}{0^2+4}}{h}$$

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\begin{aligned} \left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0} &= \lim_{h \rightarrow 0} \frac{\frac{0+h}{(0+h)^2+4} - \frac{0}{0^2+4}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{h}{h^2+4}}{h} \end{aligned}$$

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\begin{aligned} \left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0} &= \lim_{h \rightarrow 0} \frac{\frac{0+h}{(0+h)^2+4} - \frac{0}{0^2+4}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{h}{h^2+4}}{h} = \lim_{h \rightarrow 0} \frac{h}{h^2+4} \cdot \frac{1}{h} \end{aligned}$$

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\begin{aligned} \left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0} &= \lim_{h \rightarrow 0} \frac{\frac{0+h}{(0+h)^2+4} - \frac{0}{0^2+4}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{h}{h^2+4}}{h} = \lim_{h \rightarrow 0} \frac{h}{h^2+4} \cdot \frac{1}{h} = \lim_{h \rightarrow 0} \frac{1}{h^2+4} \end{aligned}$$

The Derivative

Example

Use the definition of the derivative to calculate $\left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0}$.

Solution: Put $x = 0$ in the expression for the Newton quotient before taking the limit:

$$\begin{aligned} \left. \frac{d}{dx} \left(\frac{x}{x^2+4} \right) \right|_{x=0} &= \lim_{h \rightarrow 0} \frac{\frac{0+h}{(0+h)^2+4} - \frac{0}{0^2+4}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{h}{h^2+4}}{h} = \lim_{h \rightarrow 0} \frac{h}{h^2+4} \cdot \frac{1}{h} = \lim_{h \rightarrow 0} \frac{1}{h^2+4} = \frac{1}{4}. \end{aligned}$$

The Newton quotient

$$\frac{f(x+h) - f(x)}{h}$$

can be written in the form $\Delta y / \Delta x$ where $\Delta y = f(x+h) - f(x)$ is the increment in y , and $\Delta x = (x+h) - x = h$. Using symbols:

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x}.$$

The Derivative

Differentials

The Derivative

Differentials

The differential $y = f(x)$ is defined as:

$$dy = f'(x)dx.$$

The Derivative

Differentials

The differential $y = f(x)$ is defined as:

$$dy = f'(x)dx.$$

The figure below shows the relationship between the increment Δy and the differential dy .

Δy represent the change in height of the curve $y = f(x)$.

dy represent the change in height of the tangent line when x changes by an amount $dx = \Delta x$.

