

GTU, Fall 2020, MATH 101

Continuity

Continuity at a Point

Continuity at a Point

- * Let S be interval or unions of separate intervals. A point $x \in S$ is called an *interior point* of S if there is an open interval (a, b) s.t. $x \in (a, b) \subseteq S$.

Continuity at a Point

- * Let S be interval or unions of separate intervals. A point $x \in S$ is called an *interior point* of S if there is an open interval (a, b) s.t. $x \in (a, b) \subseteq S$. If $x \in S$ is not an interior point, then x is called an *endpoint* of S .

Continuity at a Point

- * Let S be interval or unions of separate intervals. A point $x \in S$ is called an *interior point* of S if there is an open interval (a, b) s.t. $x \in (a, b) \subseteq S$. If $x \in S$ is not an interior point, then x is called an *endpoint* of S .
- * For example consider $S = [-1, 1]$.

Continuity at a Point

- * Let S be interval or unions of separate intervals. A point $x \in S$ is called an *interior point* of S if there is an open interval (a, b) s.t. $x \in (a, b) \subseteq S$. If $x \in S$ is not an interior point, then x is called an *endpoint* of S .
- * For example consider $S = [-1, 1]$.
Interior points : $(-1, 1)$ and end points: $-1, 1$

Continuity at a Point

- * Let S be interval or unions of separate intervals. A point $x \in S$ is called an *interior point* of S if there is an open interval (a, b) s.t. $x \in (a, b) \subseteq S$. If $x \in S$ is not an interior point, then x is called an *endpoint* of S .
- * For example consider $S = [-1, 1]$.
Interior points : $(-1, 1)$ and end points: $-1, 1$
- * For $S = (-\infty, 0) \cup (0, \infty)$, all points in S is an interior point.

Continuity

Definition

f is called is **continuous** at an interior point c of its domain if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

In otherwise f is called **discontinuous** at c .

Continuity


Definition

f is called is **continuous** at an interior point c of its domain if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

In otherwise f is called **discontinuous** at c .

Consider figure below and investigate the continuity of f at c .



Definition

f is called is **right continuous** at c if

$$\lim_{x \rightarrow c+} f(x) = f(c).$$

Continuity

Definition

f is called is **right continuous** at c if

$$\lim_{x \rightarrow c+} f(x) = f(c).$$

f is called is **left continuous** at c if

$$\lim_{x \rightarrow c-} f(x) = f(c).$$

Definition

f is called is **right continuous** at c if

$$\lim_{x \rightarrow c+} f(x) = f(c).$$

f is called is **left continuous** at c if

$$\lim_{x \rightarrow c-} f(x) = f(c).$$

- * f is continuous at c iff it is both right continuous and left continuous at c .

Definition

f is called is **right continuous** at c if

$$\lim_{x \rightarrow c+} f(x) = f(c).$$

f is called is **left continuous** at c if

$$\lim_{x \rightarrow c-} f(x) = f(c).$$

* f is continuous at c iff it is both right continuous and left continuous at c .

* Consider the Heaviside function $H(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$

$H(x)$ is continuous for $x \neq 0$. It is right continuous at 0 since

$$\lim_{x \rightarrow 0+} H(x) = H(0).$$

Continuity

Continuity at an endpoint

Continuity at an endpoint

- * f is called is continuous at a left endpoint c of its domain if it is right continuous at c .

Continuity at an endpoint

- * f is called is continuous at a left endpoint c of its domain if it is right continuous at c .
- * f is called is continuous at a right endpoint c of its domain if it is left continuous at c .


Continuity at an endpoint

- * f is called is continuous at a left endpoint c of its domain if it is right continuous at c .
- * f is called is continuous at a right endpoint c of its domain if it is left continuous at c .
- * For example, the domain of $f(x) = \sqrt{1 - x^2}$ is $[-1, 1]$.

Continuity

Continuity at an endpoint

- * f is called continuous at a left endpoint c of its domain if it is right continuous at c .
- * f is called continuous at a right endpoint c of its domain if it is left continuous at c .
- * For example, the domain of $f(x) = \sqrt{1-x^2}$ is $[-1, 1]$.
 f is continuous at a right endpoint 1 since $\lim_{x \rightarrow 1^-} f(x) = 0 = f(1)$.
 f is continuous at a left endpoint -1 since $\lim_{x \rightarrow -1^+} f(x) = 0 = f(-1)$.



Continuity on an interval:

- * f is continuous on (a, b) if f is continuous at any points $x \in (a, b)$.

Continuity on an interval:

- * f is continuous on (a, b) if f is continuous at any points $x \in (a, b)$.
- * f is continuous on $[a, b]$ if f is continuous on $x \in (a, b)$, and right continuous at a , and left continuous at b .

Continuity on an interval:

- * f is continuous on (a, b) if f is continuous at any points $x \in (a, b)$.
- * f is continuous on $[a, b]$ if f is continuous on $x \in (a, b)$, and right continuous at a , and left continuous at b .
- * f is **continuous function** if f is continuous at every point of its domain.

Continuity on an interval:

- * f is continuous on (a, b) if f is continuous at any points $x \in (a, b)$.
- * f is continuous on $[a, b]$ if f is continuous on $x \in (a, b)$, and right continuous at a , and left continuous at b .
- * f is **continuous function** if f is continuous at every point of its domain.

Exercise: Define continuity on $(a, b]$, $(-\infty, b]$, (a, ∞) .

Continuity on an interval:

- * f is continuous on (a, b) if f is continuous at any points $x \in (a, b)$.
- * f is continuous on $[a, b]$ if f is continuous on $x \in (a, b)$, and right continuous at a , and left continuous at b .
- * f is **continuous function** if f is continuous at every point of its domain.

Exercise: Define continuity on $(a, b]$, $(-\infty, b]$, (a, ∞) .

- * $\lim_{x \rightarrow 0+} \sqrt{x} = 0$ and $\lim_{x \rightarrow a} \sqrt{x} = \sqrt{a} \ (\forall a \in (0, \infty)) \implies \sqrt{x}$ is continuous on $[0, \infty)$.

Continuity

Note that

- * all polynomials;
- * all rational functions;
- * all rational powers $x^{m/n}$;
- * the trigonometric functions;
- * the absolute value functions $|x|$;

are continuous whenever they are defined.

Continuity

- * If f and g are both defined on an interval containing c and both are continuous at c ,

Continuity

- * If f and g are both defined on an interval containing c and both are continuous at c , then

$$\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = f(c) + g(c) = (f + g)(c)$$

Continuity

- * If f and g are both defined on an interval containing c and both are continuous at c , then

$$\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = f(c) + g(c) = (f + g)(c)$$

so $f + g$ is continuous.

Continuity

- * If f and g are both defined on an interval containing c and both are continuous at c , then

$$\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = f(c) + g(c) = (f + g)(c)$$

so $f + g$ is continuous.

Exercise: Show that $f - g$, fg , f/g (provided $g(c) \neq 0$), and $(f(x))^{1/n}$ (provided $f(c) > 0$ if n is even) are continuous.

Continuity

- * If f and g are both defined on an interval containing c and both are continuous at c , then

$$\lim_{x \rightarrow c} (f(x) + g(x)) = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = f(c) + g(c) = (f + g)(c)$$

so $f + g$ is continuous.

Exercise: Show that $f - g$, fg , f/g (provided $g(c) \neq 0$), and $(f(x))^{1/n}$ (provided $f(c) > 0$ if n is even) are continuous.

- * If $f(g(x))$ is defined on an interval containing c , and f is continuous at L and $\lim_{x \rightarrow c} g(x) = L$, then

$$\lim_{x \rightarrow c} f(g(x)) = f(L) = f\left(\lim_{x \rightarrow c} g(x)\right).$$

For example,

$$* 2x^3 + 5$$

$$* \frac{x+3}{x^4-6}$$

$$* \sqrt{x^3 + 2x + 1}$$

$$* \frac{|x|}{|x+1|}$$

are continuous everywhere on their respective domains.

- * *Removable discontinuity*: f has a removable discontinuity at a if f is undefined or discontinuous at a but can be (re)defined at that *single point* so that it becomes continuous there.

- * *Removable discontinuity*: f has a removable discontinuity at a if f is undefined or discontinuous at a but can be (re)defined at that *single point* so that it becomes continuous there.

For example,

- * *Removable discontinuity*: f has a removable discontinuity at a if f is undefined or discontinuous at a but can be (re)defined at that *single point* so that it becomes continuous there.

For example,

- *
$$f(x) = \begin{cases} x & \text{if } x \neq 1 \\ 2 & \text{if } x = 1 \end{cases}$$

has a removable discontinuity at $x = 1$.

Continuity

Continuous Functions on Closed, Finite Intervals

Continuity

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Let $f(x)$ be continuous on $[a, b]$, then there exist numbers p and q in $[a, b]$ such that $\forall x \in [a, b]$,

$$f(p) \leq f(x) \leq f(q).$$

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Let $f(x)$ be continuous on $[a, b]$, then there exist numbers p and q in $[a, b]$ such that $\forall x \in [a, b]$,

$$f(p) \leq f(x) \leq f(q).$$

Thus f has the absolute minimum value $m = f(p)$ and the absolute maximum value $M = f(q)$.

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Let $f(x)$ be continuous on $[a, b]$, then there exist numbers p and q in $[a, b]$ such that $\forall x \in [a, b]$,

$$f(p) \leq f(x) \leq f(q).$$

Thus f has the absolute minimum value $m = f(p)$ and the absolute maximum value $M = f(q)$.

- * The theorem says that minimum and maximum values exist; it does not tell us how to find them.

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Let $f(x)$ be continuous on $[a, b]$, then there exist numbers p and q in $[a, b]$ such that $\forall x \in [a, b]$,

$$f(p) \leq f(x) \leq f(q).$$

Thus f has the absolute minimum value $m = f(p)$ and the absolute maximum value $M = f(q)$.

- * The theorem says that minimum and maximum values exist; it does not tell us how to find them.
- * The theorem implies that a function that is continuous on a closed, finite interval is **bounded**.

Continuous Functions on Closed, Finite Intervals

Theorem

The Max.-Min. Theorem

Let $f(x)$ be continuous on $[a, b]$, then there exist numbers p and q in $[a, b]$ such that $\forall x \in [a, b]$,

$$f(p) \leq f(x) \leq f(q).$$

Thus f has the absolute minimum value $m = f(p)$ and the absolute maximum value $M = f(q)$.

- * The theorem says that minimum and maximum values exist; it does not tell us how to find them.
- * The theorem implies that a function that is continuous on a closed, finite interval is **bounded**. This means there must exist a number K such that

$$|f(x)| \leq K \text{ that is } -K \leq f(x) \leq K.$$

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } (0, 1].$$

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } (0, 1].$$

f is continuous on $(0, 1]$. It is not bounded and it has a minimum value but no maximum value. In fact the minimum value occurs $x = 1$.

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } (0, 1].$$

f is continuous on $(0, 1]$. It is not bounded and it has a minimum value but no maximum value. In fact the minimum value occurs $x = 1$.

* Consider $f(x) = \begin{cases} x & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$

on $[0, 1]$. f is discontinuous at $x = 1$. It has minimum value but no maximum value.

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } (0, 1].$$

f is continuous on $(0, 1]$. It is not bounded and it has a minimum value but no maximum value. In fact the minimum value occurs $x = 1$.

* Consider $f(x) = \begin{cases} x & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$

on $[0, 1]$. f is discontinuous at $x = 1$. It has minimum value but no maximum value.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } [1/2, 1].$$

Continuity

Remark: The conclusion of the theorem may fail if f is not continuous or if the interval is not closed.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } (0, 1].$$

f is continuous on $(0, 1]$. It is not bounded and it has a minimum value but no maximum value. In fact the minimum value occurs $x = 1$.

* Consider $f(x) = \begin{cases} x & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$

on $[0, 1]$. f is discontinuous at $x = 1$. It has minimum value but no maximum value.

* Consider

$$f(x) = \frac{1}{x^2} \text{ on } [1/2, 1].$$

f is continuous on $[1/2, 1]$. It is bounded. It has both a maximum and a minimum value.

Continuity

Continuous Functions on Closed, Finite Intervals

Continuity

Continuous Functions on Closed, Finite Intervals

Theorem

The Intermediate-Value Theorem

Continuous Functions on Closed, Finite Intervals

Theorem

The Intermediate-Value Theorem

Let $f(x)$ be continuous on $[a, b]$ and d be a value between $f(a)$ and $f(b)$. Then there exists $c \in (a, b)$ such that $f(c) = d$.


Continuity

Continuous Functions on Closed, Finite Intervals

Theorem

The Intermediate-Value Theorem

Let $f(x)$ be continuous on $[a, b]$ and d be a value between $f(a)$ and $f(b)$. Then there exists $c \in (a, b)$ such that $f(c) = d$.



* Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

Continuity

* Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\}$$

* Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

Continuity

- * Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

- * Show that the equation $\sin x = x - 1$ has a solution.

Continuity

- * Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

- * Show that the equation $\sin x = x - 1$ has a solution.

Define $f(x) = \sin x - x + 1 = \sin x - (x - 1)$

Continuity

- * Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

- * Show that the equation $\sin x = x - 1$ has a solution.

Define $f(x) = \sin x - x + 1 = \sin x - (x - 1)$

$$\left. \begin{array}{l} f(x) \text{ is continuous on } [0, \pi] \\ f(0) = 1 > 0, f(\pi) = -\pi + 1 < 0 \end{array} \right\}$$

Continuity

- * Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

- * Show that the equation $\sin x = x - 1$ has a solution.

Define $f(x) = \sin x - x + 1 = \sin x - (x - 1)$

$$\left. \begin{array}{l} f(x) \text{ is continuous on } [0, \pi] \\ f(0) = 1 > 0, f(\pi) = -\pi + 1 < 0 \end{array} \right\} \implies \exists c \in [0, \pi] \text{ s.t. } f(c) = 0$$

Continuity

- * Show that $P(x) = x^5 - 4x + 1$ has a root in $[0, 1]$.

$$\left. \begin{array}{l} P(x) \text{ is continuous on } [0, 1] \\ P(0) = 1 > 0 \\ P(1) = -2 < 0 \end{array} \right\} \implies \exists c \in [0, 1] \text{ s.t. } P(c) = 0$$

by IVT.

- * Show that the equation $\sin x = x - 1$ has a solution.

Define $f(x) = \sin x - x + 1 = \sin x - (x - 1)$

$$\left. \begin{array}{l} f(x) \text{ is continuous on } [0, \pi] \\ f(0) = 1 > 0, f(\pi) = -\pi + 1 < 0 \end{array} \right\} \implies \exists c \in [0, \pi] \text{ s.t. } f(c) = 0$$

by IVT.