

GTU, Fall 2021, MATH 101

Preliminaries, Real Numbers

- * Calculus I is all about the functions defined on real numbers or smaller subsets of real numbers.

Preliminaries, Real Numbers


- * Calculus I is all about the functions defined on real numbers or smaller subsets of real numbers.
- * Two basic notions: *Real Numbers* and *Functions*. Let's start by recalling real numbers.

Preliminaries, Real Numbers

- * Calculus I is all about the functions defined on real numbers or smaller subsets of real numbers.
- * Two basic notions: *Real Numbers* and *Functions*. Let's start by recalling real numbers.
- * Real numbers have decimal expansion; e.g. $-3 = -3.000000\dots$, $1/2 = 0.5$, $\pi = 3.141\dots$ (Decimal expansion is not a good way to represent real numbers).


Preliminaries, Real Numbers

- * Calculus I is all about the functions defined on real numbers or smaller subsets of real numbers.
- * Two basic notions: *Real Numbers* and *Functions*. Let's start by recalling real numbers.
- * Real numbers have decimal expansion; e.g. $-3 = -3.000000\dots$, $1/2 = 0.5$, $\pi = 3.141\dots$ (Decimal expansion is not a good way to represent real numbers).
- * A better way is to use the *real line*



Preliminaries, Real Numbers

- * Calculus I is all about the functions defined on real numbers or smaller subsets of real numbers.
- * Two basic notions: *Real Numbers* and *Functions*. Let's start by recalling real numbers.
- * Real numbers have decimal expansion; e.g. $-3 = -3.000000\dots$, $1/2 = 0.5$, $\pi = 3.141\dots$ (Decimal expansion is not a good way to represent real numbers).
- * A better way is to use the *real line*



- * There is no hole on the real line, so there is no *gap* between real numbers.

Preliminaries, Real Numbers

- * Recall some important subsets of real numbers.
 - i) Natural numbers, \mathbb{N} : The numbers $0, 1, 2, 3, \dots$
 - ii) Integers, \mathbb{Z} : $\dots, -2, -1, 0, 1, 2, \dots$ (or $0, \pm 1, \pm 2, \dots$)
 - iii) Rational Numbers, \mathbb{Q} : all numbers of the form a/b where a and b are integers and $b \neq 0$. For example, $1/2, -3/7, \dots$ are rational.

Preliminaries, Real Numbers

- * Recall some important subsets of real numbers.
 - i) Natural numbers, \mathbb{N} : The numbers $0, 1, 2, 3, \dots$
 - ii) Integers, \mathbb{Z} : $\dots, -2, -1, 0, 1, 2, \dots$ (or $0, \pm 1, \pm 2, \dots$)
 - iii) Rational Numbers, \mathbb{Q} : all numbers of the form a/b where a and b are integers and $b \neq 0$. For example, $1/2, -3/7, \dots$ are rational.
- * But the set of real numbers, \mathbb{R} is much more larger (!). If a real number is not rational then it is called irrational. For example, $\sqrt{2} = 1.41\dots$ and $\pi = 3.14\dots$ are irrational.

Preliminaries, Real Numbers

- * Recall some important subsets of real numbers.
 - i) Natural numbers, \mathbb{N} : The numbers $0, 1, 2, 3, \dots$
 - ii) Integers, \mathbb{Z} : $\dots, -2, -1, 0, 1, 2, \dots$ (or $0, \pm 1, \pm 2, \dots$)
 - iii) Rational Numbers, \mathbb{Q} : all numbers of the form a/b where a and b are integers and $b \neq 0$. For example, $1/2, -3/7, \dots$ are rational.
- * But the set of real numbers, \mathbb{R} is much more larger (!). If a real number is not rational then it is called irrational. For example, $\sqrt{2} = 1.41\dots$ and $\pi = 3.14\dots$ are irrational.
- * **Exercise:** What is the difference between the decimal expansion of a rational number and of an irrational number?

* $\sqrt{2}$ is not a rational number. Let's see why it is irrational.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $gcd(m, n) = 1$ (greatest common divisor).

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $\gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

Since m is even, we can write $m = 2k$, where $k \in \mathbb{Z}$.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

Since m is even, we can write $m = 2k$, where $k \in \mathbb{Z}$.

$$4k^2 = 2n^2 \implies n^2 = 2k^2$$

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

Since m is even, we can write $m = 2k$, where $k \in \mathbb{Z}$.

$$4k^2 = 2n^2 \implies n^2 = 2k^2$$

Thus n is also even.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $\gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

Since m is even, we can write $m = 2k$, where $k \in \mathbb{Z}$.

$$4k^2 = 2n^2 \implies n^2 = 2k^2$$

Thus n is also even. But m and n cannot both be even.

- * $\sqrt{2}$ is not a rational number. Let's see why it is irrational.
- * Suppose that it is rational. Then $\sqrt{2} = m/n$, where $m, n \in \mathbb{Z}$ and $n \neq 0$. Assume that $\gcd(m, n) = 1$ (greatest common divisor). Now

$$m^2/n^2 = 2 \implies m^2 = 2n^2 \implies m \text{ must be even}$$

Since m is even, we can write $m = 2k$, where $k \in \mathbb{Z}$.

$$4k^2 = 2n^2 \implies n^2 = 2k^2$$

Thus n is also even. But m and n cannot both be even. Accordingly, there can be no rational number whose square is 2.

Preliminaries, Real Numbers

- * For Calculus we need different special subsets of real numbers called *intervals*

Preliminaries, Real Numbers

* For Calculus we need different special subsets of real numbers called *intervals*


$$* (a, b) = \{x \in \mathbb{R} \mid a < x < b\}$$

$$* [a, b] = \{x \in \mathbb{R} \mid a \leq x \leq b\}$$

$$* (a, b] = \{x \in \mathbb{R} \mid a < x \leq b\}$$

$$* [a, b) = \{x \in \mathbb{R} \mid a \leq x < b\}$$

* These are all *finite* intervals.



Preliminaries, Real Numbers

* We may also talk about *infinite* intervals.


* $(a, +\infty) = \{x \in \mathbb{R} \mid a < x\}$

* $(-\infty, b] = \{x \in \mathbb{R} \mid x \leq b\}$

* $(-\infty, +\infty) = \mathbb{R}$

* $+\infty$: plus infinity,

* $-\infty$: minus infinity.



Preliminaries, Real Numbers

- * An important notion on real numbers is *the absolute value*.

Preliminaries, Real Numbers

- * An important notion on real numbers is *the absolute value*.
- * The absolute value of a real number x , denoted by $|x|$ is the 'distance' of x to 0 on the real line.

Preliminaries, Real Numbers

- * An important notion on real numbers is *the absolute value*.
- * The absolute value of a real number x , denoted by $|x|$ is the 'distance' of x to 0 on the real line.
- * For example; $|-3| = 3$, $|5.1| = 5.1$, $|0| = 0$
- * Absolute value can not be negative.

Preliminaries, Real Numbers


- * An important notion on real numbers is *the absolute value*.
- * The absolute value of a real number x , denoted by $|x|$ is the 'distance' of x to 0 on the real line.
- * For example; $|-3| = 3$, $|5.1| = 5.1$, $|0| = 0$
- * Absolute value can not be negative.
- * **Exercise:** Solve the following equalities/inequalities (Find all values of x which satisfy the given equality/inequality).
 - i) $x|x - 2| \leq 0$ (Answer: $x \leq 0$ or $x = 2$).
 - ii) $x^2 - 3x + 2 > 0$ (Answer: $x > 2$ or $x < 1$).
 - iii) $1 - 2x = |x|$ (Answer: $x = 1/3$ or $x = 1$).

Preliminaries, Real Numbers

- * An important notion on real numbers is *the absolute value*.
- * The absolute value of a real number x , denoted by $|x|$ is the 'distance' of x to 0 on the real line.
- * For example; $|-3| = 3$, $|5.1| = 5.1$, $|0| = 0$
- * Absolute value can not be negative.
- * **Exercise:** Solve the following equalities/inequalities (Find all values of x which satisfy the given equality/inequality).
 - i) $x|x - 2| \leq 0$ (Answer: $x \leq 0$ or $x = 2$).
 - ii) $x^2 - 3x + 2 > 0$ (Answer: $x > 2$ or $x < 1$).
 - iii) $1 - 2x = |x|$ (Answer: $x = 1/3$ or $x = 1$).

(Note: These are of high school level, you can use sign table to solve such inequalities. Similar problems will be solved in detail in problem sessions).


Preliminaries, Cartesian Coordinate System




* xy -plane is divided into four regions called *quadrants* numbered by I, II, III and IV.

- * O : The origin
- * P : a point on the xy -plane.
- * a : the x -coordinate of P
- * b : the y -coordinate of P

Preliminaries, Cartesian Coordinate System - Distance Formula



Preliminaries, Cartesian Coordinate System - Distance Formula



The distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by

$$D = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$


(This follows by Pythagorean theorem).

Preliminaries, Cartesian Coordinate System - Line Equations

- * The basic rule: There is one and only one line passing through given two distinct points.

Preliminaries, Cartesian Coordinate System - Line Equations

- * The basic rule: There is one and only one line passing through given two distinct points.



- * The slope is $m = \frac{y_2 - y_1}{x_2 - x_1}$.
- * The line is given by the equation

$$L : \frac{y - y_1}{x - x_1} = m$$

(Why?)

Preliminaries, Cartesian Coordinate System - Line Equations

- * Example; the slope of the line L passing through $P(1, 2)$ and $Q(2, 5)$ is

$$m = \frac{5 - 2}{2 - 1} = 3$$

Preliminaries, Cartesian Coordinate System - Line Equations

- * Example; the slope of the line L passing through $P(1, 2)$ and $Q(2, 5)$ is

$$m = \frac{5 - 2}{2 - 1} = 3$$

- * So $L : \frac{y - 2}{x - 1} = 3$, simplify to obtain $L : y = 3x - 1$.

Preliminaries, Cartesian Coordinate System - Line Equations

- * Example; the slope of the line L passing through $P(1, 2)$ and $Q(2, 5)$ is

$$m = \frac{5 - 2}{2 - 1} = 3$$

- * So $L : \frac{y - 2}{x - 1} = 3$, simplify to obtain $L : y = 3x - 1$.
- * In general a line on the xy -plane is represented by an equation of the form $Ax + By = C$ where A, B and C are constant and x, y represent the coordinates of the points on the line.

Preliminaries, Cartesian Coordinate System - Line Equations

- * Example; the slope of the line L passing through $P(1, 2)$ and $Q(2, 5)$ is

$$m = \frac{5 - 2}{2 - 1} = 3$$

- * So $L : \frac{y - 2}{x - 1} = 3$, simplify to obtain $L : y = 3x - 1$.
- * In general a line on the xy -plane is represented by an equation of the form $Ax + By = C$ where A, B and C are constant and x, y represent the coordinates of the points on the line.
- * A more useful form for line equations is $y = mx + n$. Here m is the slope and n is the y -intercept (the point at which the line cuts the y -axis).

Preliminaries, Cartesian Coordinate System - Line Equations

- * Example; the slope of the line L passing through $P(1, 2)$ and $Q(2, 5)$ is

$$m = \frac{5 - 2}{2 - 1} = 3$$



- * So $L : \frac{y - 2}{x - 1} = 3$, simplify to obtain $L : y = 3x - 1$.
- * In general a line on the xy -plane is represented by an equation of the form $Ax + By = C$ where A, B and C are constant and x, y represent the coordinates of the points on the line.
- * A more useful form for line equations is $y = mx + n$. Here m is the slope and n is the y -intercept (the point at which the line cuts the y -axis).
- * The x -intercept (the point at which the line cuts the x -axis) can be found by setting $y = 0$, so it is $x = 1/3$.

Preliminaries, Cartesian Coordinate System - Line Equations

- * Let's consider two specific type of lines.


Preliminaries, Cartesian Coordinate System - Line Equations

- * Let's consider two specific type of lines.
- * If $m = 0$, then the line equation is $y = n$, so that it is a *horizontal* line.



Preliminaries, Cartesian Coordinate System - Line Equations

- * Let's consider two specific type of lines.
- * If $m = 0$, then the line equation is $y = n$, so that it is a *horizontal* line.



- * We may also have *vertical* lines with equation $x = c$. Vertical lines do not have a slope (!) (or do they have? what would you choose for the 'slope' of a vertical line?).