

GTU, Fall 2020, MATH 101

Differentiation Rules

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0))$$

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0)$$

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0) = f'(x_0) \cdot 0 = 0$$

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0) = f'(x_0) \cdot 0 = 0$$

$$\implies \lim_{x \rightarrow x_0} f(x) = f(x_0).$$



Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0) = f'(x_0) \cdot 0 = 0$$

$$\implies \lim_{x \rightarrow x_0} f(x) = f(x_0).$$



Remarks:

* the converse is false.

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0) = f'(x_0) \cdot 0 = 0$$

$$\implies \lim_{x \rightarrow x_0} f(x) = f(x_0).$$



Remarks:

- * the converse is false.

Counter-example: $|x|$ is continuous at $x = 0$ but not differentiable at $x = 0$.

Differentiation Rules

Theorem

If $f(x)$ is differentiable at x_0 , then $f(x)$ is continuous at x_0 .

Proof.

We know $\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$ exist since $f(x)$ is differentiable at x_0 .

$$\lim_{x \rightarrow x_0} (f(x) - f(x_0)) = \lim_{x \rightarrow x_0} \left(\frac{f(x) - f(x_0)}{x - x_0} \right) (x - x_0) = f'(x_0) \cdot 0 = 0$$

$$\implies \lim_{x \rightarrow x_0} f(x) = f(x_0).$$



Remarks:

- * the converse is false.

Counter-example: $|x|$ is continuous at $x = 0$ but not differentiable at $x = 0$.

- * if f is not continuous then f is not differentiable

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

$$* (f + g)'(x) = f'(x) + g'(x),$$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

$$* (f + g)'(x) = f'(x) + g'(x),$$

$$* (f - g)'(x) = f'(x) - g'(x),$$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

$$* (f + g)'(x) = f'(x) + g'(x),$$

$$* (f - g)'(x) = f'(x) - g'(x),$$

$$* (Cf)'(x) = Cf'(x),$$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

- * $(f + g)'(x) = f'(x) + g'(x),$
- * $(f - g)'(x) = f'(x) - g'(x),$
- * $(Cf)'(x) = Cf'(x),$
- * $(fg)'(x) = f'(x)g(x) + f(x)g'(x),$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

- * $(f + g)'(x) = f'(x) + g'(x),$
- * $(f - g)'(x) = f'(x) - g'(x),$
- * $(Cf)'(x) = Cf'(x),$
- * $(fg)'(x) = f'(x)g(x) + f(x)g'(x),$
- * $\left(\frac{1}{f}\right)'(x) = \frac{-f'(x)}{(f(x))^2}, \quad (f(x) \neq 0)$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

$$* (f + g)'(x) = f'(x) + g'(x),$$

$$* (f - g)'(x) = f'(x) - g'(x),$$

$$* (Cf)'(x) = Cf'(x),$$

$$* (fg)'(x) = f'(x)g(x) + f(x)g'(x),$$

$$* \left(\frac{1}{f}\right)'(x) = \frac{-f'(x)}{(f(x))^2}, \quad (f(x) \neq 0)$$

$$* \left(\frac{f}{g}\right)'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{(g(x))^2} \quad (g(x) \neq 0).$$

Rules for differentiation:

Let f and g be differentiable at x , and C be a constant. Then

$$* (f + g)'(x) = f'(x) + g'(x),$$

$$* (f - g)'(x) = f'(x) - g'(x),$$

$$* (Cf)'(x) = Cf'(x),$$

$$* (fg)'(x) = f'(x)g(x) + f(x)g'(x),$$

$$* \left(\frac{1}{f}\right)'(x) = \frac{-f'(x)}{(f(x))^2}, \quad (f(x) \neq 0)$$

$$* \left(\frac{f}{g}\right)'(x) = \frac{g(x)f'(x) - f(x)g'(x)}{(g(x))^2} \quad (g(x) \neq 0).$$

Differentiation Rules

proof of the product rule:

Differentiation Rules

proof of the product rule:

$$(fg)'(x) = \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h}$$

Differentiation Rules

proof of the product rule:

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left(f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x) \right)\end{aligned}$$

Differentiation Rules

proof of the product rule:

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left(f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x) \right) \\&= \lim_{h \rightarrow 0} g(x+h) \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} f(x) \frac{g(x+h) - g(x)}{h}\end{aligned}$$

Differentiation Rules

proof of the product rule:

$$\begin{aligned}(fg)'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left(f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x) \right) \\&= \lim_{h \rightarrow 0} g(x+h) \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} f(x) \frac{g(x+h) - g(x)}{h} \\&= f'(x)g(x) + f(x)g'(x)\end{aligned}$$

since g is continuous.

Some examples:

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

Some examples:

* Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\frac{dy}{dx} = \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4)$$

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4) \\ &= \frac{3}{2x^{1/2}} + \frac{1}{x^{2/3}} - \frac{3}{x^4}.\end{aligned}$$

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4) \\ &= \frac{3}{2x^{1/2}} + \frac{1}{x^{2/3}} - \frac{3}{x^4}.\end{aligned}$$

- * Find y' for $y = \frac{x+3}{x^2+3}$.

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4) \\ &= \frac{3}{2x^{1/2}} + \frac{1}{x^{2/3}} - \frac{3}{x^4}.\end{aligned}$$

- * Find y' for $y = \frac{x+3}{x^2+3}$.

$$y' = \left(\frac{x+3}{x^2+3}\right)' = \frac{(x+3)'(x^2+3) - (x+3)(x^2+3)'}{(x^2+3)^2}$$

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4) \\ &= \frac{3}{2x^{1/2}} + \frac{1}{x^{2/3}} - \frac{3}{x^4}.\end{aligned}$$

- * Find y' for $y = \frac{x+3}{x^2+3}$.

$$\begin{aligned}y' &= \left(\frac{x+3}{x^2+3}\right)' = \frac{(x+3)'(x^2+3) - (x+3)(x^2+3)'}{(x^2+3)^2} \\ &= \frac{(x^2+3) - 2x(x+3)}{(x^2+3)^2}\end{aligned}$$

Some examples:

- * Find f' for $f(x) = x^{-n}$ where $n \in \mathbb{N}$.

$$\frac{d}{dx}(x^{-n}) = \frac{d}{dx}\left(\frac{1}{x^n}\right) = -\frac{nx^{n-1}}{(x^n)^2} = -nx^{-n-1}.$$

- * Find $\frac{dy}{dx}$ for $y = 3\sqrt{x} - \frac{2}{\sqrt{x}} + \frac{1}{x^3} + 4$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(3x^{1/2}) - \frac{d}{dx}(2x^{-1/2}) + \frac{d}{dx}(x^{-3}) + \frac{d}{dx}(4) \\ &= \frac{3}{2x^{1/2}} + \frac{1}{x^{2/3}} - \frac{3}{x^4}.\end{aligned}$$

- * Find y' for $y = \frac{x+3}{x^2+3}$.

$$\begin{aligned}y' &= \left(\frac{x+3}{x^2+3}\right)' = \frac{(x+3)'(x^2+3) - (x+3)(x^2+3)'}{(x^2+3)^2} \\ &= \frac{(x^2+3) - 2x(x+3)}{(x^2+3)^2} = \frac{3-6x-x^2}{(x^2+3)^2}.\end{aligned}$$

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Solution: We need to identify the points where $f'(a) = -3$.

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Solution: We need to identify the points where $f'(a) = -3$. So let's find those:

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Solution: We need to identify the points where $f'(a) = -3$. So let's find those:

$$y = f(x) = x + \frac{1}{x} \implies f'(x) = 1 - \frac{1}{x^2}$$

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Solution: We need to identify the points where $f'(a) = -3$. So let's find those:

$$y = f(x) = x + \frac{1}{x} \implies f'(x) = 1 - \frac{1}{x^2}$$

We want the particular points where $f'(a) = -3$.

Differentiation Rules

Example

Find the equations of any tangent line(s) to the curve $y = x + \frac{1}{x}$ at points where the tangent is parallel to the line $y = -3x$.

Solution: We need to identify the points where $f'(a) = -3$. So let's find those:

$$y = f(x) = x + \frac{1}{x} \implies f'(x) = 1 - \frac{1}{x^2}$$

We want the particular points where $f'(a) = -3$.

$$f'(a) = 1 - \frac{1}{a^2} = -3 \implies -\frac{1}{a^2} = -4 \implies a = \pm \frac{1}{2}$$

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

and so the point $(-1/2, -5/2)$ is on the line and its equation is:

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

and so the point $(-1/2, -5/2)$ is on the line and its equation is:

$$y - (-5/2) = -3(x - (-1/2)) \text{ or } y = -3x - 4$$

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

and so the point $(-1/2, -5/2)$ is on the line and its equation is:

$$y - (-5/2) = -3(x - (-1/2)) \text{ or } y = -3x - 4$$

And for $x_2 = 1/2$ we have $y_2 = 5/2$

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

and so the point $(-1/2, -5/2)$ is on the line and its equation is:

$$y - (-5/2) = -3(x - (-1/2)) \text{ or } y = -3x - 4$$

And for $x_2 = 1/2$ we have $y_2 = 5/2$ and

Differentiation Rules

We know from above the x -value of one point on each line, so we just need to find the corresponding y -value of those two points:

Recall the original function

$$f(x) = x + \frac{1}{x}$$

Hence

$$x_1 = -\frac{1}{2} \implies y_1 = f\left(-\frac{1}{2}\right) = -\frac{1}{2} + (-2) = -\frac{5}{2}$$

and so the point $(-1/2, -5/2)$ is on the line and its equation is:

$$y - (-5/2) = -3(x - (-1/2)) \text{ or } y = -3x - 4$$

And for $x_2 = 1/2$ we have $y_2 = 5/2$ and

$$y - (5/2) = -3(x - 1/2) \text{ or } y = -3x + 4.$$