

Consider the minimum spanning tree T of G under the edge weights $\{a_e\}$, and suppose T were not a minimum-altitude connected subgraph. Then there would be some pair of nodes u and v , and two u - v paths $P \neq P^*$ (represented as sets of edges), so that P is the u - v path in T but P^* has smaller height. In other words, there is an edge $e' = (u', v')$ on P that has the maximum altitude over all edges in $P \cup P^*$. Now, if we consider the edges in $(P \cup P^*) - \{e'\}$, they contain a (possibly self-intersecting) u' - v' path; we can construct such a path by walking along P from u' to u , then along P^* from u to v , and then along P from v to v' . Thus $(P \cup P^*) - \{e'\}$ contains a simple path Q . But then $Q \cup \{e'\}$ is a cycle on which e' is the heaviest edge, contradicting the Cycle Property. Thus, T must be a minimum-altitude connected subgraph.

Now consider a connected subgraph $H = (V, E')$ that does not contain all the edges of T ; let $e = (u, v)$ be an edge of T that is not part of E' . Deleting e from T partitions T into two connected components; and these two components represent a partition of V into sets A and B . The edge e is the minimum-altitude edge with one end in A and the other in B . Since any path in H from u to v must cross at some point from A to B , and it cannot use e , it must have height greater than a_e . It follows that H cannot be a minimum-altitude connected subgraph.