



Department of Computer Science  
UNIVERSITY OF COLORADO **BOULDER**



# Hypothesis Testing II: Two Sample $t$ Tests

Introduction to Data Science Algorithms

Jordan Boyd-Graber and Michael Paul

OCTOBER 6, 2016

## Comparing Two Samples

---

- Thus far, we've tested whether data is consistent with mean  $\mu$
- What if we want to test whether two samples are from the same distribution

## Comparing Two Samples

---

- Thus far, we've tested whether data is consistent with mean  $\mu$
- What if we want to test whether two samples are from the same distribution
- Two-Sample  $t$ -test

## Two-Sample (unpooled)

---

- Two samples  $X_1 = \{x_{1,1}, x_{1,2} \dots x_{1,N_1}\}$  and  $X_2 = \{x_{2,1}, x_{2,2} \dots x_{2,N_2}\}$
- Doesn't assume that variance is the same for both samples (unpooled)
- Compute mean and sample variance for sample 1 ( $\bar{x}_1, s_1^2$ ) and sample 2 ( $\bar{x}_2, s_2^2$ )

## Test Statistic

---

- T-statistic

$$T = \frac{(\bar{x}_1 - \bar{x}_2)}{\sqrt{\frac{s_1^2}{N_1} + \frac{s_2^2}{N_2}}} \quad (1)$$

- Plug into t-distribution with

$$\nu = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{N_2}\right)^2}{\frac{\left(\frac{s_1^2}{N_1}\right)^2}{N_1-1} + \frac{\left(\frac{s_2^2}{N_2}\right)^2}{N_2-1}} \quad (2)$$

- Intuition: Difference between  $\bar{x}_1$  and  $\bar{x}_2$  has variance that's an interpolation between the two samples

## Test Statistic

---

- T-statistic

$$T = \frac{(\bar{x}_1 - \bar{x}_2)}{\sqrt{\frac{s_1^2}{N_1} + \frac{s_2^2}{N_2}}} \quad (1)$$

- Plug into t-distribution with

$$\nu = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{N_2}\right)^2}{\frac{\left(\frac{s_1^2}{N_1}\right)^2}{N_1-1} + \frac{\left(\frac{s_2^2}{N_2}\right)^2}{N_2-1}} \quad (2)$$

- Intuition: Difference between  $\bar{x}_1$  and  $\bar{x}_2$  has variance that's an interpolation between the two samples
- Two-tailed vs. one-tailed distinction still applies