



# Support Vector Machines


Jordan Boyd-Graber University of Colorado Boulder LECTURE 9

Slides adapted from Tom Mitchell, Eric Xing, and Lauren Hannah


.

### **Administrative Questions**

### Find the maximum margin hyperplane




### Find the maximum margin hyperplane




Which are the support vectors?

Working geometrically:




## Working geometrically:

• The maximum margin weight vector will be parallel to the shortest line connecting points of the two classes, that is, the line between (1,1) and (2,3), giving a weight vector of (1,2).



## Working geometrically:


- The maximum margin weight vector will be parallel to the shortest line connecting points of the two classes, that is, the line between (1,1) and (2,3), giving a weight vector of (1,2).
- The optimal decision surface is orthogonal to that line and intersects it at the halfway point. Therefore, it passes through (1.5, 2).




### Working geometrically:

- The maximum margin weight vector will be parallel to the shortest line connecting points of the two classes, that is, the line between (1,1) and (2,3), giving a weight vector of (1,2).
- The optimal decision surface is orthogonal to that line and intersects it at the halfway point. Therefore, it passes through (1.5, 2).
- The SVM decision boundary is:

$$0 = \frac{1}{2}x + y - \frac{11}{4} \Leftrightarrow 0 = \frac{2}{5}x + \frac{4}{5}y - \frac{11}{5}$$






$$w_1x_1+w_2x_2+b$$

Jordan Boyd-Graber Boulder Support Vector Machines

6 of 6




 $.4x_1 + .8x_2 - 2.2$ 

Jordan Boyd-Graber |

Boulder

Support Vector Machines

#### **Cannonical Form**



 $.4x_1 + .8x_2 - 2.2$ 

- $.4 \cdot 1 + .8 \cdot 1 2.2 = -1$
- $.4 \cdot \frac{3}{2} + .8 \cdot 2 = 0$
- $.4 \cdot 2 + .8 \cdot 3 2.2 = +1$