



Department of Computer Science

UNIVERSITY OF COLORADO **BOULDER**




## Slack SVMs

Jordan Boyd-Graber  
University of Colorado Boulder

LECTURE 9


## Can SVMs Work Here?

---



## Can SVMs Work Here?


---



$$y_i(w \cdot x_i + b) \geq 1 \quad (1)$$

## Trick: Allow for a few bad apples

---



## New objective function

---

$$\min_{w,b,\xi} \frac{1}{2} \|w\|^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to  $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

## New objective function

---

$$\min_{w,b,\xi} \frac{1}{2} ||w||^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to  $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin

## New objective function

---

$$\min_{w,b,\xi} \frac{1}{2} ||w||^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to  $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin
- How wrong a point is (slack variables)

## New objective function

---

$$\min_{w,b,\xi} \frac{1}{2} ||w||^2 + \textcolor{red}{C} \sum_{i=1} \xi_i^p \quad (2)$$

subject to  $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin
- How wrong a point is (slack variables)
- **Tradeoff between margin and slack variables**



## New objective function

---

$$\min_{w,b,\xi} \frac{1}{2} ||w||^2 + C \sum_{i=1} \xi_i^p \quad (2)$$

subject to  $y_i(w \cdot x_i + b) \geq 1 - \xi_i \wedge \xi_i \geq 0, i \in [1, m]$

- Standard margin
- How wrong a point is (slack variables)
- Tradeoff between margin and slack variables
- How bad wrongness scales

## Aside: Loss Functions


---

- Losses measure how bad a mistake is
- Important for slack as well

## Aside: Loss Functions

---


- Losses measure how bad a mistake is
- Important for slack as well



## Aside: Loss Functions

---


- Losses measure how bad a mistake is
- Important for slack as well



## Aside: Loss Functions

---


- Losses measure how bad a mistake is
- Important for slack as well



## Aside: Loss Functions

---

- Losses measure how bad a mistake is
- Important for slack as well



We'll focus on linear hinge loss

## Optimizing Constrained Functions

---

### Theorem: Lagrange Multiplier Method

Given functions  $f(x_1, \dots, x_n)$  and  $g(x_1, \dots, x_n)$ , the critical points of  $f$  restricted to the set  $g = 0$  are solutions to equations:

$$\begin{aligned}\frac{\partial f}{\partial x_i}(x_1, \dots, x_n) &= \lambda \frac{\partial g}{\partial x_i}(x_1, \dots, x_n) \quad \forall i \\ g(x_1, \dots, x_n) &= 0\end{aligned}$$

This is  $n + 1$  equations in the  $n + 1$  variables  $x_1, \dots, x_n, \lambda$ .

## Lagrange Example

---

Maximize  $f(x, y) = \sqrt{xy}$  subject to the constraint  $20x + 10y = 200$ .

- Compute derivatives



## Lagrange Example

---

Maximize  $f(x, y) = \sqrt{xy}$  subject to the constraint  $20x + 10y = 200$ .

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \sqrt{\frac{x}{y}} \quad \frac{\partial g}{\partial y} = 10$$

## Lagrange Example

---

Maximize  $f(x, y) = \sqrt{xy}$  subject to the constraint  $20x + 10y = 200$ .

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2} \sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2} \sqrt{\frac{x}{y}} \quad \frac{\partial g}{\partial y} = 10$$

- Create new systems of equations

## Lagrange Example

---

Maximize  $f(x, y) = \sqrt{xy}$  subject to the constraint  $20x + 10y = 200$ .

- Compute derivatives

$$\frac{\partial f}{\partial x} = \frac{1}{2}\sqrt{\frac{y}{x}} \quad \frac{\partial g}{\partial x} = 20$$

$$\frac{\partial f}{\partial y} = \frac{1}{2}\sqrt{\frac{x}{y}} \quad \frac{\partial g}{\partial y} = 10$$

- Create new systems of equations

$$\frac{1}{2}\sqrt{\frac{y}{x}} = 20\lambda$$

$$\frac{1}{2}\sqrt{\frac{x}{y}} = 10\lambda$$

$$20x + 10y = 200$$

## Lagrange Example

---

- Dividing the first equation by the second gives us

$$\frac{y}{x} = 2 \tag{3}$$

- which means  $y = 2x$ , plugging this into the constraint equation gives:

$$20x + 20(2x) = 200$$

$$x = 5 \Rightarrow y = 10$$

## New Lagrangian

---

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\vec{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i (w \cdot x_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

## New Lagrangian

---

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\vec{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i(\vec{w} \cdot \vec{x}_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients  $(\nabla_{\vec{w}} \mathcal{L}, \nabla_b \mathcal{L}, \nabla_{\xi_i})$  and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7) \quad \vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i \quad (8) \quad \alpha_i + \beta_i = C \quad (9)$$

## New Lagrangian

---

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\vec{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i (w \cdot x_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients ( $\nabla_{\vec{w}} \mathcal{L}, \nabla_b \mathcal{L}, \nabla_{\xi_i} \mathcal{L}$ ) and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7) \quad \vec{w} = \sum_{i=1}^m \alpha_i y_i x_i \quad (8) \quad \alpha_i + \beta_i = C \quad (9)$$

## New Lagrangian

---

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\vec{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i (w \cdot x_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients  $(\nabla_{\vec{w}} \mathcal{L}, \nabla_b \mathcal{L}, \nabla_{\xi_i} \mathcal{L})$  and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7) \quad \vec{w} = \sum_{i=1}^m \alpha_i y_i x_i \quad (8) \quad \alpha_i + \beta_i = C \quad (9)$$



## New Lagrangian

---

$$\mathcal{L}(\vec{w}, b, \vec{\xi}, \vec{\alpha}, \vec{\beta}) = \frac{1}{2} \|\vec{w}\|^2 + C \sum_{i=1}^m \xi_i \quad (4)$$

$$- \sum_{i=1}^m \alpha_i [y_i (w \cdot x_i + b) - 1 + \xi_i] \quad (5)$$

$$- \sum_{i=1}^m \beta_i \xi_i \quad (6)$$

Taking the gradients  $(\nabla_w \mathcal{L}, \nabla_b \mathcal{L}, \nabla_{\xi_i})$  and solving for zero gives us

$$\sum_{i=1}^m \alpha_i y_i = 0 \quad (7) \quad \vec{w} = \sum_{i=1}^m \alpha_i y_i x_i \quad (8) \quad \alpha_i + \beta_i = C \quad (9)$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i x_i$$

$$\alpha_i + \beta_i = C$$

## Simplifying dual objective

---

$$\begin{aligned}\sum_{i=1}^m \alpha_i y_i &= 0 & \vec{w} &= \sum_{i=1}^m \alpha_i y_i \mathbf{x}_i & \alpha_i + \beta_i &= C \\ \mathcal{L} &= \frac{1}{2} \|\vec{w}_i\|^2 - \sum_i^m \alpha_i y_i \vec{w} \cdot \vec{x}_i - \sum_i^m \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i & (10)\end{aligned}$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i \quad (10)$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b - \sum_{i=1}^m \beta_i \xi_i \quad (10)$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$

## Simplifying dual objective

---

$$\sum_{i=1}^m \alpha_i y_i = 0$$

$$\vec{w} = \sum_{i=1}^m \alpha_i y_i \vec{x}_i$$

$$\alpha_i + \beta_i = C$$

$$\mathcal{L} = \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) - \sum_i^m \alpha_i y_i b + \sum_i^m \alpha_i \quad (10)$$



## Simplifying dual objective

---

$$\begin{aligned}\sum_{i=1}^m \alpha_i y_i &= 0 & \vec{w} &= \sum_{i=1}^m \alpha_i y_i \vec{x}_i & \alpha_i + \beta_i &= C \\ \mathcal{L} &= \frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i \vec{x}_i \right\|^2 - \sum_i \sum_j \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) + \sum_i \alpha_i & (10)\end{aligned}$$

First two terms are the same!

## Simplifying dual objective

---

$$\begin{aligned} \sum_{i=1}^m \alpha_i y_i &= 0 & \vec{w} &= \sum_{i=1}^m \alpha_i y_i \vec{x}_i & \alpha_i + \beta_i &= C \\ \mathcal{L} &= -\frac{1}{2} \sum_i^m \sum_j^m \alpha_i \alpha_j y_i y_j (\vec{x}_j \cdot \vec{x}_i) + \sum_i^m \alpha_i \end{aligned} \quad (10)$$

Just like separable case, except that we add the constraint that  $\alpha_i \leq C$ !

## Wrapup

---

- Adding slack variables don't break the SVM problem
- Very popular algorithm
  - SVMLight (many options)
  - Libsvm / Liblinear (very fast)
  - Weka (friendly)
  - pyml (Python focused, from Colorado)
- Next time: simple algorithm for finding SVMs