

# An Overview of Graceful Labelings

Calder Evans, Breelyn Cocke

Northern Arizona University  
Department of Mathematics and Statistics

**FAMUS**

6 December 2024



---

Department of Mathematics and Statistics

In this presentation we will:

1. Define graceful labelings.

In this presentation we will:

1. Define graceful labelings.
2. Present known results on graceful labelings.

In this presentation we will:

1. Define graceful labelings.
2. Present known results on graceful labelings.
3. Look over past research from an NAU student.

In this presentation we will:

1. Define graceful labelings.
2. Present known results on graceful labelings.
3. Look over past research from an NAU student.
4. Introduce the Ringel-Kotzig Conjecture and state recent progress.

We define the following:

We define the following:

## Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.

We define the following:

## Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.

## Definition

A **graceful labeling** is a labeling of a graph with  $n$  edges in which:

We define the following:

## Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.

## Definition

A **graceful labeling** is a labeling of a graph with  $n$  edges in which:

- the vertices are labeled with a subset of distinct integers from the set  $\{0, 1, 2, \dots, n\}$ ,

We define the following:

## Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.

## Definition

A **graceful labeling** is a labeling of a graph with  $n$  edges in which:

- the vertices are labeled with a subset of distinct integers from the set  $\{0, 1, 2, \dots, n\}$ ,
- the edges are labeled with a subset of distinct integers from the set  $\{1, 2, \dots, n\}$ ,

We define the following:

### Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.

### Definition

A **graceful labeling** is a labeling of a graph with  $n$  edges in which:

- the vertices are labeled with a subset of distinct integers from the set  $\{0, 1, 2, \dots, n\}$ ,
- the edges are labeled with a subset of distinct integers from the set  $\{1, 2, \dots, n\}$ ,
- each edge label is the absolute difference of its vertex endpoint labels,

We define the following:

## Definition

A **graph**,  $G$ , is a collection of vertices,  $V(G)$ , and a collection of edges,  $E(G)$ , that connect a possibly empty subset of vertices.


## Definition

A **graceful labeling** is a labeling of a graph with  $n$  edges in which:


- the vertices are labeled with a subset of distinct integers from the set  $\{0, 1, 2, \dots, n\}$ ,
- the edges are labeled with a subset of distinct integers from the set  $\{1, 2, \dots, n\}$ ,
- each edge label is the absolute difference of its vertex endpoint labels,
- vertex and edge labelings are unique.

Vertex Labels:  $\{0, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$   
Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, \emptyset\}$

Edge Labels:  $\{1, 2, 3, 4, 5, \emptyset\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$   
Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$




Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$   
Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$



Vertex Labels:  $\{\emptyset, 1, 2, 3, 4, 5, 6\}$

Edge Labels:  $\{1, 2, 3, 4, 5, 6\}$



## Definition

A **path** graph,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.

## Definition

A **path graph**,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

## Definition

A **path graph**,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

## Definition

A **path graph**,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .
  - Edge labels need to be from the set  $\{1, 2, 3, 4, 5\}$ .

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .
  - Edge labels need to be from the set  $\{1, 2, 3, 4, 5\}$ .
- The vertices are distinctly labeled  $\{0, 1, 2, 3, 4, 5\}$ .

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .
  - Edge labels need to be from the set  $\{1, 2, 3, 4, 5\}$ .
- The vertices are distinctly labeled  $\{0, 1, 2, 3, 4, 5\}$ .
- The edges are distinctly labeled  $\{1, 2, 3, 4, 5\}$

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .
  - Edge labels need to be from the set  $\{1, 2, 3, 4, 5\}$ .
- The vertices are distinctly labeled  $\{0, 1, 2, 3, 4, 5\}$ .
- The edges are distinctly labeled  $\{1, 2, 3, 4, 5\}$
- Each edge label is the absolute difference of its vertex endpoint labels.

## Definition

A *path graph*,  $P_n$ , is a graph with  $n$  vertices, of which two have degree 1 and all others have degree 2.




Figure: A graceful labeling of the path  $P_6$ .

Is  $P_6$  labeled gracefully?

We can see that:

- $P_6$  has five edges, so:
  - Vertex labels need to be from the set  $\{0, 1, 2, 3, 4, 5\}$ .
  - Edge labels need to be from the set  $\{1, 2, 3, 4, 5\}$ .
- The vertices are distinctly labeled  $\{0, 1, 2, 3, 4, 5\}$ .
- The edges are distinctly labeled  $\{1, 2, 3, 4, 5\}$ .
- Each edge label is the absolute difference of its vertex endpoint labels.

Thus,  $P_6$  has a graceful labeling.

## Definition

A *cycle graph*,  $C_n$ , is a graph with  $n$  vertices that has a path that starts and ends on the same vertex, in which every vertex has degree two.

## Definition

A *cycle graph*,  $C_n$ , is a graph with  $n$  vertices that has a path that starts and ends on the same vertex, in which every vertex has degree two.




Figure: A graceful labeling of the cycle  $C_8$ .

## "The Father of Graceful-ness"

Alexander Rosa received his Ph.D. in mathematics from the Slovak Academy of Science in 1966, and published his paper "On certain valuations of the vertices of a graph" in 1967.



Alexander Rosa received his Ph.D. in mathematics from the Slovak Academy of Science in 1966, and published his paper "On certain valuations of the vertices of a graph" in 1967.



Rosa was the first to study graceful labelings; he originally referred to them as  $\beta$ -valuations.

Rosa proved seven theorems in his original 1967 paper on graceful labelings, some of which included:

Rosa proved seven theorems in his original 1967 paper on graceful labelings, some of which included:

### Theorem

*All path graphs have a graceful labeling.*

Rosa proved seven theorems in his original 1967 paper on graceful labelings, some of which included:

## Theorem

*All path graphs have a graceful labeling.*

## Theorem

*A cycle with  $n$  vertices is graceful if and only if  $n \equiv 0$  or  $3 \pmod{4}$ .*

The labeling functions for graceful cycles are:

If  $n \equiv 0 \pmod{4}$ :

$$f(v_i) = \begin{cases} \frac{i-1}{2}, & \text{if } i \text{ is odd,} \\ n+1 - \frac{i}{2}, & \text{if } i \text{ is even and } i \leq \frac{n}{2}, \\ n - \frac{i}{2}, & \text{if } i \text{ is even and } i > \frac{n}{2}. \end{cases}$$


If  $n \equiv 3 \pmod{4}$ :

$$f(v_i) = \begin{cases} n+1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{n-1}{2}, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{n-1}{2}. \end{cases}$$


Here, the vertices of  $C_n$  are denoted  $v_1, v_2, \dots, v_n$ , where each  $\{v_i, v_{i+1}\}$  is an edge for all  $1 \leq i \leq n-1$ , as is  $\{v_n, v_1\}$ .

**It is important to note that a given labeling function is not necessarily unique.**

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 1 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 1 : 1 \text{ is odd and } 1 \leq 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 1 : 1$  is odd and  $1 \leq 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 1 : 1 \text{ is odd and } 1 \leq 3 \Rightarrow f(v_1) = \frac{1-1}{2}$$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 1 : 1 \text{ is odd and } 1 \leq 3 \Rightarrow f(v_1) = \frac{1-1}{2} = 0$$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 2 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 2 : 2$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 2 : 2$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 2 : 2 \text{ is even} \Rightarrow f(v_2) = 7 + 1 - \frac{2}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 2 : 2 \text{ is even} \Rightarrow f(v_2) = 7 + 1 - \frac{2}{2} = 7$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 3 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 3 : 3$  is odd and  $3 \leq 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 3 : 3$  is odd and  $3 \leq 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 3 : 3 \text{ is odd and } 3 \leq 3 \Rightarrow f(v_3) = \frac{3-1}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 3 : 3 \text{ is odd and } 3 \leq 3 \Rightarrow f(v_3) = \frac{3-1}{2} = 1$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 4 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 4 : 4$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 4 : 4$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 4 : 4 \text{ is even} \Rightarrow f(v_4) = 7 + 1 - \frac{4}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 4 : 4 \text{ is even} \Rightarrow f(v_4) = 7 + 1 - \frac{4}{2} = 6$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 5 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 5 : 5$  is odd and  $5 > 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 5 : 5$  is odd and  $5 > 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 5 : 5 \text{ is odd and } 5 > 3 \Rightarrow f(v_5) = \frac{5+1}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 5 : 5 \text{ is odd and } 5 > 3 \Rightarrow f(v_5) = \frac{5+1}{2} = 3$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 6 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 6 : 6$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 6 : 6$  is even

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 6 : 6 \text{ is even} \Rightarrow f(v_6) = 7 + 1 - \frac{6}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 6 : 6 \text{ is even} \Rightarrow f(v_6) = 7 + 1 - \frac{6}{2} = 5$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 7 :$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 7 : 7 \text{ is odd and } 7 > 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$i = 7 : 7$  is odd and  $7 > 3$

$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$i = 7 : 7 \text{ is odd and } 7 > 3 \Rightarrow f(v_7) = \frac{7+1}{2}$$


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$i = 7 : 7 \text{ is odd and } 7 > 3 \Rightarrow f(v_7) = \frac{7+1}{2} = 4$$

## Labeling Function Example


$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$




$$7 \equiv 3 \pmod{4} : f(v_i) = \begin{cases} 7 + 1 - \frac{i}{2}, & \text{if } i \text{ is even,} \\ \frac{i-1}{2}, & \text{if } i \text{ is odd and } i \leq \frac{7-1}{2} = 3, \\ \frac{i+1}{2}, & \text{if } i \text{ is odd and } i > \frac{7-1}{2} = 3. \end{cases}$$



## Shark $n$ -Cycles with $n \equiv 3 \pmod{4}$ are Graceful

# Shark $n$ -Cycles with $n \equiv 3 \pmod{4}$ are Graceful






# Graceful Work from an NAU student

## Definition

A *pendant graph* is an  $n$ -cycle graph with one pendant vertex attached to each cycle vertex.

Dr. Alessandra Graf researched graceful labelings of pendant graphs for several semesters at NAU starting in 2013. She was able to prove that all pendant graphs that have labels 0 and  $2n$ , where  $n \equiv 4$  or  $8 \pmod{8}$  and  $n \equiv 0 \pmod{16}$  can be gracefully labeled, while also finding the labeling functions for them.



## Miscellaneous Graph Families With Graceful Labelings

Since Rosa first introduced the idea of graceful labeling, many mathematicians (including Dr. Alessandra Graf) have proved that certain graphs can always be gracefully labeled.

## Miscellaneous Graph Families With Graceful Labelings

Since Rosa first introduced the idea of graceful labeling, many mathematicians (including Dr. Alessandra Graf) have proved that certain graphs can always be gracefully labeled.

Let's look at some of them...

# Miscellaneous Graphs With Graceful Labelings

## Theorem

All **wheel**, web, helm, gear, grid, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various mathematicians]




Figure: The wheel graph  $W_7$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, *web*, helm, gear, grid, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The web graph  $W_6$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, **helm**, gear, grid, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The helm graph  $H_6$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, helm, gear, grid, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The gear graph  $G_6$ .

## Theorem

All wheel, web, helm, gear, **grid**, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The grid graph  $P_5 \times P_4$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, helm, gear, grid, star, complete, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The star graph  $S_7$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, helm, gear, grid, star, **complete**, banana tree, and caterpillar tree graphs are graceful. [Various]




Figure: The complete graph  $K_6$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, helm, gear, grid, star, complete, *banana tree*, and caterpillar tree graphs are graceful. [Various]




Figure: The banana tree graph  $B_{4,5}$ .

# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All wheel, web, helm, gear, grid, star, complete, banana tree, and **caterpillar tree** graphs are graceful. [Various]





Figure: A caterpillar tree graph.


# Miscellaneous Graph Families With Graceful Labelings

## Theorem

All *n*-dimension hypercubes are graceful. [Kotzig, 1981]



3-dimensional hypercube



4-dimensional hypercube

The conjecture that all trees are graceful was proposed by Gerhard Ringel and Anton Kotzig, often identified as the [Ringel-Kotzig Conjecture](#).

The conjecture that all trees are graceful was proposed by Gerhard Ringel and Anton Kotzig, often identified as the **Ringel-Kotzig Conjecture**.

## Definition

A **tree** is an undirected acyclic graph in which any two vertices are connected by exactly one path.

The conjecture that all trees are graceful was proposed by Gerhard Ringel and Anton Kotzig, often identified as the **Ringel-Kotzig Conjecture**.

## Definition

A **tree** is an undirected acyclic graph in which any two vertices are connected by exactly one path.

- This is a major open conjecture in graph theory.

The conjecture that all trees are graceful was proposed by Gerhard Ringel and Anton Kotzig, often identified as the **Ringel-Kotzig Conjecture**.

## Definition

A **tree** is an undirected acyclic graph in which any two vertices are connected by exactly one path.

- This is a major open conjecture in graph theory.
- Kotzig commented that trying to prove this conjecture was a “disease” .

The conjecture that all trees are graceful was proposed by Gerhard Ringel and Anton Kotzig, often identified as the **Ringel-Kotzig Conjecture**.

## Definition

A **tree** is an undirected acyclic graph in which any two vertices are connected by exactly one path.

- This is a major open conjecture in graph theory.
- Kotzig commented that trying to prove this conjecture was a “disease” .
- It has several related (yet weaker) conjectures, many of which have proven partial results.

# The Ringel-Kotzig Conjecture – Recent Progress

Mathematicians have proven all trees to be graceful up to  $n$  vertices for the following values of  $n$ :

# The Ringel-Kotzig Conjecture – Recent Progress

Mathematicians have proven all trees to be graceful up to  $n$  vertices for the following values of  $n$ :

- All trees with up to 27 vertices are graceful. [Mckay et al., 1998]

Mathematicians have proven all trees to be graceful up to  $n$  vertices for the following values of  $n$ :

- All trees with up to 27 vertices are graceful. [Mckay et al., 1998]
- All trees with up to 29 vertices are graceful. [Horton, 2003]

# The Ringel-Kotzig Conjecture – Recent Progress

Mathematicians have proven all trees to be graceful up to  $n$  vertices for the following values of  $n$ :

- All trees with up to **27** vertices are graceful. [Mckay et al., 1998]
- All trees with up to **29** vertices are graceful. [Horton, 2003]
- All trees with up to **35** vertices are graceful. [Fang et al., 2010]

Mathematicians have proven all trees to be graceful up to  $n$  vertices for the following values of  $n$ :

- All trees with up to 27 vertices are graceful. [Mckay et al., 1998]
- All trees with up to 29 vertices are graceful. [Horton, 2003]
- All trees with up to 35 vertices are graceful. [Fang et al., 2010]

Questions?