

4-Output Games and 3-Colorings

Calder Evans and Taylor Smith

Northern Arizona University
Department of Mathematics and Statistics

Undergraduate Symposium
25 April 2025



Department of Mathematics and Statistics

In this presentation we will:

1. Share our research goals

In this presentation we will:

1. Share our research goals
2. Overview Non-local games

In this presentation we will:

1. Share our research goals
2. Overview Non-local games
3. Walkthrough the Mermin-Peres Magic Square Game

In this presentation we will:

1. Share our research goals
2. Overview Non-local games
3. Walkthrough the Mermin-Peres Magic Square Game
4. Explain quantum coloring and encoding of game rules

In this presentation we will:

1. Share our research goals
2. Overview Non-local games
3. Walkthrough the Mermin-Peres Magic Square Game
4. Explain quantum coloring and encoding of game rules
5. Show how our graph is Quantum 3-Colorable and not Classically 3-Colorable

In this presentation we will:

1. Share our research goals
2. Overview Non-local games
3. Walkthrough the Mermin-Peres Magic Square Game
4. Explain quantum coloring and encoding of game rules
5. Show how our graph is Quantum 3-Colorable and not Classically 3-Colorable
6. Share progress with whittling down our graph

Research Goal: Find the smallest graph that can be quantum 3-colored while being not classically 3-colorable. To do this we:

Research Goal: Find the smallest graph that can be quantum 3-colored while being not classically 3-colorable. To do this we:

- built a graph based on the game algebra of the Mermin-Peres Magic Square Game

Research Goal: Find the smallest graph that can be quantum 3-colored while being not classically 3-colorable. To do this we:

- built a graph based on the game algebra of the Mermin-Peres Magic Square Game
 - We chose the Mermin-Peres game because it is the smallest synchronous game that has a winning quantum strategy but no classical strategy

Research Goal: Find the smallest graph that can be quantum 3-colored while being not classically 3-colorable. To do this we:

- built a graph based on the game algebra of the Mermin-Peres Magic Square Game
 - We chose the Mermin-Peres game because it is the smallest synchronous game that has a winning quantum strategy but no classical strategy
- encoded the graph into SageMath and ran a multitude of constructions to identify which vertices could be removed without changing the colorabilities of the graph

Games

Games

- A game \mathcal{G} is played by two players, Alice and Bob.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$
 - If $\lambda(a, b, x, y) = 0$, then Alice and Bob lose.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$
 - If $\lambda(a, b, x, y) = 0$, then Alice and Bob lose.
 - If $\lambda(a, b, x, y) = 1$, then Alice and Bob win.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$
 - If $\lambda(a, b, x, y) = 0$, then Alice and Bob lose.
 - If $\lambda(a, b, x, y) = 1$, then Alice and Bob win.
- We use the following to define \mathcal{G} 's parameters: $\mathcal{G} = (\mathcal{O}, \mathcal{I}, \lambda)$.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$
 - If $\lambda(a, b, x, y) = 0$, then Alice and Bob lose.
 - If $\lambda(a, b, x, y) = 1$, then Alice and Bob win.
- We use the following to define \mathcal{G} 's parameters: $\mathcal{G} = (\mathcal{O}, \mathcal{I}, \lambda)$.

Non-Local Games

- We say a game $\mathcal{G} = (\mathcal{O}, \mathcal{I}, \lambda)$ is non-local if Alice and Bob can't communicate while playing, implying Bob's answer doesn't affect Alice's answer and vice-versa.

Games

- A game \mathcal{G} is played by two players, Alice and Bob.
- Alice and Bob, are given questions, x, y , by a referee, where $x, y \in \mathcal{I}$. Alice and Bob answer with $a, b \in \mathcal{O}$
 - Where \mathcal{I} and \mathcal{O} are finite sets.
- Winning the game is defined by a rule function $\lambda : \mathcal{O} \times \mathcal{O} \times \mathcal{I} \times \mathcal{I} \rightarrow \{0, 1\}$
 - If $\lambda(a, b, x, y) = 0$, then Alice and Bob lose.
 - If $\lambda(a, b, x, y) = 1$, then Alice and Bob win.
- We use the following to define \mathcal{G} 's parameters: $\mathcal{G} = (\mathcal{O}, \mathcal{I}, \lambda)$.

Non-Local Games

- We say a game $\mathcal{G} = (\mathcal{O}, \mathcal{I}, \lambda)$ is non-local if Alice and Bob can't communicate while playing, implying Bob's answer doesn't affect Alice's answer and vice-versa.
 - i.e. $P_A(a|x) = \sum_{b \in \mathcal{O}} P(a, b|x, y)$ is independent of y , and

$$P_B(b|y) = \sum_{a \in \mathcal{O}} P(a, b|x, y)$$
 is independent of x .

Synchronous Non-Local Games

- A non-local game is synchronous if whenever Alice and Bob are given the same question $x \in \mathcal{I}$ then

Synchronous Non-Local Games

- A non-local game is synchronous if whenever Alice and Bob are given the same question $x \in \mathcal{I}$ then
 - $\lambda(a, b, x, x) = 0$ if $a \neq b$

Synchronous Non-Local Games

- A non-local game is synchronous if whenever Alice and Bob are given the same question $x \in \mathcal{I}$ then
 - $\lambda(a, b, x, x) = 0$ if $a \neq b$
 - i.e. If they give different answers they lose

.

Definition

The Mermin-Peres Magic Square Game is a synchronous non-local game. In this game, Alice and Bob will each receive one equation from the following list:

$$x_1 + x_2 + x_3 \equiv 0 \pmod{2}$$

$$x_4 + x_5 + x_6 \equiv 0 \pmod{2}$$

$$x_7 + x_8 + x_9 \equiv 0 \pmod{2}$$

$$x_1 + x_4 + x_7 \equiv 0 \pmod{2}$$

$$x_2 + x_5 + x_8 \equiv 0 \pmod{2}$$

$$x_3 + x_6 + x_9 \equiv 1 \pmod{2}$$

Definition

The Mermin-Peres Magic Square Game is a synchronous non-local game. In this game, Alice and Bob will each receive one equation from the following list:

$$x_1 + x_2 + x_3 \equiv 0 \pmod{2}$$

$$x_4 + x_5 + x_6 \equiv 0 \pmod{2}$$

$$x_7 + x_8 + x_9 \equiv 0 \pmod{2}$$

$$x_1 + x_4 + x_7 \equiv 0 \pmod{2}$$

$$x_2 + x_5 + x_8 \equiv 0 \pmod{2}$$

$$x_3 + x_6 + x_9 \equiv 1 \pmod{2}$$

Alice and Bob fill out their assigned equations with 0's and 1's simultaneously without knowing how the other fills theirs out. Their goal is to satisfy each of their own equations while producing the same value for any shared variable.

The Magic Square game can be represented by a 3x3 grid by labeling each cell like so:

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

$$x_1 + x_2 + x_3 = 0$$

$$x_4 + x_5 + x_6 = 0$$

$$x_7 + x_8 + x_9 = 0$$

x_1	x_2	x_3
+	+	+
x_4	x_5	x_6
+	+	+
x_7	x_8	x_9
=	=	=
0	0	1

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

Bob

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

Bob

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9



Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

Bob

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9



Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$

Bob

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$

1

Bob

x_1	x_2	x_3
x_4	x_5	x_6
x_7	x_8	x_9

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$

$$1 + 0 + 1 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$

$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 0 + 1 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 0 + 1 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 0 + 1 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 0	x_6 1
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 0 + 1 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 1	x_6 0
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 1 + 0 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Magic Square Walkthrough

Alice

x_1	x_2	x_3
x_4 1	x_5 1	x_6 0
x_7	x_8	x_9

$$0 = x_4 + x_5 + x_6$$



$$1 + 1 + 0 \equiv 0 \pmod{2}$$

Bob

x_1	x_2	x_3 1
x_4	x_5	x_6 0
x_7	x_8	x_9 0

$$\begin{aligned} &x_3 \\ &+ \\ &x_6 \\ &+ \\ &x_9 \\ &= \\ &1 \end{aligned}$$



$$1 + 0 + 0 \equiv 1 \pmod{2}$$

Classical Strategy for Magic Square Game Strategy

Definition

A classical strategy for a Non-Local Game is represented by a function $f : \mathcal{I} \rightarrow \mathcal{O}$.

For the Magic Square Game, we can define a classical strategy using the grid below. The best strategy has them agree upon a set table of answers, this allows them to win most of the time. It is impossible to create a solution set that satisfies all nine equations at the same time, otherwise they would be able to win every single game.

x_1 1	x_2 0	x_3 1	$1 + 1 = 0$
x_4 1	x_5 1	x_6 0	$1 + 1 = 0$
x_7 0	x_8 1	x_9 0	$1 \neq 0$


1	1	1
+	+	+
1	1	0
=	=	=
0	0	1

Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.


Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.




Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.
- If vertices $v_1, v_2 \in V(G)$ have an edge $e \in E(G)$ that connects, v_1 to v_2 we say v_1 and v_2 are adjacent.




Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.
- If vertices $v_1, v_2 \in V(G)$ have an edge $e \in E(G)$ that connects v_1 to v_2 we say v_1 and v_2 are adjacent.
- To color a graph, we assign colors to each vertex of a graph such that no adjacent vertices share the same color.




Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.
- If vertices $v_1, v_2 \in V(G)$ have an edge $e \in E(G)$ that connects, v_1 to v_2 we say v_1 and v_2 are adjacent.
- To color a graph, we assign colors to each vertex of a graph such that no adjacent vertices share the same color.
- We say that a graph is k -colorable if the minimum amount of colors needed to color the graph is k .



Definitions

- A graph, G , is a collection of vertices, $V(G)$, and a collection of edges, $E(G)$, that connect a possibly empty subset of vertices.
- If vertices $v_1, v_2 \in V(G)$ have an edge $e \in E(G)$ that connects, v_1 to v_2 we say v_1 and v_2 are adjacent.
- To color a graph, we assign colors to each vertex of a graph such that no adjacent vertices share the same color.
- We say that a graph is k -colorable if the minimum amount of colors needed to color the graph is at most k .



Game Algebra

- We can represent the rules of the Mermin-Peres Magic Square Game using an algebra $\mathcal{A}(\mathcal{G})$. We use $a \in \mathcal{O}$ and $x \in \mathcal{I}$ to index our elements. $\mathcal{A}(\mathcal{G})$ has elements that satisfy the following.

Game Algebra

- We can represent the rules of the Mermin-Peres Magic Square Game using an algebra $\mathcal{A}(\mathcal{G})$. We use $a \in \mathcal{O}$ and $x \in \mathcal{I}$ to index our elements. $\mathcal{A}(\mathcal{G})$ has elements that satisfy the following.
- For all $x \in \mathcal{I}$, we have $\sum_{a \in \mathcal{O}} e_{a,x} = 1$.

$$e_{1,2} \longrightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

Game Algebra

- We can represent the rules of the Mermin-Peres Magic Square Game using an algebra $\mathcal{A}(\mathcal{G})$. We use $a \in \mathcal{O}$ and $x \in \mathcal{I}$ to index our elements. $\mathcal{A}(\mathcal{G})$ has elements that satisfy the following.
- For all $x \in \mathcal{I}$, we have $\sum_{a \in \mathcal{O}} e_{a,x} = 1$.
- For $x, y \in \mathcal{I}$ and $a, b \in \mathcal{O}$ we have $\lambda(a, b, x, y) = 0$ implies $e_{a,x} e_{b,y} = 0$.

$$e_{1,2} \longrightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

Game Algebra

- We can represent the rules of the Mermin-Peres Magic Square Game using an algebra $\mathcal{A}(\mathcal{G})$. We use $a \in \mathcal{O}$ and $x \in \mathcal{I}$ to index our elements. $\mathcal{A}(\mathcal{G})$ has elements that satisfy the following.
- For all $x \in \mathcal{I}$, we have $\sum_{a \in \mathcal{O}} e_{a,x} = 1$.
- For $x, y \in \mathcal{I}$ and $a, b \in \mathcal{O}$ we have $\lambda(a, b, x, y) = 0$ implies $e_{a,x} e_{b,y} = 0$.
- For all $x \in \mathcal{I}$ and $a \in \mathcal{O}$ we have $e_{a,x}^2 = e_{a,x} = e_{a,x}^*$

$$e_{1,2} \longrightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & -\frac{1}{2} \\ 0 & 0 & -\frac{1}{2} & \frac{1}{2} \end{bmatrix}$$

Definition


To make a graph quantum 3-colorable we assign 3-tuples of matrices to each vertex. For any vertices $v_1, v_2 \in V(G)$ that are adjacent these matrices, element by element, are orthogonal.

Definition

To make a graph quantum 3-colorable we assign 3-tuples of matrices to each vertex. For any vertices $v_1, v_2 \in V(G)$ that are adjacent these matrices, element by element, are orthogonal.


Example

$$\begin{aligned}
 & (e_{2,x} + e_{4,x}, e_{1,x}, e_{3,x}) \cdot (e_{1,x} + e_{3,x}, e_{4,x}, e_{2,x}) \\
 &= ((e_{2,x} + e_{4,x})(e_{3,x} + e_{1,x}), e_{1,x}e_{4,x}, e_{3,x}e_{2,x}) \\
 &= (0, 0, 0).
 \end{aligned}$$




Definitions

- Gadgets are subgraphs that encode the game rules into our graph.




Definitions

- Gadgets are subgraphs that encode the game rules into our graph.
- The vertices in red we denote as $V_{a,x}$, with $a \in \mathcal{O}$ and $x \in \mathcal{I}$. These sets are the same as our input and output sets from the Mermin-Peres Magic Square Game.




Definitions

- Gadgets are subgraphs that encode the game rules into our graph.
- The vertices in red we denote as $V_{a,x}$, with $a \in \mathcal{O}$ and $x \in \mathcal{I}$. These sets are the same as our input and output sets from the Mermin-Peres Magic Square Game.
- Other Gadgets such as E_λ and F_λ subgraphs enforce the graph to not be classically 3-colorable.




Definitions

- Gadgets are subgraphs that encode the game rules into our graph.
- The vertices in red we denote as $V_{a,x}$, with $a \in \mathcal{O}$ and $x \in \mathcal{I}$. These sets are the same as our input and output sets from the Mermin-Peres Magic Square Game.
- Other Gadgets such as E_λ and F_λ subgraphs enforce the graph to not be classically 3-colorable.
- We call $V_{a,x}$ and $V_{b,y}$ incompatible whenever $\lambda(a, b, x, y) = 0$.





For vertices $V_{a,x}$ and $V_{b,y}$ who are incompatible, $\lambda(a, b, x, y) = 0$, and are not orthogonal to one another we create one of the subgraphs,


if $(a, b, x, y) \in \mathcal{E}_\lambda$



if $(a, b, x, y) \in \mathcal{F}_\lambda$









297 vertices

Mermin-Peres Graph





110 vertices

Thank you!

Questions?