

CTA200 Computing Project

Caleb Lammers

May 19 2020

Introduction

The light emitted from a galaxy provides information about its physical properties, including both its star formation history and stellar mass. Light emitted from galaxies in different wavelengths contain different information about the galaxy's composition. For instance, it is well known that younger stars emit shorter wavelength light; that is, bluer light than old stars. Thus, if a galaxy emits much more shorter wavelength light, this galaxy likely has many younger stars and thus was forming stars not so long ago. Because different wavelengths help provide different insights, it is useful to represent a galaxy's emitted light as a graph of luminosity/flux density vs. wavelength, called the galaxy's Spectral Energy Distribution (SED).




Figure 1: Example SED of flux density vs. wavelength

By combining knowledge on stellar evolution, calibrated stellar libraries, initial mass functions (IMF), and dust in the interstellar medium, it is possible to predict the SED of galaxies, based on their physical properties. This is known as stellar population synthesis (SPS); a model that combines these ingredients

is called a stellar population synthesis model. In this project, the FSPS (Conroy and Gunn 2009) SPS model was used, particularly the python package python-FSPS. Being able to compute a galaxy's SED based on its physical properties is a step in the right direction, however, as astronomers, we would like to be able to determine a galaxy's physical properties from its observable SED. This project will consist of two parts. In the first, the FSPS model was set up and functions were created so that SEDs and other important graphs could be made. In the second, the goal was to work backwards and determine the physical properties of two galaxies, given their SEDs, which was done using brute force and Markov chain Monte Carlo (MCMC) methods.

Description of Script

Part 1

A StellarPopulation object was initialized in accordance with the settings provided in question 4. Now, by using the built in get spectrum method, SEDs of (bolometric) luminosity in L_{\odot}/Hz vs. wavelength in \AA can be made.




Figure 2: SED of luminosity vs. wavelength

SEDs are more commonly presented as flux density (in μJy) vs wavelength (in \AA), so, next a function was written to convert the SED from units of luminosity into units of spectral flux, according to the equation below.

$$f_v = \frac{L_{(1+z)v}}{4\pi D_L^2}$$

Figure 3: Flux density from luminosity

Because f_v depends on $(1 + z)v$ rather than v , instead of converting each f_v into L_v , it is easier to convert f_v to $L_{(1+z)v}$ and scale each λ by $(1 + z)$. Now, SEDs of this galaxy can be created in units of spectral flux by specifying a redshift. Note that at all redshifts we expect the SED to have similar behaviour as above because we have only rescaled the x and y axes.




Figure 4: SED of flux density vs. wavelength at redshift 1




Figure 5: SED of flux density vs. wavelength at redshift 2




Figure 6: SED of flux density vs. wavelength at redshift 3

Next, the goal was to fix $\lambda = 8000\text{\AA}$ and make a plot of flux density vs. redshift. Note that at each z the generated SED is not guaranteed to have a flux density value at $\lambda = 8000\text{\AA}$. To get around this, the wavelength closest to 8000\AA was linearly interpolated to find a reasonable value for flux density, giving the following graph. As you would expect, flux density decreases as redshift increases.




Figure 7: Flux density vs. redshift at $\lambda = 8000\text{\AA}$

Finally, a function to generate the SED in flux density (μJy) vs. wavelength (\AA) for a galaxy of any age at any redshift was created. Using this function, the following graphs were made.




Figure 8: SED of flux density vs. wavelength for galaxy of age 0.01 Gyr at redshift 1




Figure 9: SED of flux density vs. wavelength for galaxy of age 0.1 Gyr at redshift 1




Figure 10: SED of flux density vs. wavelength for galaxy of age 1 Gyr at redshift 1




Figure 11: SED of flux density vs. wavelength for galaxy of age 3 Gyr at redshift 1




Figure 12: SED of flux density vs. wavelength for galaxy of age 10 Gyr at redshift 1

Part 2

In order to fit a straight line to the given data, it was assumed that the error is Gaussian so that the use of weighted linear least-square fitting is justified (see answer to 2.1 for more motivation). Using linear algebra, a function to determine the slope and y-intercept (along with uncertainties) was made. Additionally, a function to calculate χ^2 was created. Then, the line may be plotted and its fit may be quantitatively measured.

$$b = -7.390^{+2.158}_{-2.158}$$

$$m = 5.277^{+0.451}_{-0.451}$$




Figure 13: Optimal line to fit the given data. $\chi^2 = 17,847$

Similarly, in order to estimate the age and redshift of the galaxy, we must have a way to quantify the goodness of fit of a theoretical SED to a noisy observed SED. By the analogy to fitting a straight line, a similar likelihood function was used:

$$Y^2 = - \sum_{i=1}^{N=1} \frac{(y_i - f(x_i))^2}{\sigma_i^2}$$

Figure 14: Likelihood function for the fit between theoretical and observational SEDs

Where now y_i (Fig. 14) represents an observed flux density and $f(x_i)$ is the theoretical flux density. Note that the negative sign is required because a higher likelihood must imply a closer fit. With this definition in place, we can now compare how a galaxy with age 0.1 Gyr at $z = 1.0$ compares with galaxy 1 and galaxy 2's SEDs. This galaxy had a likelihood of -3121221711061.225 with galaxy 1 and a likelihood of -364732.5275292772 with galaxy 2.




Figure 15: given SED of galaxy 1




Figure 16: given SED of galaxy 2

Clearly, a galaxy with age 0.1 Gyr (Fig. 9) and $z = 1.0$ has a more similar SED to galaxy 2, which makes sense when comparing the SED of galaxy 2 to the SED of this galaxy (above, include figure). By using the likelihood function, we may also try to brute force our way to an estimate of the proper age and z of these galaxies. That is, we may loop through a number of combinations of age and z to find which combination has the highest likelihood. Using this implemented function, we get that galaxy 1 has age of around 6.4 Gyr and redshift of 0.8 and galaxy 2 has age of around 3.4 Gyr and redshift 0.2.




Figure 17: SED of closest galaxy to galaxy 1 found by brute force




Figure 18: SED of closest galaxy to galaxy 2 found by brute force

As you can see, the graphs are somewhat similar but have their discrepancies. Before continuing, one should note that this is not a great way of approximating the age and redshifts of these galaxies. In fact, we have no guarantees that we are even close to correct, we may have simply found a local max of the likelihood function. To do this in a more intelligent way, Markov chain Monte Carlo (MCMC) methods were used. First, MCMC was applied to the example of fitting a line. Of course, applying MCMC to the the χ^2 model is pointless because we have an analytic solution. Instead, we now adopt a more complex model with the assumption that the errors were somewhat underestimated.

$$\ln(p) = -\frac{1}{2} \sum_{i=1}^{N=1} \frac{(y_i - f(x_i))^2}{s_i^2} + \ln(2\pi s_i^2)$$

$$s_i^2 = \sigma_i^2 + f_0^2 f(x_i)$$

Figure 19: Likelihood function for new model

Notice that this model has a so-called nuisance parameter f_0^2 . Because it is now not clear that there is an analytical solution or even how f , m , and b (of course, $f(x_i) = m * x_i + b$) should relate, MCMC is useful. To use MCMC, we define a uniform prior probability function, and use python emcee's features. With 32 walkers centered around the values found from weighted linear least-square fitting and 10000 steps we get the following Markov chain.




Figure 20: Markov chain with emcee

According to the autocorrelation time, it takes about 40 steps for the Markov chain to burn in. Removing the first 40 steps, we get the following corner plots.




Figure 21: Markov chain with emcee

These plots show that b and m are closely related while b and f/m and f/b are not. The fact that f takes on low values shows that the errors are likely not underestimated by very much, if at all. Nonetheless, MCMC gives the following values for b , m , and f and graph.

$$m = 5.261_{-0.455}^{+0.444}$$

$$b = -7.391_{-2.108}^{+2.172}$$

$$f = 0.052_{-0.037}^{+0.056}$$




Figure 22: Markov chain with emcee

The slope, y-int and uncertainties of this line are similar to the those from the weighted linear least-square fitting line. A similar method was used to find

optimal age and redshift for galaxy 1 and 2. Again, a uniform prior was used and the same likelihood function from above was used. When run with 2500 steps and 16 walkers centered around the estimated ages and redshifts, the walkers did not move around the space - they got stuck around the estimated values. So, instead, the walkers were started in random locations distributed around the estimated values. Here are the Markov chains and corner plots.




Figure 23: Markov chain of galaxy 1 data with emcee




Figure 24: Corner graph of galaxy 1 data with emcee

$$\ln(\text{age}) = 1.608_{-0.002}^{+0.105}$$

$$z = 0.853_{-0.024}^{+0.016}$$

Note that the error is likely somewhat higher than calculated here (which is due to how the distribution looks).




Figure 25: Markov chain of galaxy 2 data with emcee




Figure 26: Corner graph of galaxy 2 data with emcee

$$\ln(\text{age}) = 1.843^{+0.559}_{-0.001}$$

$$z = 0.162^{+0.000}_{-0.029}$$

Due to the difficulty of calculation, the Markov chain and corner plots were not as continuous but it is interesting how they still find a similar answer to the brute force method

The next step would be to add stellar mass and dust attenuation to the free variables in MCMC. Though it would be easy to extend the code to do

this, the MCMC calculations above already took several hours, so, this would undoubtedly take unreasonably long in the current setup. To make this feasible, one could use the Scinet cluster or try to further optimize the code.

Question Answers

1.0. As briefly mentioned in the introduction, stellar population synthesis (SPS) refers to combining information on stellar evolution, calibrated stellar libraries, initial mass functions (IMF), and dust in the interstellar medium to construct the SED of a galaxy. Many different SPS models exist, and they have their own assumptions, tools, and calibrations. By using an SPS model, specifying a galaxy's physical properties allows for reconstruction of its SED. Of course, this is only half the battle because we would like to go from the SED to a galaxy's physical properties. There have been many approaches to solving this problem, including Monte Carlo methods, machine learning, and numerical methods. No matter the tool used, the role of the SPS model is the same. The SPS model will generate SEDs given specified physical parameters which will be compared to the SED of the desired galaxy (there are many ways to "compare," of course). Then, adjustments will be made to the physical parameters in hopes of finding a better fit between the generated and desired SED. In this way, SPS models provide the foundation for this method of determining a galaxy's physical properties from its emitted light.

1.3. The StellarPopulation can take in a variety of inputs about the galaxy, including its dust content, initial mass function, metallicity, the type of star formation history and more.

1.4. As discussed above, the units of the output spectrum are solar luminosity per hertz (L_{\odot}/Hz). The StellarPopulation object is using MESA Isochrones and Stellar Tracks (MIST) and the MILES spectral library. At 100 Myr, the stellar mass of the galaxy is about $0.92 M_{\odot}$.

1.7. As galaxies cannot be older than the universe, when a galaxy is observed at some redshift z , we are effectively looking back in time t at its light. Then, $t + t_g < t_u$ (there are equations to calculate t from z) where t_u is about 13.8 billion years. So, the larger a galaxy's redshift, the smaller its possible maximum age is.

2.1 There are many possible techniques to fit a straight line to a set of sampled points. As Hogg, Lang, and Bovy point out, the standard method when there is negligible error in the x data and known Gaussian error in the y data is weighted linear least-square fitting. This type of fitting minimizes χ^2 which can be intuitively understood as a measure of the line from the data. It is mathematically defined as:

$$\chi^2 = \sum_{i=1}^{N=1} \frac{(y_i - f(x_i))^2}{\sigma_{y_i}^2}$$

Part of the motivation for using this measure of goodness of fit is that it has an easily calculable minimum with linear algebra. With this method, we get a reasonable looking line of best fit. It is difficult to say whether a line is the best curve to represent the data without any physical knowledge about the data, however, a line appears to fit the data well with a relatively low χ^2 .