



CamTrapAsia: 210 full wildlife capture lists from camera trapping studies

Journal:	<i>Ecology</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Data Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Pereira Mendes, Calebe; Nanyang Technological University, Asian School of the Environment Thapa, Arjun; University of Queensland, School of the Environment Albert, Wido; Fauna & Flora International Shia, Amanda; Hutan Amir, Zachary; The University of Queensland, School of the Environment Ancrenaz, Marc; Hutan Ash, Eric; WildCRU Azhar, Badrul; Universiti Putra Malaysia, Department of Forest Science and Biodiversity, Faculty of Forestry and Environment Bernard, Henry; Universiti Malaysia Sabah Brodie, Jedediah; University of Montana, Wildlife Biology Carr, Elliot; The University of Queensland Clements, Gopalasamy Reuben; Sunway University Davies, Glyn; World Wildlife Fund Malaysia, Sabah Landscape Programme Deere, Nicolas; University of Kent, Durrell Institute of Conservation and Ecology (DICE) Dinata, Yoan; The Zoological Society of London, Indonesia program Donnelly, Christl; University of Oxford, Department of Statistics Duangchantrasiri, Somphot; Department of National Parks, Plant, and Wildlife Conservation, Wildlife Research Division Fredriksson, Gabriella ; Pro Natura Foundation Goossens, Benoit; Cardiff University, School of Biosciences Granados, Alys; Felidae Conservation Fund Hearn, Andrew; WildCRU Hon, Jason; World Wildlife Fund Malaysia, Malaysia program Hughes, Tom; Conservation Medicine Jansen, Patrick; Wageningen University, Wildlife Ecology & Conservation; Smithsonian Tropical Research Institute, Kawanishi, Kae; Malaysian Conservation Alliance for Tigers (MYCAT) Kinnaird, Margaret; Mpala Research Center Koh, Sharon; World Wildlife Fund Malaysia, Malaysia program Latinne, Alice; Wildlife Conservation Society, Viet Nam Country Program Linkie, Matthew; Wildlife Conservation Society, Indonesia program Loi, Federica; Istituto Zooprofilattico Sperimentale della Sardegna G Pegreff, Regional Veterinary Epidemiological Observatory Lynam, Anthony; Wildlife Conservation Society, Thailand program Meijaard, Erik; Borneo Futures</p>

	<p>Mohd-Azlan, Jayasilan; Universiti Malaysia Sarawak, Department of Biology</p> <p>Moore, Jonathan; University of East Anglia, ; SUSTech,</p> <p>Nathan, Senthilvel; Sabah Wildlife Department, Department of Veterinary Services</p> <p>Ngoprasert, Dusit; King Mongkut's University of Technology Thonburi, Conservation Ecology Program</p> <p>Novarino, Wilson; Andalas University, Department of Biology</p> <p>Nursamsi, Ilyas; The University of Queensland, School of the Environment</p> <p>O'Brien, Tim; Wildlife Conservation Society, Global Conservation Program Department</p> <p>Ong, Robert; Sabah State Government, Department of Veterinary Services</p> <p>Payne, John; Sabah State Government, Department of Veterinary Services</p> <p>Priatna, Dolly; Pakuan University, Graduate School of Environmental Management</p> <p>Rayan, D. Mark; Wildlife Conservation Society, Malaysia Program</p> <p>Reynolds, Glen; South East Asia Rainforest Research Partnership, Conservation Programme</p> <p>Rustam, Rustam; Mulawarman University, Faculty of Forestry</p> <p>Selvadurai, Sasidhran; Universiti Putra Malaysia, Department of Forest Science and Biodiversity, Faculty of Forestry and Environment</p> <p>Silmi, Muhammad; United Plantations Berhad, Biodiversity Division</p> <p>Sinovas, Pablo; Fauna and Flora International, Cambodia Programme</p> <p>Sribuarod, Kriangsak; Royal Thai Government Department of National Park Wildlife and Plant Conservation, Khlong Saeng Wildlife Research Station</p> <p>Steinmetz, Robert; World Wildlife Fund, Cambodia program</p> <p>Struebig, Matthew; University of Kent, Durrell Institute of Conservation and Ecology (DICE)</p> <p>Sukmasuang, Ronglarp; Kasetsart University</p> <p>Sunarto, Sunarto; World wildlife Fund, Indonesia program</p> <p>Tarmizi, Tarmizi; Leuser International Foundation</p> <p>Traeholt, Carl; Copenhagen Zoo, Research and Conservation Division</p> <p>Wearn, Oliver; Fauna and Flora International, Vietnam Programme</p> <p>Wibisono, Hariyo; San Diego Zoo, Conservation Programme</p> <p>Wilting, Andreas; Leibniz Institute for Zoo and Wildlife Research, Department of Ecological Dynamics</p> <p>Wong, Seth; Leibniz Institute of Zoo and Wildlife Research, Department of Ecological Dynamics</p> <p>Wong, Siew Te ; Bornean Sun Bear Conservation Centre</p> <p>Word, Jettie; Borneo Project</p> <p>Chiok, Wen; Nanyang Technological University, Asian School of the Environment</p> <p>Luskin, Matthew; The University of Queensland, School of Biological Sciences;</p>
Substantive Area:	Conservation < Population Ecology < Substantive Area, Distributions/Patchiness/Marginal Populations < Population Dynamics and Life History < Population Ecology < Substantive Area, Data paper < Data < Substantive Area
Organism:	Vertebrates < Animals, Mammals < Vertebrates < Animals, Birds (specify type in field below) < Vertebrates < Animals, Reptiles < Vertebrates < Animals
Habitat:	Rain Forest < Subtropical Zone < Terrestrial < Habitat, Tropical Zone < Terrestrial < Habitat
Geographic Area:	Southeast Asia < Asia < Geographic Area

Key words/phrases:	data paper, vertebrate, mammal, bird, richness, biodiversity, IUCN Red List, Threatened, Endangered, distribution, Animal, Trail camera
Abstract:	<p>Southeast Asia holds some of the most diverse rainforests on the planet. Despite their importance, information on biodiversity is scattered across published, peer-reviewed, and gray literature and in unpublished raw data. Camera traps are an effective non-invasive method of surveying vertebrates. In this data paper, we compiled and standardized 210 camera trap surveys from across Southeast Asia. The camera deployments included in this study used relatively standardized methods, providing a consistent and reliable data set relative to other large-scale occurrence datasets such as online citizen science repositories. The complete data set comprises 276,805 records of 360 species (225 mammals, 129 birds, and 6 reptiles), making it one of the most extensive and comprehensive biodiversity inventories for the region. The information detailed in this data paper opens opportunities for single-species ecological or conservation studies as well as community ecology and macroecology investigations. For example, the dataset may be useful to understand the effects of habitat loss, fragmentation, climate change, and other human-mediated processes on species and communities. for future ecological research that could be replicated through time and in other regions.</p>
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
Data S1.rar	

Title: CamTrapAsia: 210 full wildlife capture lists from camera trapping studies

Authors:

Mendes, Calebe P¹; Thapa, Arjun²; Albert, Wido R³; Amanda, Shia K. P. ⁴; Amir, Zachary²; Ancrenaz, Marc⁴; Ash, Eric⁵; Azhar, Badrul⁶; Bernard, Henry⁷; Brodie, Jedediah⁸; Carr, Elliot²; Clements, Gopalasamy Reuben⁹; Davies, Glyn¹⁰; Deere, Nicolas J¹¹; Dinata, Yoan¹²; Donnelly, Christl A¹³; Duangchantrasiri, Somphot¹⁴; Fredriksson, Gabriella¹⁵; Goossens, Benoit¹⁶; Granados, Alys¹⁷; Hearn, Andrew⁵; Hon, Jason¹⁸; Hughes, Tom¹⁹; Jansen, Patrick²⁰; Kawanishi, Kae²¹; Kinnaird, Margaret²²; Koh, Sharon¹⁸; Latinne, Alice²³; Linkie, Matthew²⁴; Loi, Federica²⁵; Lynam, Anthony J²⁶; Meijaard, Erik²⁷; Mohd-Azlan, Jayasilan²⁸; Moore, Jonathan H²⁹; Nathan, Senthilvel KSS³⁰; Ngoprasert, Dusit³¹; Novarino, Wilson³²; Nursamsi, Ilyas²; O'Brien, Timothy³³; Ong, Robert³⁰; Payne, John³⁰; Priatna, Dolly³⁴; Rayan, Mark³⁵; Reynolds, Glen³⁶; Rustam, Rustam³⁷; Selvadurai, Sasidhran⁶; Silmi, Muhammad³⁸; Sinovas, Pablo³⁹; Sribuarod, Kriangsak⁴⁰; Steinmetz, Robert⁴¹; Struebig, Matthew J¹¹; Sukmasuang, Ronglarp⁴²; Sunarto, Sunarto⁴³; Tarmizi, Tarmizi⁴⁴; Traeholt, Carl⁴⁵; Wearn, Oliver R⁴⁶; Wibisono, Hariyo Beebach⁴⁷; Wilting, Andres⁴⁸; Wong, Seth Timothy⁴⁸; Wong, Siew Te⁴⁹; Word, Jettie⁵⁰; Xuan, Chiok Wen¹; Luskin, Matthew Scott⁵¹

Author Affiliations:

1. Asian School of the Environment, Nanyang Technological University
2. School of the Environment, University of Queensland, Australia
3. Fauna & Flora International, Indonesia
4. HUTAN, Malaysia
5. WildCRU, Oxford, England
6. Department of Forest Science and Biodiversity, Faculty of Forestry and Environment, Universiti Putra Malaysia
7. Institute for Tropical Biology and Conservation, Universiti Malaysia Sabah
8. Wildlife Biology, University of Montana
9. Department of Biological Sciences, Sunway University
10. Sabah Landscape Programme, World Wildlife Fund
11. Durrell Institute of Conservation and Ecology (DICE), University of Kent
12. Indonesia program, Zoological Society of London

- 32 13. Department of Statistics, University of Oxford
- 33 14. Wildlife Research Division, Department of National Parks, Plant, and Wildlife Conservation,
- 34 Bangkok, Thailand
- 35 15. Pro Natura Foundation
- 36 16. School of Biosciences, Cardiff University
- 37 17. Felidae Conservation Fund
- 38 18. Malaysia program, World Wildlife Fund
- 39 19. Conservation Medicine
- 40 20. Wildlife Ecology & Conservation, Wageningen University
- 41 21. Malaysian Conservation Alliance for Tigers (MYCAT).
- 42 22. Mpala Research Centre
- 43 23. Viet Nam Country Program, Wildlife Conservation Society
- 44 24. Indonesia program, Wildlife Conservation Society
- 45 25. Regional Veterinary Epidemiological Observatory, Istituto Zooprofilattico Sperimentale
- 46 della Sardegna
- 47 26. Thailand program, Wildlife Conservation Society
- 48 27. Borneo Futures, Brunei Darussalam
- 49 28. Department of Biology, Universiti Malaysia Sarawak
- 50 29. Environmental Science, SUSTech University
- 51 30. Department of Veterinary Services, Sabah Government
- 52 31. Conservation Ecology, King Mongkut's University of Technology Thonburi
- 53 32. Department of Biology, Andalas University
- 54 33. Global Conservation Program Department, Wildlife Conservation Society
- 55 34. Graduate School of Environmental Management, Pakuan University
- 56 35. Malaysia program, Wildlife Conservation Society Malaysia Program
- 57 36. Conservation Programme, SEARRP
- 58 37. Fakulty of Forestry, Mulawarman University
- 59 38. Biodiversity Division, United Plantations Berhad / PT Surya Sawit Sejati
- 60 39. Cambodia Programme, Fauna & Flora International
- 61 40. Klongsang Wildlife Research Station, Department of National Park Wildlife and Plant
- 62 41. Cambodia program, World Wildlife Fund

42. Kasetsart University

43. Indonesia program, World Wildlife Fund

44. Leuser International Foundation

45. Research and Conservation Division, Copenhagen Zoo

46. Vietnam Programme, Fauna & Flora International

47. Conservation Programme, San Diego Zoo

48. Department of Ecological Dynamics, Leibniz Institute of Zoo and Wildlife Research

49. Bornean Sun Bear Conservation Centre

50. Borneo Project

51. Department of Biological Sciences, University of Queensland

Corresponding Author:

Luskin, Matthew Scott <mattluskin@gmail.com>

Open Research statement:

The data and code are available in the GitHub directory
<<https://github.com/CalebePMendes/CamTrapAsia.git>>

Class I. Data Set Descriptors

A. Data set identity:

CamTrapAsia: 210 full wildlife capture lists from camera trapping studies

B. Data set identification code:

Metadata S1

Data S1.zip

A. Data set description

Originators:

Mendes, Calebe Pereira – Asian School of the environment, Nanyang Technological University – calebepm3@gmail.com

Tharpa, Arjun – School of the Environment, University of Queensland, Australia – tharpa.nature@gmail.com

Luskin, Matthew – School of Biological Sciences, University of Queensland – m.luskin@uq.edu.au

Abstract:

Southeast Asia holds some of the most diverse rainforests on the planet. Despite their importance, information on biodiversity is scattered across published, peer-reviewed, and gray literature and in unpublished raw data. Camera traps are an effective non-invasive method of surveying vertebrates. In this data paper, we compiled and standardized 210 camera trap surveys from across Southeast Asia. The camera deployments included in this study used relatively standardized methods, providing a consistent and reliable data set

relative to other large-scale occurrence datasets such as online citizen science repositories. The complete data set comprises 276,805 records of 360 species (225 mammals, 129 birds, and 6 reptiles), making it one of the most extensive and comprehensive biodiversity inventories for the region. The information detailed in this data paper opens opportunities for single-species ecological or conservation studies as well as community ecology and macroecology investigations. For example, the dataset may be useful to understand the effects of habitat loss, fragmentation, climate change, and other human-mediated processes on species and communities. for future ecological research that could be replicated through time and in other regions.

B. **Key words/phrases:** Animal, vertebrate, mammal, bird, biodiversity, richness, Southeast Asia, distribution, camera trap, trail camera, covariates, IUCN red list

Class II. Research origin descriptors

A. Overall project description:

1. Identity: CamTrapAsia: 210 full wildlife capture lists from camera trapping studies.

2. Originators: Calebe Pereira Mendes, Arjun Tharpa, Matthew Luskin.

3. Period of study: The data range from 1987 to 2020.

4. Site: Southeast Asia

5. Objectives: The CamTrapAsia dataset aims to facilitate the access and use of camera trap data previously scattered in the literature.

6. Sources of funding:

The research was funded by the Smithsonian Institution's ForestGEO program, Nanyang Technological University in Singapore, the University of Queensland (UQ) Centre for Biodiversity and Conservation Science, National Geographic Society #9384-13 and ARC DECRA #DE210101440, and BIFA6_005: Biodiversity Information Fund for Asia "Sharing vertebrate occurrence data from camera traps in Asia".

B. Specific subproject description

1. Rationale:

Camera traps have existed in some form since 1890 and revolutionized wildlife research since the late 1990s (O'Connell, Nichols, and Karanth 2010). There are numerous advantages to cameras, including researcher time or capacity, automatically triggered cameras remove much of the observer bias present in humans. Since 2005, most cameras are digital and detect the heat signature from

animals moving within the sensor range, photos are later sorted, and captures (or detections) are noted when a species is present. Cameras monitor wildlife with minimal disturbance, both day and night, contributing to biodiversity inventories (species richness) and images are time-stamped to allow for investigations on topics like wildlife behaviour (e.g. temporal activity patterns). The development of hierarchical modelling methods including mark-recapture for density and occupancy analyses - and the inclusion of covariates into these models - have given scientists a robust glimpse into the ecology of cryptic animals in their natural environments (O'Connell, Nichols, and Karanth 2010; Sollmann 2018).

Camera traps and associated analyses play an important role in monitoring animal populations during the Anthropocene 6th mass extinction (Ceballos et al. 2015). Over 60% of the earth's forest landscapes suffer some degree of degradation (Grantham et al. 2020) and widespread poaching has driven declines >25% of the vertebrate species (Dirzo et al. 2014). Climate change is the next great threat (IPCC 2014). Addressing these challenges requires far larger data volumes than any single research team or organization can collect or manage. Further, monitoring for global and regional trends requires long time series, so there needs to be systematic handing down of projects and datasets to the next generation of scientists, and ideally redundancy to ensure the continuation of critical longitudinal measurements (Beaudrot et al. 2016). These aims can only be met via collaboration and data sharing. The CamTrapAsia dataset for Southeast Asian tropical and subtropical

forests joins other open online camera trap images and data repositories (Ahumada et al. 2020; Lima et al. 2017).

CamTrapAsia only mobilizes the full capture lists (e.g. we excluded studies that only reported a single species or subset of species) and includes numerous covariates for each study, in a single standardized and accessible format. Providing these covariates is aimed to remove barriers to science. While we endeavoured to exhaustively search the white and grey literature for all available information and requested missing information from authors, we estimate our dataset contains just 20% of the camera trapping conducted to date in the region as much is never published in any form or only selective animal detections are published. There is work yet to be done.

2. Research methods

We searched for Asian camera trap studies using many approaches. We started with a Web of Science using terms “camera*” and Asia* or Thai* or Malaysia* or Indonesia* or Singapore* or Borneo* or Cambodia* or Vietnam* or Lao* or Myanmar* or Burm* or Sumatra* or Borne*. We also used Google Scholar using the terms camera* AND vertebrate* or mammal* or bird* or biodiverse* or richness or Endangered or Threatened or terrestrial or distribution* or abundance* or carnivor* or herbiv* or omnivor* or predator* tiger* or leopard* or rhino* or elephant* or tapir* or deer* or civet*.

Criteria for inclusion of published data were: (i) a complete list of the vertebrate species >1 kg detected, the number of independent records, the trapping effort (with a minimum of 25 trap nights), the number of cameras/stations deployed, the coordinates of the study site and the period when the trapping survey was performed (with a minimum temporal precision of year). The data matching the required criteria was collated in a single table, together with the references needed to locate the original publication. The collated data set was standardized, with all coordinates set to WGS 84, and the species binomial names were verified using the R package Taxize (Chamberlain et al. 2022), based in the taxonomic databases from the National Center for Biotechnology Information and the Global Biodiversity Information Facility (Schoch et al. 2020; GBIF 2022).

To facilitate the usage of this dataset, we also made available a set of 13 spatial covariables including the percentage of forest cover, altitude, terrain roughness, forest landscape integrity index (FLII), human footprint index, ecoregion intactness index, average precipitation, average temperature, night light emissivity, human population, percentage of oil palm, percentage of urban areas and, percentage of protected areas), extracted in 3 spatial scales around the survey locations (10, 20 and 30 km radius). We also added a set of biological traits (adult body mass, percentage of diet composed by invertebrates, percentage of diet composed by vertebrates, percentage of diet composed by plants, diet breadth, trophic level, activity cycle and habitat breadth) for the recorded species extracted from the combine dataset (Soria et al. 2021). Finally, the resulting data set is

provided as a three of .csv tables and as a Darwin Core archive, and available on GBIF. The R code used to assemble the dataset and standardize the species names is also provided.

3. Results

CamTrapAsia contains 210 surveys from 133 sites (as labelled in the original papers) from 89 landscapes (here defined as an area encompassing one or more forest patches within 20 km of each other, and which share similar socio-environmental conditions). The geographical coverage was 11 countries including Indonesia, Singapore, Malaysia, Bhutan, Thailand, Myanmar, Cambodia, Laos, Vietnam, Nepal and far-eastern India (Fig 1 and 2). The temporal coverage spans from 1987 to 2020 (Fig 3). There were 276,805 records from 10,024 camera stations over 561,292 trap nights. A total of 360 species were recorded, from 173 genera and 70 families. Mammals comprised 62.5% of the species (fig 4 and 5), birds comprised 35.8% (fig 6 and 7), and the remaining were reptiles. The most recorded species was *Sus scrofa* with 42,269 records, followed by *Macaca nemestrina* ($n = 32,964$) and *Muntiacus muntjac* ($n = 23,796$). A total of 45 species were recorded just once (singletons). The most recorded families were Cervidae, Suidae and Cercopithecidae, with 52,087, 52,076 and 45,891 records respectively. There were 8 families with species recorded once.

4. Project personnel:

Calebe Pereira Mendes, Arjun Tharpa, Matthew Luskin.

5. Acknowledgements

We would like to acknowledge the incredible work that went into fieldwork to collect this data, including Gregoire Bertagnolio, F. Cheong, Roshan Guharajan, Gurutzeta Guillera, Ferran Jori, FR Khakim, Salwa Khalid, Saurav Hari Kumar, AH Lubis, T Lubiz, Sugesti Marif, Maryati Maryati, Mumponjan Mumponjan, Helga Peters, Alice Porco, Roshan Guharajan, Roslina Ragai, Mila Redkani, N Seuaturien, NM Shwe, Wong Siew, Cedric Tan, Joe Smith, Melvin E Sunquist, Cedric Tan, Shuwoan Teoh, Ha Wahyudi, C.T. Wong, Zulfahmi Zulfahmit. . We also thank Yayasan Sabah, the Sabah Forest Department, the Sabah Biodiversity Council, and the Danum Valley Management Committee, Abdul Hamid, Glen Reynolds, Jedediah Brodie, Katie Doehla, and Tombi Karolus for permission and help conducting fieldwork at Danum Valley. We thank Patrick Jansen, the Smithsonian Institute and the Tropical Ecology Assessment and Monitoring (TEAM) network for collecting data from Pasoh, as well as Yao Tse Leong and the Forest Research Institute Malaysia (FRIM) for permission to work at Pasoh. We thank M. Bt. Mohamad, J. Kulis, and the Sarawak Forestry Department for permission to conduct fieldwork at Lambir Hills and Piyasart ‘Guide’ Kumhom and NTU field ecology courses for help collecting data. We thank S. Lum, A. Loo, M. Khoo, B. Lee, J. Chan, A. Goh, and N. Parks for permission and help with fieldwork in Singapore. We thank W. Chantorn, A. Nathalang, S. Bunyavejchewin, R. Sukmasuang, F. Gutierrez, C. Scanlon for permissions and help at Khao Yai and Khao Ban Tat. In Sumatra, we thank W. R. Albert, M.

Linkie, Y. Dinata, H. Wibisono, and H. Kita for facilitating fieldwork in Sumatra.
We thank E. S. Sembiring, T. Ramadiyanta, E. Ramadiyanta, S., I. Tanjung, and
C. Decky for assistance with fieldwork associated with the Ecological Cascades
Lab.

6. Conflict of Interest Statement

The authors have no conflicts of interest to declare.

For Review Only

Class III. Data set status and accessibility

C. Status

1. Latest update: 03-04-2023
2. Latest archive date: 03-04-2023
3. Metadata status: 13-04-2023
4. Data verification: 13-04-2023

D. Accessibility

1. Storage location and medium:

The data available in the GitHub directory
<<https://github.com/CalebePMendes/CamTrapAsia.git>>

2. Contact persons:

Calebe Pereira Mendes, Asian School of the environment -Nanyang Technological
University, calebepm3@gmail.com
Matthew Luskin, School of Biological Sciences - University of Queensland,
m.luskin@uq.edu.au

3. Copyright restrictions:

The data is fully available to the public for utilization and research. Kindly
acknowledge the data paper when utilizing the data.

Costs:

The data is available for free.

Class IV. Data structural descriptors

A. Data set file

1. Identity:

CamTrapAsia_Captures_20230403.csv

CamTrapAsia_Metadata_20230403.csv

Captures_raw_20230403.csv

Metadata_raw_20230403.csv

species_traits.csv

Dataset merging and Standardization – 20230123.r

2. Size:

904 KB [4458 rows, 16 columns]

212 KB [210 rows, 69 columns]

389 KB [4536 rows, 8 columns]

212 KB [210 rows, 69 columns]

65 KB [369 rows, 18 columns]

11 KB [318 lines]

3. Format and storage mode:

Comma-separated values (.csv)

B. Variable information

306 metadata.csv

Field Name	Description	Range	NAs	Example	Source
survey_id	Unique ID which connects the records from the metadata table with the capture table.	NA	0	Lambir2017.ECL	compiler
region	General region where sampling was performed	NA	0	Southeast_Asia	original authors
country	Country where sampling was performed	NA	0	Indonesia	original authors
site	Name of the location where sampling was performed	NA	0	Leuser_forest_fragments	original authors
effort	Trapping effort, in trap-nights	28 - 32027	0	45	original authors
size_km2	Size of the forest where the sampling was performed, in km ²	0.14 - 278889.37	12	1.80	original authors
Protected_area	Whether the sampling happened within a protected area	"y" for protected areas, "n" for not protected areas and "Mixed" when only a part of the area is protected	22	y	original authors
Y_lat	Latitude in decimal degrees, using WGS84	-8.7 to 33.0	0	3.972066	original authors
X_long	Longitude in decimal degrees, using WGS84	80.2 to 140.1	0	98.08855	original authors
logging	Whether the area was previously logged, not-logged or in a plantation	"logged", "not_logged", "plantation" and "mixed" which includes cameras in both logged and unlogged areas	93	logged	original authors
logging_obs	Observations about logging	text	97	logged	original authors or compiler
edge_1km	Whether the sapling was performed within 1 km from an edge	"edge" for sampling within 1 km of the nearest edge, "interior" for sites farther than 1 km, "both" for	123	edge	original authors

		sampling with cameras in both edge and interior.			
year_start	Year of the start of the trapping survey	1987 - 2020	0	2014	original authors
year_end	Year of the ending of the trapping survey	1988 - 2020	0	2014	original authors
monthstart	Month of the start of the trapping survey	1 to 12	4	1	original authors
monthfinish	Month of the ending of the trapping survey	1 to 12	6	2	original authors
n_points	Number of sampling points in the survey	1 - 310	3	1	original authors
n_cameras	Number of cameras deployed in the survey	1 - 600	0	1	original authors
cam_spacing	Minimum space between cameras, in meters	60 - 4000	46	1000	original authors
area_cover_km2	Area of the minimum convex polygon of the sampling points in a survey, in km ²	0.04 - 1357	89	NA	original authors
indent_cap_mins	Minimum time allowed between two independent records, in minutes	0 - 60	21	60	original authors
forest_type	Type of forest in which the sampling was deployed	NA	77	Evergreen broadleaf	original authors
study_notes	Notes about the study from which the camera trap data was obtained	NA	172	NA	compiler
veg_notes	Notes about the vegetation where the sampling was deployed	NA	72	Frag	NA
study_author	Correspondent author of the study/data	NA	0	Luskin et al. 2017	original authors
Source	Link or citation to the original publication or the author contact.	NA	0	https://www.nature.com/articles/s41467-017-01656-4	original authors

forest_cover_10km	Percentage of area covered by vegetation above 4m in a 10km buffer around the sampling coordinates	0 to 78.22	0	60.26213169	(Sexton et al. 2013)
forest_cover_20km	Percentage of area covered by vegetation above 4m in a 20km buffer around the sampling coordinates	0 to 75.79	0	57.71243796	(Sexton et al. 2013)
forest_cover_30km	Percentage of area covered by vegetation above 4m in a 30km buffer around the sampling coordinates	0 to 74.92	0	54.05373898	(Sexton et al. 2013)
altitude_10km	Average altitude in a 10km buffer around the sampling coordinates, in meters	8 to 4798.25	0	56.88501509	(Jaxa 2015)
altitude_20km	Average altitude in a 20km buffer around the sampling coordinates, in meters	9.39 to 4960.17	0	107.5634248	(Jaxa 2015)
altitude_30km	Average altitude in a 30km buffer around the sampling coordinates, in meters	9.13 to 5005.22	0	219.5147203	(Jaxa 2015)
roughness_10km	Roughness index, calculated using the R function terra::terrain() with option v = "roughness", in 10km buffer around the sampling coordinates, in meters	2.85 to 68.80	0	14.36782614	(Jaxa 2015; Hijmans 2022)
roughness_20km	Roughness index, calculated using the R function terra::terrain() with option v = "roughness", in 20km buffer around the sampling coordinates, in meters	2.62 to 59.13	0	17.58147274	(Jaxa 2015; Hijmans 2022)
roughness_30km	Roughness index, calculated using the R function terra::terrain() with option v = "roughness", in 30km buffer around the sampling coordinates, in meters	2.94 to 56.25	0	19.45138369	(Jaxa 2015; Hijmans 2022)

FLII_10km	Forest Landscape Integrity Index in 10 km buffer around the sampling coordinates, unitless	0 to 9.98	2	5.459989231	(Grantham et al. 2020)
FLII_20km	Forest Landscape Integrity Index in 20 km buffer around the sampling coordinates, unitless	0 to 9.75	1	5.175927237	(Grantham et al. 2020)
FLII_30km	Forest Landscape Integrity Index in 30 km buffer around the sampling coordinates, unitless	0 to 9.63	1	5.403120846	(Grantham et al. 2020)
human_footprint_10km	Human Footprint Index in 10 km buffer around the sampling coordinates, unitless	0.14 to 94.66	0	9.853968254	(Venter et al. 2018)
human_footprint_20km	Human Footprint Index in 20 km buffer around the sampling coordinates, unitless	1.16 to 95.77	0	11.97931583	(Venter et al. 2018)
human_footprint_30km	Human Footprint Index in 30 km buffer around the sampling coordinates, unitless	1.42 to 103.79	0	15.08300954	(Venter et al. 2018)
ecoregion_intactness_10km	Ecoregion Intactness Index in 10 km buffer around the sampling coordinates, unitless	0 to 787.88	0	53.77821012	(Beyer et al. 2020)
ecoregion_intactness_20km	Ecoregion Intactness Index in 20 km buffer around the sampling coordinates, unitless	0 to 661.38	0	57.35282651	(Beyer et al. 2020)
ecoregion_intactness_30km	Ecoregion Intactness Index in 30 km buffer around the sampling coordinates, unitless	0.12 to 636.78	0	87.49029982	(Beyer et al. 2020)
precipitation_10km	Mean annual precipitation in 10km buffer around the sampling coordinates, in mm	585.362637362637 - 4105.63561643836	0	2616.955801	WorldClim 2.1 (http://worldclim.org)
precipitation_20km	Mean annual precipitation in 20km buffer around the sampling coordinates, in mm	567.015068493151 - 4134.70378006873	0	2614.322536	WorldClim 2.1 (http://worldclim.org)

precipitation_30km	Mean annual precipitation in 30km buffer around the sampling coordinates, in mm	567.131466828971 - 3935.06654456654	0	2588.659534	WorldClim 2.1 (http://worldclim.org)
temperature_10km	Mean annual temperature in 10km buffer around the sampling coordinates, in degrees Celsius	-1.81 to 27.98	0	25.97048802	WorldClim 2.1 (http://worldclim.org)
temperature_20km	Mean annual temperature in 20km buffer around the sampling coordinates, in degrees Celsius	-2.36 to 28.04	0	25.72089934	WorldClim 2.1 (http://worldclim.org)
temperature_30km	Mean annual temperature in 30km buffer around the sampling coordinates, in degrees Celsius	-2.26 to 28.03	0	25.14833225	WorldClim 2.1 (http://worldclim.org)
nighttime_lights_10km	Mean artificial light emissivity during nighttime in 10km buffer around the sampling coordinates, in microflicks	-0.31 to 50.49	0	0.30145052	(Elvidge et al. 2021)
nighttime_lights_20km	Mean artificial light emissivity during nighttime in 20km buffer around the sampling coordinates, in microflicks	-0.41 to 38.77	0	0.420156367	(Elvidge et al. 2021)
nighttime_lights_30km	Mean artificial light emissivity during nighttime in 30km buffer around the sampling coordinates, in microflicks	-0.39 to 23.99	0	0.527438576	(Elvidge et al. 2021)
human_population_10km	Human population in a 10km buffer around the sampling coordinates	0 to 3418112	0	8533.319525	(European Commission et al. 2019)
human_population_20km	Human population in a 20km buffer around the sampling coordinates	0 to 6413067	0	77634.95748	(European Commission et al. 2019)
human_population_30km	Human population in a 30km buffer around the sampling coordinates	335 to 7188317	0	265187.7767	(European Commission et al. 2019)

oil_palm_10km	Percentage of area covered by oil palm plantations in 10km buffer around the sampling coordinates	0 to 62.58	0	28.0820553674567	(Miettinen, Shi, and Liew 2016)
oil_palm_20km	Percentage of area covered by oil palm plantations in 20km buffer around the sampling coordinates	0 to 64	0	31.3603581198707	(Miettinen, Shi, and Liew 2016)
oil_palm_30km	Percentage of area covered by oil palm plantations in 30km buffer around the sampling coordinates	0 to 31.39	0	24.9347604936087	(Miettinen, Shi, and Liew 2016)
urban_areas_10km	Percentage of area covered human settlements in 10km buffer around the sampling coordinates	0 to 46.08	0	0	(FAO et al. 2014)
urban_areas_20km	Percentage of area covered human settlements in 20km buffer around the sampling coordinates	0 to 24.17	0	0.12303486	(FAO et al. 2014)
urban_areas_30km	Percentage of area covered human settlements in 30km buffer around the sampling coordinates	0 to 11.80	0	0.579732197	(FAO et al. 2014)
protected_areas_10km	Percentage of area protected in 10km buffer around the sampling coordinates	0 to 100	0	57.70151636	(UNEP-WCMC and IUCN 2021)
protected_areas_20km	Percentage of area protected in 20km buffer around the sampling coordinates	0 to 100	0	35.98726115	(UNEP-WCMC and IUCN 2021)
protected_areas_30km	Percentage of area protected in 30km buffer around the sampling coordinates	0 to 100	0	28.62700027	(UNEP-WCMC and IUCN 2021)

307

308

309 captures.csv

Field Name	Description	Range	NAs	Example	Source
survey_id	Unique ID which connects the records from the capture table with the metadata table.	NA	0	Lambir2017.ECL	compiler
records	Number of independent records	0 - 3644	0	3	original authors
Y_lat	Latitude in decimal degrees, using WGS84	-8.71 to 33.02	0	3.972066	original authors
X_long	Longitude in decimal degrees, using WGS84	80.25 to 140.09	0	98.08855	original authors
year_start	Year of the start of the trapping survey	1987 - 2020	0	2014	original authors
country	Country where sampling was performed	NA	0	Malaysia	original authors
domestic	States whether the species recorded is domestic or wild	“domestic” or “wild”	0	wild	IUCN Red List
uri	Link for the species webpage at the NCBI or GBIF	NA	0	https://www.ncbi.nlm.nih.gov/taxonomy/37029	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
species	Species name of the recorded animal (when identified to the species taxonomic level)	NA	549	bengalensis	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
genus	Genus name of the recorded animal (when identified at least to the genus taxonomic level)	NA	234	Prionailurus	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
family	Family name of the recorded animal (when identified at least to the family taxonomic level)	NA	102	Felidae	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
order	Order name of the recorded animal (when identified at least to the order taxonomic level)	NA	0	Carnivora	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)

class	Class name of the recorded animal (when identified at least to the class taxonomic level)	Aves, Mammalia or Reptilia	0	Mammalia	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
binomial_verified	Binomial name of the animal recorded, verified in the NCBI or GBIF database.	NA	0	Prionailurus bengalensis	original authors, verified by NCBI or GBIF
taxonomic_level	Taxonomic level in which the animal record was identified	species, genus, order or family	0	species	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)

NCBI - National Center for Biotechnology Information (www.ncbi.nlm.nih.gov)

GBIF - Global Biodiversity Information Facility (www.gbif.org)

species_traits.csv

Field Name	Description	Range	NAs	Example	Source
uri	Link for the species webpage at the NCBI or GBIF	NA	0	https://www.ncbi.nlm.nih.gov/taxonomy/9691	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
species	Species name of the recorded animal (when identified to the species taxonomic level)	NA	63	pardus	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
genus	Genus name of the recorded animal (when identified at least to the genus taxonomic level)	NA	33	Panthera	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
family	Family name of the recorded animal (when identified at least to the family taxonomic level)	NA	8	Felidae	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
order	Order name of the recorded animal (when identified at least to the order taxonomic level)	NA	0	Carnivora	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)

class	Class name of the recorded animal (when identified at least to the class taxonomic level)	Aves, Mammalia or Reptilia	0	Mammalia	(Schoch et al. 2020; GBIF 2022; Chamberlain et al. 2022)
binomial_verified	Binomial name of the animal recorded, verified in the NCBI or GBIF database.	NA	0	Panthera pardus	original authors, verified by NCBI or GBIF
taxonomic_level	Taxonomic level in which the animal record was identified	species, genus, order or family	0	species	original authors, verified by NCBI or GBIF
adult_mass_g	Average body mass of an adult individual, in grams	25.99 to 3220000	192	53075	(Soria et al. 2021)
dphy_invertebrate	Percentage of the diet composed by invertebrates	0 to 100	192	0	(Soria et al. 2021)
dphy_vertibrate	Percentage of the diet composed by vertebrates	0 to 100	192	100	(Soria et al. 2021)
dphy_plant	Percentage of the diet composed by plants	0 to 100	192	0	(Soria et al. 2021)
det_diet_breadth_n	Number of dietary categories with consumption above 20%, based on EltonTraits	1 to 4	192	1	(Soria et al. 2021)
trophic_level	Trophic level of the species recorded	1 for herbivores, 2 for omnivores and 3 for carnivores	192	3	(Soria et al. 2021)
activity_cycle	Dial activity cycle of the recorded species	1 for strictly nocturnal, 2 for cathemeral and crepuscular, 3 for strictly diurnal species	192	1	(Soria et al. 2021)
habitat_breadth_n	Number of habitats suitable for the species, based on IUCN	1-7	198	6	(Soria et al. 2021)

iucn2020_binomial	Binomial name of the recorded animal, based on the IUCN taxonomic database	Accipiter gentilis - Zoothera dauma	0	Panthera pardus	(IUCN 2022; Chamberlain et al. 2022)
IUCN	IUCN redlist status	CR - VU	84	VU	(IUCN 2022)

313 NCBI - National Center for Biotechnology Information (www.ncbi.nlm.nih.gov)

314 GBIF - Global Biodiversity Information Facility (www.gbif.org)

315 IUCN - The International Union for Conservation of Nature (www.iucn.org)

316

For Review Only

E. Data anomalies:

Not available data is indicated by “NA”.

Literature Citations

Ahumada, Jorge A., Eric Fegraus, Tanya Birch, Nicole Flores, Roland Kays, Timothy G. O'Brien, Jonathan Palmer, Stephanie Schuttler, Jennifer Y. Zhao, Walter Jetz, Margaret Kinnaird, Sayali Kulkarni, Arnaud Lyet, David Thau, Michelle Duong, Ruth Oliver, and Anthony Dancer. 2020. "Wildlife Insights: A Platform to Maximize the Potential of Camera Trap and Other Passive Sensor Wildlife Data for the Planet." *Environmental Conservation* 47 (1): 1-6. <https://doi.org/10.1017/S0376892919000298>. <https://www.cambridge.org/core/article/wildlife-insights-a-platform-to-maximize-the-potential-of-camera-trap-and-other-passive-sensor-wildlife-data-for-the-planet/98295387F86A977F2ECD96CCC5705CCC>.

Beaudrot, Lydia, Jorge A. Ahumada, Timothy O'Brien, Patricia Alvarez-Loayza, Kelly Boekee, Ahimsa Campos-Arceiz, David Eichberg, Santiago Espinosa, Eric Fegraus, Christine Fletcher, Krisna Gajapersad, Chris Hallam, Johanna Hurtado, Patrick A. Jansen, Amit Kumar, Eileen Larney, Marcela Guimarães Moreira Lima, Colin Mahony, Emanuel H. Martin, Alex McWilliam, Badru Mugerwa, Mireille Ndoundou-Hockemba, Jean Claude Razafimahaimodison, Hugo Romero-Saltos, Francesco Rovero, Julia Salvador, Fernanda Santos, Douglas Sheil, Wilson R. Spironello, Michael R. Willig, Nurul L. Winarni, Alex Zvoleff, and Sandy J. Andelman. 2016. "Standardized Assessment of Biodiversity Trends in Tropical Forest Protected Areas: The End Is Not in Sight." *PLOS Biology* 14 (1): e1002357. <https://doi.org/10.1371/journal.pbio.1002357>. <https://doi.org/10.1371/journal.pbio.1002357>.

Beyer, Hawthorne L., Oscar Venter, Hedley S. Grantham, and James E.M. Watson. 2020. "Substantial losses in ecoregion intactness highlight urgency of globally coordinated action." *Conservation Letters* 13 (2): e12692. <https://doi.org/https://doi.org/10.1111/conl.12692>. <https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/conl.12692>.

Ceballos, Gerardo, Paul R. Ehrlich, Anthony D. Barnosky, Andrés García, Robert M. Pringle, and Todd M. Palmer. 2015. "Accelerated modern human-induced species losses: Entering the sixth mass extinction." *Science Advances* 1 (5): e1400253. <https://doi.org/doi:10.1126/sciadv.1400253>. <https://www.science.org/doi/abs/10.1126/sciadv.1400253>.

taxize: Taxonomic Information from Around the Web 0.9.100. CRAN.

Dirzo, Rodolfo, Hillary S. Young, Mauro Galetti, Gerardo Ceballos, Nick J. B. Isaac, and Ben Collen. 2014. "Defaunation in the Anthropocene." *Science* 345 (6195): 401-406. <https://doi.org/doi:10.1126/science.1251817>. <https://www.science.org/doi/abs/10.1126/science.1251817>.

Elvidge, Christopher D., Mikhail Zhizhin, Tilottama Ghosh, Feng-Chi Hsu, and Jay Taneja. 2021. "Annual Time Series of Global VIIRS Nighttime Lights Derived from Monthly Averages: 2012 to 2019." *Remote Sensing* 13 (5): 922. <https://www.mdpi.com/2072-4292/13/5/922>.

European Commission, Joint Research Centre, D. Ehrlich, A. Florczyk, M. Pesaresi, L. Maffenini, M. Schiavina, L. Zanchetta, P. Politis, T. Kemper, F. Sabo, S. Freire, C. Corbane, and M. Melchiorri. 2019. *GHSL data package 2019 : public release GHS P2019*. Publications Office.

FAO, John Latham, Renato Cumani, Ilaria Rosati, and Mario Bloise. 2014. Global Land Cover SHARE (GLC-SHARE) database Beta-Release Version 1.0 - 2014. edited by FAO.

GBIF. 2022. The Global Biodiversity Information Facility. GBIF.

Grantham, H. S., A. Duncan, T. D. Evans, K. R. Jones, H. L. Beyer, R. Schuster, J. Walston, J. C. Ray, J. G. Robinson, M. Callow, T. Clements, H. M. Costa, A. DeGemmis, P. R. Elsen, J. Ervin, P. Franco, E. Goldman, S. Goetz, A. Hansen, E. Hofsvang, P. Jantz, S. Jupiter, A. Kang, P. Langhammer, W. F. Laurance, S. Lieberman, M. Linkie, Y. Malhi, S. Maxwell, M. Mendez, R. Mittermeier, N. J. Murray, H. Possingham, J. Radachowsky, S. Saatchi, C. Samper, J. Silverman, A. Shapiro, B. Strassburg, T. Stevens, E. Stokes, R. Taylor, T. Tear, R. Tizard, O. Venter, P. Visconti, S. Wang, and J. E. M. Watson. 2020. "Anthropogenic modification of forests means only 40% of remaining forests have high ecosystem integrity." *Nature Communications* 11 (1): 5978. <https://doi.org/10.1038/s41467-020-19493-3>. <https://doi.org/10.1038/s41467-020-19493-3>.

terra: Spatial Data Analysis. R package version 1.6-17.

IPCC. 2014. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland.

- IUCN. 2022. The IUCN Red List of Threatened Species. Version 2022-1.
- Jaxa. 2015. ALOS Global Digital Surface Model "ALOS World 3D-30m"(AW3D30).
- Lima, Fernando, Gabrielle Beca, Renata Muylaert, Clinton Jenkins, Miriam Perilli, Ana Paschoal, Rodrigo Massara, Adriano Paglia, Adriano Chiarello, Mauricio Graipel, Jorge Cherem, Andre Regolin, Luiz Oliveira-Santos, Carlos Brocardo, Agustin Paviolo, Mario Di Bitetti, Leandro Moraes, Fabiana Lopes Rocha, Roberto Fusco-Costa, and Mauro Galetti. 2017. "ATLANTIC-CAMTRAPS: a dataset of medium and large terrestrial mammal communities in the Atlantic Forest of South America." *Ecology* 98. <https://doi.org/10.1002/ecy.1998>.
- Miettinen, Jukka, Chenghua Shi, and Soo Chin Liew. 2016. "2015 Land cover map of Southeast Asia at 250 m spatial resolution." *Remote Sensing Letters* 7: 701-710. <https://doi.org/10.1080/2150704X.2016.1182659>.
- O'Connell, Allan F., James D. Nichols, and K. Ullas Karanth. 2010. *Camera Traps in Animal Ecology*. Springer Tokyo.
- Schoch, C. L., S. Ciufu, M. Domrachev, C. L. Hutton, S. Kannan, R. Khovanskaya, D. Leipe, R. McVeigh, K. O'Neill, B. Robbertse, S. Sharma, V. Soussov, J. P. Sullivan, L. Sun, S. Turner, and I. Karsch-Mizrachi. 2020. "NCBI Taxonomy: a comprehensive update on curation, resources and tools." *Database (Oxford)* 2020. <https://doi.org/10.1093/database/baaa062>.
- Sexton, Joseph O., Xiao-Peng Song, Min Feng, Praveen Noojipady, Anupam Anand, Chengquan Huang, Do-Hyung Kim, Kathrine M. Collins, Saurabh Channan, Charlene DiMiceli, and John R. Townshend. 2013. "Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error." *International Journal of Digital Earth* 6 (5): 427-448. <https://doi.org/10.1080/17538947.2013.786146>. <https://doi.org/10.1080/17538947.2013.786146>.
- Sollmann, Rahel. 2018. "A gentle introduction to camera-trap data analysis." *African Journal of Ecology* 56: 740-749. <https://doi.org/https://doi.org/10.1111/aje.12557>.
- Soria, C. D., M. Pacifici, M. Di Marco, S. M. Stephen, and C. Rondinini. 2021. "COMBINE: a coalesced mammal database of intrinsic and extrinsic traits." *Ecology* 102 (6): e03344. <https://doi.org/10.1002/ecy.3344>.
- UNEP-WCMC, and IUCN. 2021. Protected Planet: The World Database on Protected Areas (WDPA). Cambridge.
- Venter, O., E. W. Sanderson, A. Magrach, J. R. Allan, J. Beher, K. R. Jones, H. P. Possingham, W. F. Laurance, P. Wood, B. M. Fekete, M. A. Levy, and J. E. Watson. 2018. Last of the Wild Project, Version 3 (LWP-3): 2009 Human Footprint, 2018 Release. Palisades, New York: NASA Socioeconomic Data and Applications Center (SEDAC).




Figure 1. Map of the collated camera trap records.

249x249mm (300 x 300 DPI)




Figure 2. Species richness per survey.

249x249mm (300 x 300 DPI)




Figure 3. Temporal distribution of the deployment dates for surveys. Note that it often takes years for surveys to be published and that covid shut down most research in 2020, explaining the relatively low number of surveys included with cameras deployed since 2020.

349x199mm (300 x 300 DPI)





Figure 4. Independent detections per mammalian genus. A total of 104 genera of mammals were observed, from 13 orders.

449x299mm (300 x 300 DPI)






Figure 6. Independent detection per avian genus. A total of 72 genera of birds were observed from 14 orders. Note that many studies did not identify birds.

449x299mm (300 x 300 DPI)




Figure 7. Species, genus and order of the recorded birds. Note that birds were not exhaustively identified in all surveys.

357x357mm (300 x 300 DPI)