

# HeKA Staff Meeting

**Simon Bussy**

[simon.bussy@gmail.com](mailto:simon.bussy@gmail.com)

## **Lights: a generalized joint model for high-dimensional multivariate longitudinal data and censored durations**

S. Bussy <sup>(1,2)</sup>, V.T. Nguyen <sup>(1,2)</sup>, A. Barbieri <sup>(3)</sup>, S. Zohar <sup>(4)</sup>, A.S. Jannot <sup>(4,5)</sup>

<sup>(1)</sup>LOPF, Califrais' Machine Learning Lab, Paris, France

<sup>(2)</sup>LPSM, UMR 8001, CNRS, Sorbonne University, Paris, France

<sup>(3)</sup>INSERM, UMR 1219, Bordeaux Population Health Research Center, Univ. Bordeaux, France

<sup>(4)</sup>INSERM, UMRS 1138, Centre de Recherche des Cordeliers, Paris, France

<sup>(5)</sup>Biomedical Informatics and Public Health Department, EGP, APHP, Paris, France

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# I. Introduction

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- Deal with the problem of joint modeling of longitudinal data and censored durations

## Introduction

### Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- ▶ Deal with the problem of joint modeling of longitudinal data and censored durations
- ▶ Large number of both longitudinal and time-independent features are available

## Introduction

### Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- ▶ Deal with the problem of joint modeling of longitudinal data and censored durations
- ▶ Large number of both longitudinal and time-independent features are available
- ▶ Flexibility in modeling the dependency between the longitudinal features and the event time with appropriate penalties

## Introduction

### Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- ▶ Deal with the problem of joint modeling of longitudinal data and censored durations
- ▶ Large number of both longitudinal and time-independent features are available
- ▶ Flexibility in modeling the dependency between the longitudinal features and the event time with appropriate penalties
- ▶ Inference achieved using an efficient and novel Proximal Quasi-Newton Expectation Maximization algorithm

## Introduction

### Overview

### Use cases

### Framework

## Model

### Submodels

### Notations

### Likelihood

## Inference

### Penalization

### prox-QNEM

### Evaluation

## Application

### Competitors

### Simulations

### Results

## Conclusion

## References

- Predict the risk for an event of interest to occur quickly, taking into account simultaneously a huge number of longitudinal signals in a high-dimensional context

## Introduction

Overview

**Use cases**

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- ▶ Predict the risk for an event of interest to occur quickly, taking into account simultaneously a huge number of longitudinal signals in a high-dimensional context
- ▶ Provides powerful interpretability by automatically pinpointing significant time-dependent and time-independent features

## Introduction

Overview

**Use cases**

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References



- ▶ Predict the risk for an event of interest to occur quickly, taking into account simultaneously a huge number of longitudinal signals in a high-dimensional context
- ▶ Provides powerful interpretability by automatically pinpointing significant time-dependent and time-independent features

## Real-time decision support

- ▶ Medical context → event of interest: survival time, re-hospitalization, relapse or disease progression ; longitudinal data: biomarkers or vital parameters measurements

### Introduction

Overview

**Use cases**

Framework

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

- ▶ Predict the risk for an event of interest to occur quickly, taking into account simultaneously a huge number of longitudinal signals in a high-dimensional context
- ▶ Provides powerful interpretability by automatically pinpointing significant time-dependent and time-independent features

## Real-time decision support

- ▶ Medical context → event of interest: survival time, re-hospitalization, relapse or disease progression ; longitudinal data: biomarkers or vital parameters measurements
- ▶ Customer's satisfaction monitoring context → event of interest: time when a client churns ; longitudinal data: the client's activity recorded from account opening throughout the duration of the business relationship

### Introduction

Overview

**Use cases**

Framework

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

## ► Survival analysis

$$T = T^* \wedge C \quad \text{and} \quad \Delta = \mathbb{1}_{\{T^* \leq C\}}$$

### Introduction

Overview

Use cases

**Framework**

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

# High-dimensional framework

- Survival analysis

$$T = T^* \wedge C \quad \text{and} \quad \Delta = \mathbb{1}_{\{T^* \leq C\}}$$

- Time-independent features  $X \in \mathbb{R}^p$  with  $p \gg n$

## Introduction

Overview

Use cases

**Framework**

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

# High-dimensional framework

- Survival analysis

$$T = T^* \wedge C \quad \text{and} \quad \Delta = \mathbb{1}_{\{T^* \leq C\}}$$

- Time-independent features  $X \in \mathbb{R}^p$  with  $p \gg n$
- $L$  longitudinal outcomes such that  $L \gg n$  and

$$Y(t) = (Y^1(t), \dots, Y^L(t))^T \in \mathbb{R}^L$$

## Introduction

Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

# High-dimensional framework

- Survival analysis

$$T = T^* \wedge C \quad \text{and} \quad \Delta = \mathbb{1}_{\{T^* \leq C\}}$$

- Time-independent features  $X \in \mathbb{R}^p$  with  $p \gg n$

- $L$  longitudinal outcomes such that  $L \gg n$  and

$$Y(t) = (Y^1(t), \dots, Y^L(t))^T \in \mathbb{R}^L$$

- Heterogeneity of the population: latent subgroups

$$G \in \{0, \dots, K-1\}$$

## Introduction

Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- Survival analysis

$$T = T^* \wedge C \quad \text{and} \quad \Delta = \mathbb{1}_{\{T^* \leq C\}}$$

- Time-independent features  $X \in \mathbb{R}^p$  with  $p \gg n$

- $L$  longitudinal outcomes such that  $L \gg n$  and

$$Y(t) = (Y^1(t), \dots, Y^L(t))^T \in \mathbb{R}^L$$

- Heterogeneity of the population: latent subgroups

$$G \in \{0, \dots, K-1\}$$

- Softmax link function for the latent class membership probability given time-independent features

$$\pi_{\xi_k}(x) = \mathbb{P}[G = k | X = x] = \frac{e^{x^T \xi_k}}{\sum_{k=0}^{K-1} e^{x^T \xi_k}}$$

## Introduction

Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# II. Model



## Group-specific marker trajectories

- $h_l(\mathbb{E}[Y^l(t)|b^l, G = k]) = m_k^l(t) = u^l(t)^\top \beta_k^l + v^l(t)^\top b^l$   
with fixed effect parameters  $\beta_k^l \in \mathbb{R}^{q_l}$  and subject-and-longitudinal outcome specific random effects  $b^l \in \mathbb{R}^{r_l} \sim \mathcal{N}(0, D_{ll})$

### Introduction

Overview

Use cases

Framework

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

# Submodels

## Group-specific marker trajectories

- ▶  $h_l(\mathbb{E}[Y^l(t)|b^l, G = k]) = m_k^l(t) = u^l(t)^\top \beta_k^l + v^l(t)^\top b^l$   
with fixed effect parameters  $\beta_k^l \in \mathbb{R}^{q_l}$  and subject-and-longitudinal outcome specific random effects  $b^l \in \mathbb{R}^{r_l} \sim \mathcal{N}(0, D_{ll})$
- ▶  $\text{Cov}[b^l, b^{l'}] = D_{ll'}$  and

$$D = \begin{bmatrix} D_{11} & \cdots & D_{1L} \\ \vdots & \ddots & \vdots \\ D_{1L}^\top & \cdots & D_{LL} \end{bmatrix}$$

the global variance-covariance matrix

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# Submodels

## Group-specific marker trajectories

- ▶  $h_l(\mathbb{E}[Y^l(t)|b^l, G = k]) = m_k^l(t) = u^l(t)^\top \beta_k^l + v^l(t)^\top b^l$   
with fixed effect parameters  $\beta_k^l \in \mathbb{R}^{q_l}$  and subject-and-longitudinal outcome specific random effects  $b^l \in \mathbb{R}^{r_l} \sim \mathcal{N}(0, D_{ll})$
- ▶  $\text{Cov}[b^l, b^{l'}] = D_{ll'}$

## Group-specific risk of event

$$\lambda(t|\mathcal{Y}(t), G = k) = \lambda_0(t) \exp \left\{ \underbrace{\sum_{l=1}^L \sum_{a=1}^{\mathcal{A}} \gamma_{k,a}^l \Psi_a^l(t)}_{\gamma_k^\top \Psi(t)} \right\}$$

with  $\mathcal{Y}(t) = \{Y(u), 0 \leq u < t\}$  and for each  $l$ -th longitudinal outcome, we consider  $\mathcal{A} \in \mathbb{N}_+$  known functionals  $\Psi_a^l$  extracted from  $\mathcal{Y}^l(t)$  through a given representation mapping

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

## Group-specific marker trajectories

- ▶  $h_l(\mathbb{E}[Y^l(t)|b^l, G = k]) = m_k^l(t) = u^l(t)^\top \beta_k^l + v^l(t)^\top b^l$   
with fixed effect parameters  $\beta_k^l \in \mathbb{R}^{q_l}$  and subject-and-longitudinal outcome specific random effects  $b^l \in \mathbb{R}^{r_l} \sim \mathcal{N}(0, D_{ll})$
- ▶  $\text{Cov}[b^l, b^{l'}] = D_{ll'}$

## Group-specific risk of event

- ▶  $\lambda(t|\mathcal{Y}(t), G = k) = \lambda_0(t) \exp \left\{ \underbrace{\sum_{l=1}^L \sum_{a=1}^{\mathcal{A}} \gamma_{k,a}^l \Psi_a^l(t)}_{\gamma_k^\top \Psi(t)} \right\}$

with  $\mathcal{Y}(t) = \{Y(u), 0 \leq u < t\}$  and for each  $l$ -th longitudinal outcome, we consider  $\mathcal{A} \in \mathbb{N}_+$  known functionals  $\Psi_a^l$  extracted from  $\mathcal{Y}^l(t)$  through a given representation mapping

- ▶  $\Psi(t) \in \mathbb{R}^{L\mathcal{A}}$  : high-dim representation features vector, highly flexible and independent of any modeling assumption !

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

- ▶ JLCMs [8] : homogeneous latent subgroups sharing the same marker trajectories and the same prognostic

## Introduction

Overview  
Use cases  
Framework

## Model

**Submodels**  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# Generalization of SREMs and JLCMs

- ▶ JLCMs [8] : homogeneous latent subgroups sharing the same marker trajectories and the same prognostic
- ▶ SREMs [9] : characteristics of the longitudinal processes (e.g. functions of the random effects) are included as features in the survival model

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# Generalization of SREMs and JLCMs

- ▶ JLCMs [8] : homogeneous latent subgroups sharing the same marker trajectories and the same prognostic
- ▶ SREMs [9] : characteristics of the longitudinal processes (e.g. functions of the random effects) are included as features in the survival model
- ▶ Usual functionals would be

Description	$\tilde{\Psi}_a^l(t, \beta_k^l, b^l)$	Reference
Linear predictor	$m_k^l(t)$	Chi and Ibrahim [2]
Random effects	$b^l$	Hatfield et al. [3]
Time-dependent slope	$\frac{d}{dt} m_k^l(t)$	Rizopoulos and Ghosh [6]
Cumulative effect	$\int_0^t m_k^l(s) ds$	Andrinopoulou et al. [1]

## Introduction

Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

Simulations

Results

## Conclusion

## References

- ▶ JLCMs [8] : homogeneous latent subgroups sharing the same marker trajectories and the same prognostic
- ▶ SREMs [9] : characteristics of the longitudinal processes (e.g. functions of the random effects) are included as features in the survival model
- ▶ Usual functionals would be


Description	$\tilde{\Psi}_a^l(t, \beta_k^l, b^l)$	Reference
Linear predictor	$m_k^l(t)$	Chi and Ibrahim [2]
Random effects	$b^l$	Hatfield et al. [3]
Time-dependent slope	$\frac{d}{dt} m_k^l(t)$	Rizopoulos and Ghosh [6]
Cumulative effect	$\int_0^t m_k^l(s) ds$	Andrinopoulou et al. [1]

- ▶ Depend on  $\beta_k$  (then on  $k$ ) which leads to complicated updates, and on  $b$  which leads to untrackable integrals that requires approximation methods : computationally intensive and not scalable



# Graphical model

Lights  
7/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# Notations

- $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_i^l}^l)^\top \in \mathbb{R}^{n_i^l}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
**Notations**  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# Notations

- ▶  $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_l}^l)^\top \in \mathbb{R}^{n_l^l}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$
- ▶  $y_i = (y_i^{1^\top} \cdots y_i^{L^\top})^\top \in \mathbb{R}^{n_i}$  with  $n_i = \sum_{l=1}^L n_l^l$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
**Notations**  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# Notations

- ▶  $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_l}^l)^\top \in \mathbb{R}^{n_l'}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$
- ▶  $y_i = (y_i^{1^\top} \cdots y_i^{L^\top})^\top \in \mathbb{R}^{n_i}$  with  $n_i = \sum_{l=1}^L n_l'$
- ▶  $b_i = (b_i^{1^\top} \cdots b_i^{L^\top})^\top \in \mathbb{R}^r$  with  $r = \sum_{l=1}^L r_l$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
**Notations**  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶  $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_l}^l)^\top \in \mathbb{R}^{n_l'}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$
- ▶  $y_i = (y_i^{1\top} \cdots y_i^{L\top})^\top \in \mathbb{R}^{n_i}$  with  $n_i = \sum_{l=1}^L n_l'$
- ▶  $b_i = (b_i^{1\top} \cdots b_i^{L\top})^\top \in \mathbb{R}^r$  with  $r = \sum_{l=1}^L r_l$
- ▶ Design matrices

$$U_i = \begin{bmatrix} U_{i1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & U_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times q} \text{ and } V_i = \begin{bmatrix} V_{i1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & V_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times r}$$

with  $q = \sum_{l=1}^L q_l$  and where for all  $l = 1, \dots, L$ , one writes

$$\begin{cases} U_{il} = (u_i^l(t_{i1}^l)^\top \cdots u_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l' \times q_l}, \\ V_{il} = (v_i^l(t_{i1}^l)^\top \cdots v_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l' \times r_l}. \end{cases}$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
**Notations**  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶  $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_l}^l)^\top \in \mathbb{R}^{n_l'}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$
- ▶  $y_i = (y_i^{1\top} \dots y_i^{L\top})^\top \in \mathbb{R}^{n_i}$  with  $n_i = \sum_{l=1}^L n_l'$
- ▶  $b_i = (b_i^{1\top} \dots b_i^{L\top})^\top \in \mathbb{R}^r$  with  $r = \sum_{l=1}^L r_l$
- ▶ Design matrices

$$U_i = \begin{bmatrix} U_{i1} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & U_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times q} \text{ and } V_i = \begin{bmatrix} V_{i1} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & V_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times r}$$

with  $q = \sum_{l=1}^L q_l$  and where for all  $l = 1, \dots, L$ , one writes

$$\begin{cases} U_{il} = (u_i^l(t_{i1}^l)^\top \dots u_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l' \times q_l}, \\ V_{il} = (v_i^l(t_{i1}^l)^\top \dots v_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l' \times r_l}. \end{cases}$$

- ▶  $\beta_k = (\beta_k^{1\top} \dots \beta_k^{L\top})^\top \in \mathbb{R}^q$

- ▶  $\mathcal{D}_n = \{(x_1, y_1^1, \dots, y_1^L, t_1, \delta_1), \dots, (x_n, y_n^1, \dots, y_n^L, t_n, \delta_n)\}$  with  $y_i^l = (y_{i1}^l, \dots, y_{in_l}^l)^\top \in \mathbb{R}^{n_l^l}$  and  $y_{ij}^l = Y_i^l(t_{ij}^l)$
- ▶  $y_i = (y_i^{1^\top} \dots y_i^{L^\top})^\top \in \mathbb{R}^{n_i}$  with  $n_i = \sum_{l=1}^L n_l^l$
- ▶  $b_i = (b_i^{1^\top} \dots b_i^{r^\top})^\top \in \mathbb{R}^r$  with  $r = \sum_{l=1}^L r_l$
- ▶ Design matrices

$$U_i = \begin{bmatrix} U_{i1} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & U_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times q} \text{ and } V_i = \begin{bmatrix} V_{i1} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & V_{iL} \end{bmatrix} \in \mathbb{R}^{n_i \times r}$$

with  $q = \sum_{l=1}^L q_l$  and where for all  $l = 1, \dots, L$ , one writes

$$\begin{cases} U_{il} = (u_i^l(t_{i1}^l)^\top \dots u_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l^l \times q_l}, \\ V_{il} = (v_i^l(t_{i1}^l)^\top \dots v_i^l(t_{in_l}^l)^\top)^\top \in \mathbb{R}^{n_l^l \times r_l}. \end{cases}$$

- ▶  $\beta_k = (\beta_k^{1^\top} \dots \beta_k^{L^\top})^\top \in \mathbb{R}^q$
- ▶  $M_{ik} = U_i \beta_k + V_i b_i \in \mathbb{R}^{n_i}$

$$\blacktriangleright \theta = (\xi_0^\top \cdots \xi_{K-1}^\top, \beta_0^\top \cdots \beta_{K-1}^\top, \phi^\top, \text{vech}(D), \lambda_0^\top, \gamma_0^\top \cdots \gamma_{K-1}^\top) \in \mathbb{R}^\vartheta$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
**Likelihood**

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References



- ▶  $\theta = (\xi_0^\top \cdots \xi_{K-1}^\top, \beta_0^\top \cdots \beta_{K-1}^\top, \phi^\top, \text{vech}(D), \lambda_0^\top, \gamma_0^\top \cdots \gamma_{K-1}^\top) \in \mathbb{R}^\vartheta$
- ▶  $f(y_i | b_i, G_i = k) = \exp \left\{ (y_i \odot \Phi_i)^\top M_{ik} - c_\phi(M_{ik}) + d_\phi(y_i) \right\}$  with  $\Phi_i = (\phi_1^{-1} \mathbf{1}_{n_i^1}^\top \cdots \phi_L^{-1} \mathbf{1}_{n_i^L}^\top)^\top \in \mathbb{R}^{n_i}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
**Likelihood**

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶  $\theta = (\xi_0^\top \cdots \xi_{K-1}^\top, \beta_0^\top \cdots \beta_{K-1}^\top, \phi^\top, \text{vech}(D), \lambda_0^\top, \gamma_0^\top \cdots \gamma_{K-1}^\top) \in \mathbb{R}^\vartheta$
- ▶  $f(y_i | b_i, G_i = k) = \exp \left\{ (y_i \odot \Phi_i)^\top M_{ik} - c_\phi(M_{ik}) + d_\phi(y_i) \right\}$  with  $\Phi_i = (\phi_1^{-1} \mathbf{1}_{n_i^1}^\top \cdots \phi_L^{-1} \mathbf{1}_{n_i^L}^\top)^\top \in \mathbb{R}^{n_i}$
- ▶ Survival part:

$$f(t_i, \delta_i | G_i = k; \theta) = [\lambda(t_i | \mathcal{Y}(t_i), G_i = k)]^{\delta_i} \\ \times \exp \left\{ - \int_0^{t_i} \lambda(s | \mathcal{Y}(s), G_i = k) ds \right\}$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
**Likelihood**

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶  $\theta = (\xi_0^\top \cdots \xi_{K-1}^\top, \beta_0^\top \cdots \beta_{K-1}^\top, \phi^\top, \text{vech}(D), \lambda_0^\top, \gamma_0^\top \cdots \gamma_{K-1}^\top) \in \mathbb{R}^\vartheta$
- ▶  $f(y_i|b_i, G_i = k) = \exp \left\{ (y_i \odot \Phi_i)^\top M_{ik} - c_\phi(M_{ik}) + d_\phi(y_i) \right\}$  with  $\Phi_i = (\phi_1^{-1} \mathbf{1}_{n_i^1}^\top \cdots \phi_L^{-1} \mathbf{1}_{n_i^L}^\top)^\top \in \mathbb{R}^{n_i}$
- ▶ Survival part:

$$f(t_i, \delta_i | G_i = k; \theta) = [\lambda(t_i | \mathcal{Y}(t_i), G_i = k)]^{\delta_i} \times \exp \left\{ - \int_0^{t_i} \lambda(s | \mathcal{Y}(s), G_i = k) ds \right\}$$

- ▶ Then, the log-likelihood (JLCMs type) writes

$$\ell_n(\theta) = n^{-1} \sum_{i=1}^n \log \sum_{k=0}^{K-1} \pi_{\xi_k}(x_i) f_\theta(t_i, \delta_i | G_i = k) f_\theta(y_i | G_i = k) db_i$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
**Likelihood**

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# III. Inference

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

## ► Penalized objective

$$\ell_n^{\text{pen}}(\theta) = -\ell_n(\theta) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en},\eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sg}l_1,\tilde{\eta}}$$

with the elasticnet penalty

$$\|z\|_{\text{en},\eta} = (1 - \eta)\|z\|_1 + \frac{\eta}{2}\|z\|_2^2$$

and the sparse group lasso penalty

$$\|z\|_{\text{sg}l_1,\tilde{\eta}} = (1 - \tilde{\eta})\|z\|_1 + \tilde{\eta} \sum_{l=1}^L \|z^l\|_2$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

**Penalization**  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

## ► Penalized objective

$$\ell_n^{\text{pen}}(\theta) = -\ell_n(\theta) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en},\eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sg}l_1,\tilde{\eta}}$$

with the elasticnet penalty

$$\|z\|_{\text{en},\eta} = (1 - \eta)\|z\|_1 + \frac{\eta}{2}\|z\|_2^2$$

and the sparse group lasso penalty

$$\|z\|_{\text{sg}l_1,\tilde{\eta}} = (1 - \tilde{\eta})\|z\|_1 + \tilde{\eta} \sum_{l=1}^L \|z^l\|_2$$

## ► Resulting optimization problem

$$\hat{\theta} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \ell_n^{\text{pen}}(\theta)$$

### Introduction

Overview

Use cases

Framework

### Model

Submodels

Notations

Likelihood

### Inference

**Penalization**

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

# prox-QNEM algorithm (1/2)

Lights  
11/20

$$\blacktriangleright \ell_n^{\text{comp}}(\theta) = \ell_n^{\text{comp}}(\theta; \mathcal{D}_n, \mathbf{b}, \mathbf{G})$$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# prox-QNEM algorithm (1/2)

Lights  
11/20

$$\blacktriangleright \ell_n^{\text{comp}}(\theta) = \ell_n^{\text{comp}}(\theta; \mathcal{D}_n, \mathbf{b}, \mathbf{G})$$

## Monte Carlo E-step

$$\blacktriangleright Q_n(\theta, \theta^{(w)}) = \mathbb{E}_{\theta^{(w)}}[\ell_n^{\text{comp}}(\theta) | \mathcal{D}_n]$$

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
**prox-QNEM**  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References



- ▶  $\ell_n^{\text{comp}}(\theta) = \ell_n^{\text{comp}}(\theta; \mathcal{D}_n, \mathbf{b}, \mathbf{G})$

## Monte Carlo E-step

- ▶  $\mathcal{Q}_n(\theta, \theta^{(w)}) = \mathbb{E}_{\theta^{(w)}}[\ell_n^{\text{comp}}(\theta) | \mathcal{D}_n]$
- ▶ Requires to compute expectations of the form

$$\mathbb{E}_{\theta^{(w)}}[g(\mathbf{b}_i, \mathbf{G}_i) | t_i, \delta_i, y_i] = \sum_{k=0}^{K-1} \pi_{ik}^{\theta^{(w)}} \int_{\mathbb{R}^r} g(\mathbf{b}_i, \mathbf{G}_i) f_{\theta^{(w)}}(\mathbf{b}_i | t_i, \delta_i, y_i) d\mathbf{b}_i$$

for different functions  $g$ , where we denote

$$\pi_{ik}^{\theta^{(w)}} = \mathbb{P}_{\theta^{(w)}}[G_i = k | t_i, \delta_i, y_i]$$

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
**prox-QNEM**  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

$$\blacktriangleright \theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$$

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
**prox-QNEM**  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}}[b_i b_i^\top | t_i, \delta_i, y_i]$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}}[b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

## prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}}[b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$
- ▶  $\xi_k^{(w+1)} \in \operatorname{argmin}_{\xi_k \in \mathbb{R}^p} P_{n,k}^{(w)}(\xi_k) + \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}} [b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$
- ▶  $\xi_k^{(w+1)} \in \operatorname{argmin}_{\xi_k \in \mathbb{R}^p} P_{n,k}^{(w)}(\xi_k) + \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta}$
- ▶ L-BFGS-B to solve the problem

$$\text{minimize} \quad P_{n,k}^{(w)}(\xi_k^+ - \xi_k^-) + \zeta_{1,k} \left( (1 - \eta) \sum_{j=1}^p (\xi_{k,j}^+ + \xi_{k,j}^-) + \frac{\eta}{2} \|\xi_k^+ - \xi_k^-\|_2^2 \right)$$

$$\text{subject to} \quad \xi_{k,j}^+ \geq 0 \text{ and } \xi_{k,j}^- \geq 0 \text{ for } j = 1, \dots, p$$

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
**prox-QNEM**  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}} [b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$
- ▶  $\xi_k^{(w+1)} \in \operatorname{argmin}_{\xi_k \in \mathbb{R}^p} P_{n,k}^{(w)}(\xi_k) + \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta}$
- ▶ L-BFGS-B to solve the problem

$$\text{minimize} \quad P_{n,k}^{(w)}(\xi_k^+ - \xi_k^-) + \zeta_{1,k} \left( (1 - \eta) \sum_{j=1}^p (\xi_{k,j}^+ + \xi_{k,j}^-) + \frac{\eta}{2} \|\xi_k^+ - \xi_k^-\|_2^2 \right)$$

$$\text{subject to} \quad \xi_{k,j}^+ \geq 0 \text{ and } \xi_{k,j}^- \geq 0 \text{ for } j = 1, \dots, p$$

- ▶ Closed-form update for  $\beta_k^{(w+1)}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}} [b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$
- ▶  $\xi_k^{(w+1)} \in \operatorname{argmin}_{\xi_k \in \mathbb{R}^p} P_{n,k}^{(w)}(\xi_k) + \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta}$
- ▶ L-BFGS-B to solve the problem

$$\text{minimize} \quad P_{n,k}^{(w)}(\xi_k^+ - \xi_k^-) + \zeta_{1,k} \left( (1 - \eta) \sum_{j=1}^p (\xi_{k,j}^+ + \xi_{k,j}^-) + \frac{\eta}{2} \|\xi_k^+ - \xi_k^-\|_2^2 \right)$$

$$\text{subject to} \quad \xi_{k,j}^+ \geq 0 \text{ and } \xi_{k,j}^- \geq 0 \text{ for } j = 1, \dots, p$$

- ▶ Closed-form update for  $\beta_k^{(w+1)}$
- ▶ Proximal gradient (ISTA) for the  $\gamma_k^{(w+1)}$  update, based on Lemma 1 that states  $\operatorname{prox}_{\text{sgl}_1, \tilde{\eta}, \zeta} = \operatorname{prox}_{\zeta \tilde{\eta} \sum_{l=1}^L \|\cdot\|_{2,l}} \circ \operatorname{prox}_{\zeta(1-\tilde{\eta})\|\cdot\|_1}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References



# prox-QNEM algorithm (2/2)

## Proximal Quasi-Newton M-step

- ▶  $\theta^{(w+1)} \in \operatorname{argmin}_{\theta \in \mathbb{R}^{\vartheta}} \mathcal{Q}_n(\theta, \theta^{(w)}) + \sum_{k=0}^{K-1} \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta} + \zeta_{2,k} \|\gamma_k\|_{\text{sgl}_1, \tilde{\eta}}$
- ▶  $D^{(w+1)} = n^{-1} \sum_{i=1}^n \hat{\mathbb{E}}_{\theta^{(w)}} [b_i b_i^\top | t_i, \delta_i, y_i]$
- ▶  $P_{n,k}^{(w)}(\xi_k) = -n^{-1} \sum_{i=1}^n \hat{\pi}_{ik}^{\theta^{(w)}} \log \pi_{\xi_k}(x_i)$
- ▶  $\xi_k^{(w+1)} \in \operatorname{argmin}_{\xi_k \in \mathbb{R}^p} P_{n,k}^{(w)}(\xi_k) + \zeta_{1,k} \|\xi_k\|_{\text{en}, \eta}$
- ▶ L-BFGS-B to solve the problem

$$\text{minimize} \quad P_{n,k}^{(w)}(\xi_k^+ - \xi_k^-) + \zeta_{1,k} \left( (1 - \eta) \sum_{j=1}^p (\xi_{k,j}^+ + \xi_{k,j}^-) + \frac{\eta}{2} \|\xi_k^+ - \xi_k^-\|_2^2 \right)$$

$$\text{subject to} \quad \xi_{k,j}^+ \geq 0 \text{ and } \xi_{k,j}^- \geq 0 \text{ for } j = 1, \dots, p$$

- ▶ Closed-form update for  $\beta_k^{(w+1)}$
- ▶ Proximal gradient (ISTA) for the  $\gamma_k^{(w+1)}$  update, based on Lemma 1 that states  $\operatorname{prox}_{\text{sgl}_1, \tilde{\eta}, \zeta} = \operatorname{prox}_{\zeta \tilde{\eta} \sum_{l=1}^L \|\cdot\|_{2,l}} \circ \operatorname{prox}_{\zeta(1-\tilde{\eta})\|\cdot\|_1}$
- ▶  $\lambda_0^{(w+1)}(t) = \frac{\sum_{i=1}^n \delta_i \mathbb{1}_{\{t=t_i\}}}{\sum_{i=1}^n \sum_{k=0}^{K-1} \hat{\pi}_{ik}^{\theta^{(w)}} \exp \left\{ \gamma_k^{(w+1)\top} \Psi_i(t_i) \right\} \mathbb{1}_{\{t_i \geq t\}}}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
**prox-QNEM**  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM

### Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM

### Evaluation

### Application

Competitors  
Simulations  
Results


### Conclusion

### References

# Performance evaluation

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data
- ▶ On test data : sample  $t_i^{max} \sim T_i(1 - \text{Beta}(2, 5))$  the time for subject  $i$  when one wants to perform the risk prediction – the “present” time



### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM

### Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data
- ▶ On test data : sample  $t_i^{max} \sim T_i(1 - \text{Beta}(2, 5))$  the time for subject  $i$  when one wants to perform the risk prediction – the “present” time
- ▶ Truncate  $y_i$  up to  $t_i^{max}$  to get  $\tilde{y}_i$

### Introduction

Overview

Use cases

Framework

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data
- ▶ On test data : sample  $t_i^{max} \sim T_i(1 - \text{Beta}(2, 5))$  the time for subject  $i$  when one wants to perform the risk prediction – the “present” time
- ▶ Truncate  $y_i$  up to  $t_i^{max}$  to get  $\tilde{y}_i$

## Predictive marker

$$\text{▶ } \mathbb{P}_{\hat{\theta}}[G_i = k | T_i > t_i^{max}, \tilde{y}_i] = \frac{\pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}{\sum_{k=0}^{K-1} \pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}$$

### Introduction

Overview

Use cases

Framework

### Model

Submodels

Notations

Likelihood

### Inference

Penalization

prox-QNEM

Evaluation

### Application

Competitors

Simulations

Results

### Conclusion

### References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data
- ▶ On test data : sample  $t_i^{max} \sim T_i(1 - \text{Beta}(2, 5))$  the time for subject  $i$  when one wants to perform the risk prediction – the “present” time
- ▶ Truncate  $y_i$  up to  $t_i^{max}$  to get  $\tilde{y}_i$

## Predictive marker

- ▶ 
$$\mathbb{P}_{\hat{\theta}}[G_i = k | T_i > t_i^{max}, \tilde{y}_i] = \frac{\pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}{\sum_{k=0}^{K-1} \pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}$$
- ▶ In practice  $K = 2$

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

## Real-time prediction paradigm

- ▶ Practitioners need predictive prognostic tools to be used in real-time
- ▶ Obtain  $\hat{\theta}$  on training data
- ▶ On test data : sample  $t_i^{max} \sim T_i(1 - \text{Beta}(2, 5))$  the time for subject  $i$  when one wants to perform the risk prediction – the “present” time
- ▶ Truncate  $y_i$  up to  $t_i^{max}$  to get  $\tilde{y}_i$

## Predictive marker

- ▶ 
$$\mathbb{P}_{\hat{\theta}}[G_i = k | T_i > t_i^{max}, \tilde{y}_i] = \frac{\pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}{\sum_{k=0}^{K-1} \pi_{\hat{\xi}_k}(x_i) f_{\hat{\theta}}(t_i^{max}, \delta_i = 0, \tilde{y}_i | G_i = k)}$$
- ▶ In practice  $K = 2$
- ▶ C-index metric  $\mathcal{C}_\tau = \mathbb{P}[\hat{\mathcal{R}}_i > \hat{\mathcal{R}}_j | T_i < T_j, T_i < \tau]$



- ▶ Landmark Cox model : survival R package [10]

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

**Competitors**  
Simulations  
Results

## Conclusion

## References

- ▶ Landmark Cox model : survival R package [10]
- ▶ The time-dependent Cox model [7] : survival R package

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

**Competitors**  
Simulations  
Results

## Conclusion

## References

- ▶ Landmark Cox model : survival R package [10]
- ▶ The time-dependent Cox model [7] : survival R package
- ▶ Multivariate joint latent class model : R package lcmm [4]

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶ Landmark Cox model : survival R package [10]
- ▶ The time-dependent Cox model [7] : survival R package
- ▶ Multivariate joint latent class model : R package lcmm [4]
- ▶ Multivariate shared random effect model : JMbayes package [5]

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

### **Simulations**

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

### **Simulations**

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

### **Simulations**

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap}$  for  $j = 1, \dots, s$

## Introduction

Overview

Use cases

Framework

## Model

Submodels

Notations

Likelihood

## Inference

Penalization

prox-QNEM

Evaluation

## Application

Competitors

**Simulations**

Results

## Conclusion

## References



# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

## Simulations

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

## Simulations

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_s}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

## Simulations

Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_s}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
**Simulations**  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_s}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
**Simulations**  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_s}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$
- ▶  $(\tilde{\Psi}_{k,a}(t, \beta_k, b_i^l))_{a \in \{1,2\}} = (\beta_{k,1}^l + \beta_{k,2}^l t + b_{i,1}^l + b_{i,2}^l t, b_i^{l\top})$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
**Simulations**  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$
- ▶  $(\tilde{\Psi}_{k,a}(t, \beta_k, b_i^l))_{a \in \{1,2\}} = (\beta_{k,1}^l + \beta_{k,2}^l t + b_{i,1}^l + b_{i,2}^l t, b_i^{l\top})$
- ▶  $\tilde{\gamma}_{k,a}^l = \varsigma_2 \mathbb{1}_{\{l \in \mathcal{S}_k, a \in \{1,2\}\}}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$
- ▶  $(\tilde{\Psi}_{k,a}(t, \beta_k, b_i^l))_{a \in \{1,2\}} = (\beta_{k,1}^l + \beta_{k,2}^l t + b_{i,1}^l + b_{i,2}^l t, b_i^{l\top})$
- ▶  $\tilde{\gamma}_{k,a}^l = \varsigma_2 \mathbb{1}_{\{l \in \mathcal{S}_k, a \in \{1,2\}\}}$
- ▶  $\lambda_i(t|G_i = k) = \lambda_0(t) \exp \{ \iota_{i,k,1} + \iota_{i,k,2} t \}$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References



# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{ \lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$
- ▶  $(\tilde{\Psi}_{k,a}(t, \beta_k, b_i^l))_{a \in \{1,2\}} = (\beta_{k,1}^l + \beta_{k,2}^l t + b_{i,1}^l + b_{i,2}^l t, b_i^{l\top})$
- ▶  $\tilde{\gamma}_{k,a}^l = \varsigma_2 \mathbb{1}_{\{l \in \mathcal{S}_k, a \in \{1,2\}\}}$
- ▶  $\lambda_i(t|G_i = k) = \lambda_0(t) \exp \{ \iota_{i,k,1} + \iota_{i,k,2} t \}$
- ▶ Gompertz baseline  $\lambda_0(t) = \kappa_1 \kappa_2 \exp(\kappa_2 t)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

# High-dimensional simulation study (1/2)

- ▶ High-risk subjects proportion  $\pi_1 \in [0, 1]$
- ▶  $\mathcal{H} = \{\lfloor \pi_1 n \rfloor \text{ random samples without replacement} \}$
- ▶  $[x_{ij}] \in \mathbb{R}^{n \times p} \sim \mathcal{N}(0, \Sigma_1(\rho_1))$
- ▶  $x_{ij} \leftarrow x_{ij} + (-1)^{\mathbb{1}_{\{i \notin \mathcal{H}\}}} \text{gap for } j = 1, \dots, s$
- ▶  $\xi = (\underbrace{\varsigma_1, \dots, \varsigma_1}_s, 0, \dots, 0) \in \mathbb{R}^p$
- ▶  $G_i \sim \mathcal{B}(\pi_\xi(x_i))$
- ▶  $\mathcal{S}_k = \left\{ k \lfloor \frac{Lr_s}{K} \rfloor + 1, \dots, (k+1) \lfloor \frac{Lr_s}{K} \rfloor \right\}$
- ▶  $\beta_k^l \sim \mathbb{1}_{\{l \in \mathcal{S}_k\}} \mathcal{N}\left(\mu_k, \begin{bmatrix} \rho_3 & 0 \\ 0 & \rho_3 \end{bmatrix}\right)$
- ▶  $Y_i^l(t) = \epsilon_i^l(t) + \mathbb{1}_{\{l \in \mathcal{S}_k\}} \sum_{k=0}^{K-1} \mathbb{1}_{\{G_i=k\}} ((1, t)^\top \beta_k^l + (1, t)^\top b_i^l)$
- ▶  $(\tilde{\Psi}_{k,a}(t, \beta_k, b_i^l))_{a \in \{1,2\}} = (\beta_{k,1}^l + \beta_{k,2}^l t + b_{i,1}^l + b_{i,2}^l t, b_i^{l\top})$
- ▶  $\tilde{\gamma}_{k,a}^l = \varsigma_2 \mathbb{1}_{\{l \in \mathcal{S}_k, a \in \{1,2\}\}}$
- ▶  $\lambda_i(t|G_i = k) = \lambda_0(t) \exp\{\iota_{i,k,1} + \iota_{i,k,2} t\}$
- ▶ Gompertz baseline  $\lambda_0(t) = \kappa_1 \kappa_2 \exp(\kappa_2 t)$
- ▶  $T_i^* | G_i = k \sim \frac{1}{\iota_{i,k,2} + \kappa_2} \log \left( 1 - \frac{(\iota_{i,k,2} + \kappa_2) \log U_i}{\kappa_1 \kappa_2 \exp \iota_{i,k,1}} \right)$

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application






Competitors  
Simulations  
Results

## Conclusion

## References

# High-dimensional simulation study (2/2)

Lights  
16/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors

## Simulations

Results

## Conclusion

## References

# High-dimensional simulation study (2/2)

Lights  
16/20

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application


Competitors

## Simulations

Results


## Conclusion

## References



# High-dimensional simulation study (2/2)

Lights  
16/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application



Competitors  
**Simulations**  
Results

## Conclusion

## References

# First results (1/2)

Lights  
17/20



## Introduction

- Overview
- Use cases
- Framework

## Model

- Submodels
- Notations
- Likelihood

## Inference

- Penalization
- prox-QNEM
- Evaluation

## Application

- Competitors
- Simulations




## Results

## Conclusion

## References

# First results (1/2)

Lights  
17/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations





## Results

## Conclusion

## References

# First results (1/2)

Lights  
17/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations

## Results







## Conclusion

## References



# First results (1/2)

Lights  
17/20



## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations

## Results


## Conclusion

## References


# First results (2/2)

Lights  
18/20

Support of  $\gamma_0$  and its estimation



Support of  $\gamma_1$  and its estimation



## Introduction

- Overview
- Use cases
- Framework

## Model

- Submodels
- Notations
- Likelihood

## Inference

- Penalization
- prox-QNEM
- Evaluation

## Application

- Competitors
- Simulations

## Results

## Conclusion

## References

# First results (2/2)

Lights  
18/20

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application


Competitors  
Simulations

## Results


## Conclusion

## References


Support of  $\gamma_0$  and its estimation




Support of  $\gamma_1$  and its estimation



Frequency histogram of T



Estimated survival curves



## V. Conclusion

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available
- ▶ Penalization of the likelihood in order to perform feature selection and to prevent overfitting

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available
- ▶ Penalization of the likelihood in order to perform feature selection and to prevent overfitting
- ▶ New efficient estimation algorithm (prox-QNEM) has been derived

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available
- ▶ Penalization of the likelihood in order to perform feature selection and to prevent overfitting
- ▶ New efficient estimation algorithm (prox-QNEM) has been derived
- ▶ Automatically determines significant prognostic longitudinal features

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References



# Conclusion

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available
- ▶ Penalization of the likelihood in order to perform feature selection and to prevent overfitting
- ▶ New efficient estimation algorithm (prox-QNEM) has been derived
- ▶ Automatically determines significant prognostic longitudinal features

## Python 3 package

- ▶ Available at <https://github.com/Califrais/lights>

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

# Conclusion

- ▶ Prognostic method called lights introduced to deal with the problem of joint modeling of longitudinal data and censored durations, where a large number of longitudinal features are available
- ▶ Penalization of the likelihood in order to perform feature selection and to prevent overfitting
- ▶ New efficient estimation algorithm (prox-QNEM) has been derived
- ▶ Automatically determines significant prognostic longitudinal features

## Python 3 package

- ▶ Available at <https://github.com/Califrais/lights>
- ▶ Applications of the model available soon

### Introduction

Overview  
Use cases  
Framework

### Model

Submodels  
Notations  
Likelihood

### Inference

Penalization  
prox-QNEM  
Evaluation

### Application

Competitors  
Simulations  
Results

### Conclusion

### References

Thank you!

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References

- [1] Eleni-Rosalina Andrinopoulou, Dimitris Rizopoulos, Johanna JM Takkenberg, and Emmanuel Lesaffre. Combined dynamic predictions using joint models of two longitudinal outcomes and competing risk data. *Statistical methods in medical research*, 26(4):1787–1801, 2017.
- [2] Yueh-Yun Chi and Joseph G Ibrahim. Joint models for multivariate longitudinal and multivariate survival data. *Biometrics*, 62(2):432–445, 2006.
- [3] Laura A Hatfield, Mark E Boye, and Bradley P Carlin. Joint modeling of multiple longitudinal patient-reported outcomes and survival. *Journal of Biopharmaceutical Statistics*, 21(5):971–991, 2011.
- [4] Cécile Proust-Lima, Viviane Philipps, and Benoit Liqueur. Estimation of extended mixed models using latent classes and latent processes: The r package lcmm. *Journal of Statistical Software, Articles*, 78(2):1–56, 2017. ISSN 1548-7660. doi: 10.18637/jss.v078.i02. URL <https://www.jstatsoft.org/v078/i02>.
- [5] Dimitris Rizopoulos. The r package jmbayes for fitting joint models for longitudinal and time-to-event data using mcmc. *Journal of Statistical Software, Articles*, 72(7):1–46, 2016. ISSN 1548-7660. doi: 10.18637/jss.v072.i07. URL <https://www.jstatsoft.org/v072/i07>.
- [6] Dimitris Rizopoulos and Pulak Ghosh. A bayesian semiparametric multivariate joint model for multiple longitudinal outcomes and a time-to-event. *Statistics in medicine*, 30(12):1366–1380, 2011.
- [7] Glenn T Sueyoshi. Semiparametric proportional hazards estimation of competing risks models with time-varying covariates. *Journal of econometrics*, 51(1-2):25–58, 1992.
- [8] Jeroen K Vermunt and Jay Magidson. Latent class models for classification. *Computational Statistics & Data Analysis*, 41(3-4):531–537, 2003.
- [9] Michael S Wulfsohn and Anastasios A Tsiatis. A joint model for survival and longitudinal data measured with error. *Biometrics*, pages 330–339, 1997.
- [10] Zhongheng Zhang, Jaakko Reinikainen, Kazeem Adedayo Adeleke, Marcel E Pieterse, and Catharina GM Groothuis-Oudshoorn. Time-varying covariates and coefficients in cox regression models. *Annals of translational medicine*, 6(7), 2018.

## Introduction

Overview  
Use cases  
Framework

## Model

Submodels  
Notations  
Likelihood

## Inference

Penalization  
prox-QNEM  
Evaluation

## Application

Competitors  
Simulations  
Results

## Conclusion

## References