

Three Magnetic Dipole Problem

Callebe R. Reis

June 7, 2023

1 Introduction

Magnetic systems often exhibit nonlinear responses, which means that the relationship between an applied magnetic field and the resulting magnetization is not proportional [1]. Nonlinearity can arise due to various factors such as magnetic anisotropy, domain wall motion, and magnetic interactions [2].

This work focuses on investigating the dynamics of a system composed of three equidistantly spaced bar magnets affixed to a table, each free to rotate about its center. Starting from an initial equilibrium state, the system exhibits unexpected and chaotic behavior when a small increase in kinetic energy is applied to a single magnet, resulting in a small initial spin. This behavior depends on specific values of the physical parameters, such as the dipole moments and moments of inertia of the dipoles. The mechanism by which kinetic energy transfers among the magnets in the system is not fully understood. Understanding this phenomenon is crucial for gaining insights into the dynamics of magnet systems and their potential applications.

In order to better analyze this system, we

consider the length of the magnets to be small compared to the distance between them, allowing us to approximate the bar magnets for magnetic dipoles.

A system composed of two magnetic dipoles exhibits a nonlinear coupling, and presents two different time scales depending on the magnitudes of the magnetic interactions, because they are of different nature [4].

Previous works shown that the influence of small fluctuations in a system composed of two magnetic dipoles can lead to two different behaviors: the dipoles fluctuate around stable fixed points (with low amplitude fluctuations), or stochastic reversals occurs between stable fixed points (with strong fluctuations). Low energy fluctuations lead to disjoint basins of attraction near stable fixed points, while higher fluctuations connect basins (including an unstable fixed point) with stochastic reversals and Poisson-distributed waiting time [5].

was extensively studied in [3, 6, 7], in [7] equations for the motion of two dipoles are derived, and [3, 6] investigated the damping version as well as the system in a presence of a external magnetic field.

As any dynamical system, there could be stable or unstable points of equilibrium for this system, and as shown in the work of Carlos for the system with two magnetic dipoles there are eight equilibrium points, and for a ring made out of magnetic dipoles

References

- [1] E. M. Clements, Raja Das, Manh-Huong Phan, Ling Li, Veerle Keppens, David Mandrus, Michael Osofsky, and Hariharan Srikanth. Magnetic field dependence of nonlinear magnetic response and tricritical point in the monoaxial chiral helimagnet $\text{Cr}_{1/3}\text{NbS}_2$. *Phys. Rev. B*, 97:214438, Jun 2018.
- [2] Barnabás Horváth, Péter Decsi, and István Szalai. Nonlinear contributions to the dynamic magnetic susceptibility of magnetic fluids. *Journal of Molecular Liquids*, 359:119279, 2022.
- [3] JG Ku, XY Liu, HH Chen, RD Deng, and QX Yan. Interaction between two magnetic dipoles in a uniform magnetic field. *AIP Advances*, 6(2):025004, 2016.
- [4] D. Laroze, P. Vargas, C. Cortes, and G. Gutierrez. Dynamics of two interacting dipoles. *Journal of Magnetism and Magnetic Materials*, 320(8):1440–1448, 2008.
- [5] Nicolas Plihon, Sophie Miralles, Mickael Bourgoin, and Jean-François Pinton. Stochastic reversal dynamics of two interacting magnetic dipoles: A simple model experiment. *Phys. Rev. E*, 94:012224, Jul 2016.
- [6] Carlos Henrique Moraes Santos. Dinâmica e estabilidade da interação de dois dipolos magnéticos. Master’s thesis, Universidade de Brasília, 2019.
- [7] Kar W Yung, Peter B Landecker, and Daniel D Villani. An analytic solution for the force between two magnetic dipoles. *Magnetic and electrical Separation*, 9, 1970.