

Universal Inference

Behrad Moniri

Dept. of Electrical and Systems Engineering
University of Pennsylvania
`bemoniri@seas.upenn.edu`

April 12, 2022

Universal Inference

Larry Wasserman Aaditya Ramdas Sivaraman Balakrishnan

Department of Statistics and Data Science
Machine Learning Department
Carnegie Mellon University,
Pittsburgh, PA 15213.

`{larry, aramdas, siva}@stat.cmu.edu`

June 4, 2020

Published in the Proceedings of the National Academy of Sciences (PNAS).

- ▶ **Pillars of classical statistics:** Likelihood ratio test, and confidence intervals obtained from asymptotically pivotal estimators.

- ▶ **Pillars of classical statistics:** Likelihood ratio test, and confidence intervals obtained from asymptotically pivotal estimators.
- ▶ These methods rely on large sample asymptotic theory and this often need regularity conditions.

- ▶ **Pillars of classical statistics:** Likelihood ratio test, and confidence intervals obtained from asymptotically pivotal estimators.
- ▶ These methods rely on large sample asymptotic theory and this often need regularity conditions.
- ▶ When these conditions do not hold, there is no general method for statistical inference, with provable guarantees and these settings are typically considered in an *ad-hoc* manner.

- ▶ **One-sentence summary:**

► **One-sentence summary:**

They propose a general method for constructing confidence sets and hypothesis tests that have **finite-sample** guarantees **without** regularity conditions.

► **One-sentence summary:**

They propose a general method for constructing confidence sets and hypothesis tests that have **finite-sample** guarantees **without** regularity conditions. \rightsquigarrow *Universal Inference*.

- ▶ **One-sentence summary:**
They propose a general method for constructing confidence sets and hypothesis tests that have **finite-sample** guarantees **without** regularity conditions. \rightsquigarrow *Universal Inference*.
- ▶ Based on a modified version of the usual likelihood ratio statistic, called “the split likelihood ratio statistics”.

- ▶ **One-sentence summary:**
They propose a general method for constructing confidence sets and hypothesis tests that have **finite-sample** guarantees **without** regularity conditions. \rightsquigarrow *Universal Inference*.
- ▶ Based on a modified version of the usual likelihood ratio statistic, called “the split likelihood ratio statistics”.
- ▶ They also develop various extensions of this basic methods.

- ▶ Consider a parametric family $\{P_\theta : \theta \in \Theta\}$, for some set Θ .

- ▶ Consider a parametric family $\{P_\theta : \theta \in \Theta\}$, for some set Θ .
- ▶ Assume that each distribution has density with respect to some fixed measure μ . Let the corresponding densities be p_θ .

- ▶ Consider a parametric family $\{P_\theta : \theta \in \Theta\}$, for some set Θ .
- ▶ Assume that each distribution has density with respect to some fixed measure μ . Let the corresponding densities be p_θ .
- ▶ We are given $Y_1, \dots, Y_{2n} \sim P_{\theta^*}$ for some $\theta^* \in \Theta$.

- ▶ Consider a parametric family $\{P_\theta : \theta \in \Theta\}$, for some set Θ .
- ▶ Assume that each distribution has density with respect to some fixed measure μ . Let the corresponding densities be p_θ .
- ▶ We are given $Y_1, \dots, Y_{2n} \sim P_{\theta^*}$ for some $\theta^* \in \Theta$.
- ▶ We want to construct confidence intervals for θ^* .

Recap: Regular Models

For regular models, we proceed as follows:

For regular models, we proceed as follows:

- ▶ If $\Theta = \mathbb{R}^d$, set

$$A_n = \left\{ \theta : 2 \log \frac{\mathcal{L}(\hat{\theta})}{\mathcal{L}(\theta)} \leq c_{\alpha,d} \right\},$$

- ▶ $c_{\alpha,d}$ is the α -quantile of a χ_d^2 distribution.
- ▶ $\mathcal{L}(\cdot)$ is the likelihood function.
- ▶ $\hat{\theta}$ is the MLE.

For regular models, we proceed as follows:

- ▶ If $\Theta = \mathbb{R}^d$, set

$$A_n = \left\{ \theta : 2 \log \frac{\mathcal{L}(\hat{\theta})}{\mathcal{L}(\theta)} \leq c_{\alpha,d} \right\},$$

- ▶ $c_{\alpha,d}$ is the α -quantile of a χ_d^2 distribution.
- ▶ $\mathcal{L}(\cdot)$ is the likelihood function.
- ▶ $\hat{\theta}$ is the MLE.

Wilks' Theorem (Wilks, 1938)

For regular models,

$$P_{\theta^*}(\theta^* \in A_n) \rightarrow 1 - \alpha.$$

Universal Confidence Intervals

Confidence Intervals with Split Likelihood-Ratio Statistics

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0 , D_1 randomly.

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0, D_1 randomly.
- ▶ Let $\hat{\theta}_1$ be **any** estimator constructed from D_1 .

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0, D_1 randomly.
- ▶ Let $\hat{\theta}_1$ be **any** estimator constructed from D_1 .
This can be the MLE, a Bayes estimator that utilizes prior knowledge, a robust estimator, etc.

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0, D_1 randomly.
- ▶ Let $\hat{\theta}_1$ be **any** estimator constructed from D_1 .
This can be the MLE, a Bayes estimator that utilizes prior knowledge, a robust estimator, etc.
- ▶ The likelihood function based on D_0 is $\mathcal{L}_0(\theta) = \prod_{i \in D_0} p_\theta(Y_i)$

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0, D_1 randomly.
- ▶ Let $\hat{\theta}_1$ be **any** estimator constructed from D_1 .
This can be the MLE, a Bayes estimator that utilizes prior knowledge, a robust estimator, etc.
- ▶ The likelihood function based on D_0 is $\mathcal{L}_0(\theta) = \prod_{i \in D_0} p_\theta(Y_i)$
- ▶ Define the split likelihood ratio statistic as

$$T_n(\theta) = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta)}$$

Confidence Intervals with Split Likelihood-Ratio Statistics

- ▶ Split data into two sets D_0, D_1 randomly.
- ▶ Let $\hat{\theta}_1$ be **any** estimator constructed from D_1 .
This can be the MLE, a Bayes estimator that utilizes prior knowledge, a robust estimator, etc.
- ▶ The likelihood function based on D_0 is $\mathcal{L}_0(\theta) = \prod_{i \in D_0} p_\theta(Y_i)$
- ▶ Define the split likelihood ratio statistic as

$$T_n(\theta) = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta)}$$

- ▶ The universal confidence set is

$$\mathcal{C}_n = \left\{ \theta \in \Theta : T_n(\theta) \leq \frac{1}{\alpha} \right\}$$

- ▶ If we did not split the data and $\hat{\theta}_1$ was the MLE, then $T_n(\theta)$ would have been the usual likelihood ratio statistic.

- ▶ If we did not split the data and $\hat{\theta}_1$ was the MLE, then $T_n(\theta)$ would have been the usual likelihood ratio statistic.
- ▶ Can we prove an analog of Wilks' theorem here?

- ▶ If we did not split the data and $\hat{\theta}_1$ was the MLE, then $T_n(\theta)$ would have been the usual likelihood ratio statistic.
- ▶ Can we prove an analog of Wilks' theorem here? The answer is yes.

- ▶ If we did not split the data and $\hat{\theta}_1$ was the MLE, then $T_n(\theta)$ would have been the usual likelihood ratio statistic.
- ▶ Can we prove an analog of Wilks' theorem here? The answer is yes.
- ▶ Finding or approximating the distribution of the likelihood ratio statistic is highly nontrivial in irregular models. The split LRS avoids these complications.

Theorem

\mathcal{C}_n is a **finite-sample** valid $1 - \alpha$ confidence set for θ^* , meaning that

$$P_{\theta^*}(\theta^* \in \mathcal{C}_n) \geq 1 - \alpha.$$

The proof is extremely simple.

Proof.

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] = \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right]$$

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\begin{aligned}\mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right] \\ &= \int_A \frac{\prod_{i \in D_0} p_{\psi}(y_i)}{\prod_{i \in D_0} p_{\theta^*}(y_i)} \prod_{i \in D_0} p_{\theta^*}(y_i) dy_1 \cdots dy_n\end{aligned}$$

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\begin{aligned}\mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right] \\ &= \int_A \frac{\prod_{i \in D_0} p_{\psi}(y_i)}{\prod_{i \in D_0} p_{\theta^*}(y_i)} \prod_{i \in D_0} p_{\theta^*}(y_i) dy_1 \cdots dy_n \\ &= \int_A \prod_{i \in D_0} p_{\psi}(y_i) dy_1 \cdots dy_n\end{aligned}$$

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\begin{aligned}
 \mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right] \\
 &= \int_A \frac{\prod_{i \in D_0} p_{\psi}(y_i)}{\prod_{i \in D_0} p_{\theta^*}(y_i)} \prod_{i \in D_0} p_{\theta^*}(y_i) dy_1 \cdots dy_n \\
 &= \int_A \prod_{i \in D_0} p_{\psi}(y_i) dy_1 \cdots dy_n \leq \prod_{i \in D_0} \left[\int p_{\psi}(y_i) dy_i \right]
 \end{aligned}$$

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\begin{aligned}\mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right] \\ &= \int_A \frac{\prod_{i \in D_0} p_{\psi}(y_i)}{\prod_{i \in D_0} p_{\theta^*}(y_i)} \prod_{i \in D_0} p_{\theta^*}(y_i) dy_1 \cdots dy_n \\ &= \int_A \prod_{i \in D_0} p_{\psi}(y_i) dy_1 \cdots dy_n \leq \prod_{i \in D_0} \left[\int p_{\psi}(y_i) dy_i \right] = 1\end{aligned}$$

Proof.

Consider any fixed $\psi \in \Theta$ and let A denote the support of P_{θ^*} .

$$\begin{aligned}
 \mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\psi)}{\mathcal{L}_0(\theta^*)} \right] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} p_{\psi}(Y_i)}{\prod_{i \in D_0} p_{\theta^*}(Y_i)} \right] \\
 &= \int_A \frac{\prod_{i \in D_0} p_{\psi}(y_i)}{\prod_{i \in D_0} p_{\theta^*}(y_i)} \prod_{i \in D_0} p_{\theta^*}(y_i) dy_1 \cdots dy_n \\
 &= \int_A \prod_{i \in D_0} p_{\psi}(y_i) dy_1 \cdots dy_n \leq \prod_{i \in D_0} \left[\int p_{\psi}(y_i) dy_i \right] = 1
 \end{aligned}$$

$\hat{\theta}_1$ is fixed when we condition on D_1 . So we have

$$\mathbb{E}_{\theta^*} [T_n(\theta^*) \mid D_1] = \mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta^*)} \mid D_1 \right] \leq 1.$$

Now, using Markov's inequality,

$$P_{\theta^*}(\theta^* \notin \mathcal{C}_n) = P_{\theta^*}\left(T_n(\theta^*) > \frac{1}{\alpha}\right) \leq \alpha \mathbb{E}_{\theta^*}[T_n(\theta^*)]$$

Now, using Markov's inequality,

$$\begin{aligned} P_{\theta^*}(\theta^* \notin \mathcal{C}_n) &= P_{\theta^*}\left(T_n(\theta^*) > \frac{1}{\alpha}\right) \leq \alpha \mathbb{E}_{\theta^*}[T_n(\theta^*)] \\ &= \alpha \mathbb{E}_{\theta^*}\left[\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta^*)}\right] = \alpha \mathbb{E}_{\theta^*}\left(\mathbb{E}_{\theta^*}\left[\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta^*)} \middle| D_1\right]\right) \leq \alpha \end{aligned}$$

This completes the proof.

- ▶ The parametric setup adopted above generalizes easily to nonparametric settings as long as we can calculate a likelihood.

- ▶ The parametric setup adopted above generalizes easily to nonparametric settings as long as we can calculate a likelihood.
- ▶ For a collection of densities \mathcal{P} , and a true density $p^* \in \mathcal{P}$, suppose we use D_1 to identify $\hat{p}_1 \in \mathcal{P}$, and D_0 to calculate

$$T_n(p) = \prod_{i \in D_0} \frac{\hat{p}_1(Y_i)}{p(Y_i)}.$$

- ▶ We then define, $\mathcal{C}_n = \{p \in \mathcal{P} : T_n(p) \leq \frac{1}{\alpha}\}$, and our previous argument ensures that

$$P_{p^*}(p^* \in \mathcal{C}_n) \geq 1 - \alpha.$$

Universal Hypothesis Testing

- ▶ Let $\Theta_0 \subset \Theta$ be a null-set and consider testing

$$H_0 : \theta^* \in \Theta_0 \quad \text{versus} \quad \theta^* \notin \Theta_0$$

- ▶ **Using the duality between hypothesis testing and confidence intervals:**

We simply reject the null hypothesis if $\mathcal{C}_n \cap \Theta_0 = \emptyset$. The type I error of this test is clearly at most α .

- ▶ Let $\Theta_0 \subset \Theta$ be a null-set and consider testing

$$H_0 : \theta^* \in \Theta_0 \quad \text{versus} \quad \theta^* \notin \Theta_0$$

- ▶ **Using the duality between hypothesis testing and confidence intervals:**

We simply reject the null hypothesis if $\mathcal{C}_n \cap \Theta_0 = \emptyset$. The type I error of this test is clearly at most α .

- ▶ Can we find a computationally efficient way?

- ▶ Let $\hat{\theta}_1$ be any estimator constructed from D_1 .

- ▶ Let $\hat{\theta}_1$ be any estimator constructed from D_1 .
- ▶ Let $\hat{\theta}_0 := \operatorname{argmax}_{\theta \in \Theta_0} \mathcal{L}_0(\theta)$ be the MLE under null from D_0 .

- ▶ Let $\hat{\theta}_1$ be any estimator constructed from D_1 .
- ▶ Let $\hat{\theta}_0 := \operatorname{argmax}_{\theta \in \Theta_0} \mathcal{L}_0(\theta)$ be the MLE under null from D_0 .
- ▶ Reject H_0 if

$$\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} > \frac{1}{\alpha}.$$

- ▶ Let $\hat{\theta}_1$ be any estimator constructed from D_1 .
- ▶ Let $\hat{\theta}_0 := \operatorname{argmax}_{\theta \in \Theta_0} \mathcal{L}_0(\theta)$ be the MLE under null from D_0 .
- ▶ Reject H_0 if

$$\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} > \frac{1}{\alpha}.$$

Theorem

This test controls the type I error at level α .

- ▶ Let $\hat{\theta}_1$ be any estimator constructed from D_1 .
- ▶ Let $\hat{\theta}_0 := \operatorname{argmax}_{\theta \in \Theta_0} \mathcal{L}_0(\theta)$ be the MLE under null from D_0 .
- ▶ Reject H_0 if

$$\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} > \frac{1}{\alpha}.$$

Theorem

This test controls the type I error at level α .

Proof.

The proof is one line.

$$P_{\theta^*} \left(\mathcal{L}_0(\hat{\theta}_1) / \mathcal{L}_0(\hat{\theta}_0) > 1/\alpha \right) \leq \alpha \mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} \right] \leq \alpha \mathbb{E}_{\theta^*} \left[\frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\theta^*)} \right] \leq \alpha$$

Some Discussions

► **Regular models:**

Compare the log-likelihood ratio to the $(1 - \alpha)$ -quantile of a χ^2 distribution (dof = dimension of null - dimension of alternative)

- ▶ **Regular models:**

Compare the log-likelihood ratio to the $(1 - \alpha)$ -quantile of a χ^2 distribution (dof = dimension of null - dimension of alternative)

- ▶ **This paper:**

Compare the **split**-log-split-likelihood ratio to $\log(1/\alpha) \rightsquigarrow (1 - \alpha)$ -quantile of a χ^2 distribution with **one** degree of freedom.

- ▶ You are only using Markov?! This isn't tight enough!

- ▶ You are only using Markov?! This isn't tight enough!
Yes and No!

- ▶ You are only using Markov?! This isn't tight enough!
Yes and No!
- ▶ We are really using the fact that $\log \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$ has an exponential tail, just as an asymptotic argument would.

- ▶ You are only using Markov?! This isn't tight enough!
Yes and No!
- ▶ We are really using the fact that $\log \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$ has an exponential tail, just as an asymptotic argument would.
- ▶ In true Chernoff bounds:

$$\mathbb{E}_{\theta^*} \left[\exp \left(a \log \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} \right) \right] \leq \text{MGF of } \chi^2, \mathcal{N}, \dots$$

- ▶ One should view this proof as a **poor man's Chernoff bound**:

$$\mathbb{E}_{\theta^*} \left[\exp \left(\log \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)} \right) \right] \leq 1$$

Sanity Check: Regular Models

- Suppose that $Y_1, \dots, Y_n \sim \mathcal{N}_d(\theta, I)$ where $\theta \in \mathbb{R}^d$.

- ▶ Suppose that $Y_1, \dots, Y_n \sim \mathcal{N}_d(\theta, I)$ where $\theta \in \mathbb{R}^d$.
- ▶ Let $c_{\alpha,d}$ and z_α denote the upper α quantiles of the χ_d^2 and standard Gaussian respectively.

- ▶ Suppose that $Y_1, \dots, Y_n \sim \mathcal{N}_d(\theta, I)$ where $\theta \in \mathbb{R}^d$.
- ▶ Let $c_{\alpha,d}$ and z_α denote the upper α quantiles of the χ_d^2 and standard Gaussian respectively.
- ▶ The usual confidence set for θ based on the LRT can be computed as follows:
 - ▶ The likelihood function and MLE:

$$\mathcal{L}(\theta) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(Y_i - \mu)^2}{2}\right), \quad \hat{\theta}_{MLE} = \bar{Y}$$

$$\begin{aligned} A_n &= \left\{ \theta : \|\theta - \bar{Y}\|^2 \leq \frac{c_{\alpha,d}}{n} \right\} \\ &= \left\{ \theta : \|\theta - \bar{Y}\|^2 \leq \frac{d + \sqrt{2d}z_\alpha + o(\sqrt{d})}{n} \right\}. \end{aligned}$$

- Denoting the sample means \bar{Y}_1 and \bar{Y}_0 we see that:

$$\log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) = - \left(\frac{n}{2}\right) \frac{\|\bar{Y}_0 - \bar{Y}_1\|^2}{2} + \left(\frac{n}{2}\right) \frac{\|\theta - \bar{Y}_0\|^2}{2}.$$

- ▶ Denoting the sample means \bar{Y}_1 and \bar{Y}_0 we see that:

$$\log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) = -\left(\frac{n}{2}\right) \frac{\|\bar{Y}_0 - \bar{Y}_1\|^2}{2} + \left(\frac{n}{2}\right) \frac{\|\theta - \bar{Y}_0\|^2}{2}.$$

- ▶ The universal confidence set is

$$\begin{aligned} C_n &= \{\theta : \log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) \leq \log(1/\alpha)\} \\ &= \left\{ \theta : \|\theta - \bar{Y}_0\|^2 \leq \frac{4}{n} \log\left(\frac{1}{\alpha}\right) + \|\bar{Y}_0 - \bar{Y}_1\|^2 \right\}. \end{aligned}$$

- ▶ Denoting the sample means \bar{Y}_1 and \bar{Y}_0 we see that:

$$\log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) = - \left(\frac{n}{2}\right) \frac{\|\bar{Y}_0 - \bar{Y}_1\|^2}{2} + \left(\frac{n}{2}\right) \frac{\|\theta - \bar{Y}_0\|^2}{2}.$$

- ▶ The universal confidence set is

$$\begin{aligned} C_n &= \{\theta : \log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) \leq \log(1/\alpha)\} \\ &= \left\{ \theta : \|\theta - \bar{Y}_0\|^2 \leq \frac{4}{n} \log\left(\frac{1}{\alpha}\right) + \|\bar{Y}_0 - \bar{Y}_1\|^2 \right\}. \end{aligned}$$

- ▶ Note that $\|\bar{Y}_0 - \bar{Y}_1\|^2 = O_p(d/n)$, so both sets have radii $O_p(d/n)$.

- ▶ Denoting the sample means \bar{Y}_1 and \bar{Y}_0 we see that:

$$\log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) = - \left(\frac{n}{2}\right) \frac{\|\bar{Y}_0 - \bar{Y}_1\|^2}{2} + \left(\frac{n}{2}\right) \frac{\|\theta - \bar{Y}_0\|^2}{2}.$$

- ▶ The universal confidence set is

$$\begin{aligned} C_n &= \{\theta : \log \mathcal{L}_0(\bar{Y}_1) - \log \mathcal{L}_0(\theta) \leq \log(1/\alpha)\} \\ &= \left\{ \theta : \|\theta - \bar{Y}_0\|^2 \leq \frac{4}{n} \log\left(\frac{1}{\alpha}\right) + \|\bar{Y}_0 - \bar{Y}_1\|^2 \right\}. \end{aligned}$$

- ▶ Note that $\|\bar{Y}_0 - \bar{Y}_1\|^2 = O_p(d/n)$, so both sets have radii $O_p(d/n)$.
- ▶ For constant α , the radius is four times larger.

1. **Identifiable:** any $\theta \neq \theta^*$ it is the case that $P_\theta \neq P_{\theta^*}$.
2. Differentiable in quadratic mean (**DQM**) at θ^* : there exists a function s_{θ^*} such that:

$$\int \left[\sqrt{p_\theta} - \sqrt{p_{\theta^*}} - \frac{1}{2}(\theta - \theta^*)^T s_{\theta^*} \sqrt{p_{\theta^*}} \right]^2 d\mu = o(\|\theta - \theta^*\|^2), \text{ as } \theta \rightarrow \theta^*.$$

3. The parameter space $\Theta \subset \mathbb{R}^d$ is **compact**.
4. **Smoothness:** There is a function ℓ with $\sup_\theta \mathbb{E}_{x \sim P_\theta} \ell^2(X) < \infty$ s.t.

$$\forall \theta_1, \theta_2 \in \Theta : |\log p_{\theta_1}(x) - \log p_{\theta_2}(x)| \leq \ell(x) \|\theta_1 - \theta_2\|.$$

5. A consequence of the DQM condition is that the Fisher information matrix is well-defined, and we assume it is **non-degenerate**.

Theorem

Under the regularity conditions in the previous slide, and $\|\hat{\theta}_1 - \theta^\| = O_p(1/\sqrt{n})$, the split LRT has diameter $O_p(\sqrt{\log(1/\delta)/n})$*

Theorem

Under the regularity conditions in the previous slide, and $\|\hat{\theta}_1 - \theta^\| = O_p(1/\sqrt{n})$, the split LRT has diameter $O_p(\sqrt{\log(1/\delta)/n})$*

Proof.

The high level idea: it suffices to show that for all θ sufficiently far from θ^* , we have

$$\frac{\mathcal{L}_0(\theta)}{\mathcal{L}_0(\hat{\theta}_1)} \leq \alpha.$$



Example of an Irregular Model

- ▶ Let $Y_1, \dots, Y_{2n} \sim P$ where $Y_i \in \mathbb{R}$.
- ▶ We want to test

$$H_0 : P \in \mathcal{M}_1 \text{ versus } H_1 : P \in \mathcal{M}_2,$$

where \mathcal{M}_k denotes the set of mixtures of k Gaussians, with an appropriately restricted parameter space Θ .

- ▶ Let $Y_1, \dots, Y_{2n} \sim P$ where $Y_i \in \mathbb{R}$.
- ▶ We want to test

$$H_0 : P \in \mathcal{M}_1 \text{ versus } H_1 : P \in \mathcal{M}_2,$$

where \mathcal{M}_k denotes the set of mixtures of k Gaussians, with an appropriately restricted parameter space Θ .

- ▶ LRT has an intractable limiting distribution. There is no known confidence set for mixture problems with guaranteed coverage properties.

- ▶ The true model is assumed to be $\frac{1}{2}\phi(y; -\mu, 1) + \frac{1}{2}\phi(y; \mu, 1)$
- ▶ The null: $\mu = 0$. We set $\alpha = 0.1$ and $n = 200$.
- ▶ Let $\hat{\theta}_1$ be the MLE under \mathcal{M}_2 .
- ▶ This MLE is calculated using the EM algorithm (does it converge? IDK!)




Figure: *Black = Universal / Red = Bootstrap*




Figure: Black = Universal / Red = Bootstrap

The bootstrap test does not have any guarantee on the type I error.

Extensions

- ▶ The universal method involves randomly splitting the data and the final inferences will depend on the randomness of the split.

- ▶ The universal method involves randomly splitting the data and the final inferences will depend on the randomness of the split.
- ▶ For the test to work, we needed $\mathbb{E}_{\theta^*}[T_n] \leq 1$ where $T_n = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$.

- ▶ The universal method involves randomly splitting the data and the final inferences will depend on the randomness of the split.
- ▶ For the test to work, we needed $\mathbb{E}_{\theta^*}[T_n] \leq 1$ where $T_n = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$.
- ▶ Imagine that we obtained B such statistics $T_{n,1}, \dots, T_{n,B}$ with the same property. Let

$$\bar{T}_n = B^{-1} \sum_{j=1}^B T_{n,j}.$$

Then we still have that $\mathbb{E}_{\theta^*}[\bar{T}_n]$.

- ▶ The universal method involves randomly splitting the data and the final inferences will depend on the randomness of the split.
- ▶ For the test to work, we needed $\mathbb{E}_{\theta^*}[T_n] \leq 1$ where $T_n = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$.
- ▶ Imagine that we obtained B such statistics $T_{n,1}, \dots, T_{n,B}$ with the same property. Let

$$\bar{T}_n = B^{-1} \sum_{j=1}^B T_{n,j}.$$

Then we still have that $\mathbb{E}_{\theta^*}[\bar{T}_n]$.

- ▶ K-fold and All split.

- ▶ The universal method involves randomly splitting the data and the final inferences will depend on the randomness of the split.
- ▶ For the test to work, we needed $\mathbb{E}_{\theta^*}[T_n] \leq 1$ where $T_n = \frac{\mathcal{L}_0(\hat{\theta}_1)}{\mathcal{L}_0(\hat{\theta}_0)}$.
- ▶ Imagine that we obtained B such statistics $T_{n,1}, \dots, T_{n,B}$ with the same property. Let

$$\bar{T}_n = B^{-1} \sum_{j=1}^B T_{n,j}.$$

Then we still have that $\mathbb{E}_{\theta^*}[\bar{T}_n]$.

- ▶ K-fold and All split.
- ▶ **Broader Impact:**
These methods will potentially lead to cherry-picking :)

- ▶ Computing the maximum likelihood (under the null) is sometimes computationally hard.

- ▶ Computing the maximum likelihood (under the null) is sometimes computationally hard.
- ▶ Suppose one could come up with a relaxation F_0 of the null likelihood \mathcal{L}_0 :

$$\max_{\theta} F_0(\theta) \geq \max_{\theta} \mathcal{L}_0(\theta).$$

- ▶ Computing the maximum likelihood (under the null) is sometimes computationally hard.
- ▶ Suppose one could come up with a relaxation F_0 of the null likelihood \mathcal{L}_0 :

$$\max_{\theta} F_0(\theta) \geq \max_{\theta} \mathcal{L}_0(\theta).$$

- ▶ Define $\hat{\theta}_0^F := \operatorname{argmax}_{\theta} F_0(\theta)$, and consider the statistics

$$T'_n := \frac{\mathcal{L}_0(\hat{\theta}_1)}{F_0(\hat{\theta}_0^F)}$$

- ▶ Computing the maximum likelihood (under the null) is sometimes computationally hard.
- ▶ Suppose one could come up with a relaxation F_0 of the null likelihood \mathcal{L}_0 :

$$\max_{\theta} F_0(\theta) \geq \max_{\theta} \mathcal{L}_0(\theta).$$

- ▶ Define $\hat{\theta}_0^F := \operatorname{argmax}_{\theta} F_0(\theta)$, and consider the statistics

$$T'_n := \frac{\mathcal{L}_0(\hat{\theta}_1)}{F_0(\hat{\theta}_0^F)}$$

- ▶ then the split LRT may proceed using T' instead of T . This is because $F(\hat{\theta}_0^F) \geq \mathcal{L}(\hat{\theta})$, and hence $T'_n \leq T_n$.

- Sometimes the MLE may not exist since the likelihood function is unbounded. A smoothed likelihood has been proposed as an alternative.

- ▶ Sometimes the MLE may not exist since the likelihood function is unbounded. A smoothed likelihood has been proposed as an alternative.
- ▶ Consider a kernel $k(x, y)$ such that $\int k(x, y) dy = 1$ for any x .

$$\tilde{p}_\theta(y) := \int k(x, y) p_\theta(x) dx.$$

- ▶ Sometimes the MLE may not exist since the likelihood function is unbounded. A smoothed likelihood has been proposed as an alternative.
- ▶ Consider a kernel $k(x, y)$ such that $\int k(x, y)dy = 1$ for any x .

$$\tilde{p}_\theta(y) := \int k(x, y)p_\theta(x)dx.$$

- ▶ Denote the smoothed empirical density based on D_0 as

$$\tilde{p}_n := \frac{1}{|D_0|} \sum_{i \in D_0} k(X_i, \cdot).$$

- ▶ Sometimes the MLE may not exist since the likelihood function is unbounded. A smoothed likelihood has been proposed as an alternative.
- ▶ Consider a kernel $k(x, y)$ such that $\int k(x, y) dy = 1$ for any x .

$$\tilde{p}_\theta(y) := \int k(x, y) p_\theta(x) dx.$$

- ▶ Denote the smoothed empirical density based on D_0 as

$$\tilde{p}_n := \frac{1}{|D_0|} \sum_{i \in D_0} k(X_i, \cdot).$$

- ▶ Define the smoothed likelihood on D_0 as

$$\tilde{\mathcal{L}}_0(\theta) := \prod_{i \in D_0} \exp \int k(X_i, y) \log \tilde{p}_\theta(y) dy \rightsquigarrow \tilde{\theta}_0 := \arg \min_{\theta \in \Theta_0} KL(\tilde{p}_n, \tilde{p}_\theta)$$

- ▶ As before, let $\hat{\theta}_1 \in \Theta$ be any estimator based on D_1 . The smoothed split LRT:

$$\text{reject } H_0 \text{ if } \tilde{U}_n > 1/\alpha, \text{ where } \tilde{U}_n = \frac{\tilde{\mathcal{L}}_0(\hat{\theta}_1)}{\tilde{\mathcal{L}}_0(\tilde{\theta}_0)}.$$

- ▶ As before, let $\hat{\theta}_1 \in \Theta$ be any estimator based on D_1 . The smoothed split LRT:

$$\text{reject } H_0 \text{ if } \tilde{U}_n > 1/\alpha, \text{ where } \tilde{U}_n = \frac{\tilde{\mathcal{L}}_0(\hat{\theta}_1)}{\tilde{\mathcal{L}}_0(\tilde{\theta}_0)}.$$

Fix $\psi \in \Theta$, we have

$$\begin{aligned} \mathbb{E}_{\theta^*} \left[\frac{\tilde{\mathcal{L}}_0(\psi)}{\tilde{\mathcal{L}}_0(\tilde{\theta}_0)} \right] &\stackrel{(i)}{\leq} \mathbb{E}_{\theta^*} \left[\frac{\tilde{\mathcal{L}}_0(\psi)}{\tilde{\mathcal{L}}_0(\theta^*)} \right] = \mathbb{E}_{\theta^*} \left[\frac{\prod_{i \in D_0} \exp \int k(X_i, y) \log \tilde{p}_\psi(y) dy}{\prod_{i \in D_0} \exp \int k(X_i, y) \log \tilde{p}_{\theta^*}(y) dy} \right] \\ &= \prod_{i \in D_0} \int \exp \left(\int k(x, y) \log \frac{\tilde{p}_\psi(y)}{\tilde{p}_{\theta^*}(y)} dy \right) p_{\theta^*}(x) dx \leq \dots \leq 1. \end{aligned}$$

Sequential Testing

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .
- ▶ Let $\hat{\theta}_{1,t-1}$ be any *non-anticipating* estimator based on the first $t - 1$ samples.

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .
- ▶ Let $\hat{\theta}_{1,t-1}$ be any *non-anticipating* estimator based on the first $t-1$ samples.
- ▶ Denote the null MLE as $\hat{\theta}_{0,t} = \arg \max_{\theta \in \Theta_0} \prod_{i=1}^t p_{\theta}(Y_i)$.

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .
- ▶ Let $\hat{\theta}_{1,t-1}$ be any *non-anticipating* estimator based on the first $t-1$ samples.
- ▶ Denote the null MLE as $\hat{\theta}_{0,t} = \arg \max_{\theta \in \Theta_0} \prod_{i=1}^t p_{\theta}(Y_i)$.
- ▶ At any time t , reject the null and stop if

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} > 1/\alpha.$$

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .
- ▶ Let $\hat{\theta}_{1,t-1}$ be any *non-anticipating* estimator based on the first $t-1$ samples.
- ▶ Denote the null MLE as $\hat{\theta}_{0,t} = \arg \max_{\theta \in \Theta_0} \prod_{i=1}^t p_{\theta}(Y_i)$.
- ▶ At any time t , reject the null and stop if

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} > 1/\alpha.$$

- ▶ Let τ_{θ} denote the stopping time when the data is drawn from P_{θ} .

- ▶ Consider the following, standard, sequential testing/estimation setup:
- ▶ We observe an i.i.d. sequence Y_1, Y_2, \dots from P_{θ^*} .
- ▶ Let $\hat{\theta}_{1,t-1}$ be any *non-anticipating* estimator based on the first $t-1$ samples.
- ▶ Denote the null MLE as $\hat{\theta}_{0,t} = \arg \max_{\theta \in \Theta_0} \prod_{i=1}^t p_{\theta}(Y_i)$.
- ▶ At any time t , reject the null and stop if

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} > 1/\alpha.$$

- ▶ Let τ_{θ} denote the stopping time when the data is drawn from P_{θ} .

Theorem

The running MLE LRT has type I error at most α , meaning that $\sup_{\theta^* \in \Theta_0} P_{\theta^*}(\tau_{\theta^*} < \infty) \leq \alpha$.

- For M_t we can write:

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} \leq \underbrace{\frac{\prod_{i=1}^t p_{\hat{\theta}_{i-1}}(Y_i)}{\prod_{i=1}^t p_{\theta^*}(Y_i)}}_{L_t} = L_{t-1} \frac{p_{\hat{\theta}_{t-1}}(Y_t)}{p_{\theta^*}(Y_t)}.$$

- For M_t we can write:

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} \leq \underbrace{\frac{\prod_{i=1}^t p_{\hat{\theta}_{i-1}}(Y_i)}{\prod_{i=1}^t p_{\theta^*}(Y_i)}}_{L_t} = L_{t-1} \frac{p_{\hat{\theta}_{t-1}}(Y_t)}{p_{\theta^*}(Y_t)}.$$

- It is easy to verify that L_t is a nonnegative super-martingale with respect to the natural filtration $\mathcal{F}_t = \sigma(Y_1, \dots, Y_t)$:

$$\begin{aligned} \mathbb{E}_{\theta^*}[L_t | \mathcal{F}_{t-1}] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i=1}^t p_{\hat{\theta}_{i-1}}(Y_i)}{\prod_{i=1}^t p_{\theta^*}(Y_i)} \middle| \mathcal{F}_{t-1} \right] \\ &= L_{t-1} \mathbb{E}_{\theta^*} \left[\frac{p_{\hat{\theta}_{t-1}}(Y_t)}{p_{\theta^*}(Y_t)} \middle| \mathcal{F}_{t-1} \right] \leq L_{t-1} \rightsquigarrow \text{Super-Martingale} \end{aligned}$$

- For M_t we can write:

$$M_t := \frac{\prod_{i=1}^t p_{\hat{\theta}_{1,i-1}}(Y_i)}{\prod_{i=1}^t p_{\hat{\theta}_{0,t}}(Y_i)} \leq \underbrace{\frac{\prod_{i=1}^t p_{\hat{\theta}_{i-1}}(Y_i)}{\prod_{i=1}^t p_{\theta^*}(Y_i)}}_{L_t} = L_{t-1} \frac{p_{\hat{\theta}_{t-1}}(Y_t)}{p_{\theta^*}(Y_t)}.$$

- It is easy to verify that L_t is a nonnegative super-martingale with respect to the natural filtration $\mathcal{F}_t = \sigma(Y_1, \dots, Y_t)$:

$$\begin{aligned} \mathbb{E}_{\theta^*}[L_t | \mathcal{F}_{t-1}] &= \mathbb{E}_{\theta^*} \left[\frac{\prod_{i=1}^t p_{\hat{\theta}_{i-1}}(Y_i)}{\prod_{i=1}^t p_{\theta^*}(Y_i)} \middle| \mathcal{F}_{t-1} \right] \\ &= L_{t-1} \mathbb{E}_{\theta^*} \left[\frac{p_{\hat{\theta}_{t-1}}(Y_t)}{p_{\theta^*}(Y_t)} \middle| \mathcal{F}_{t-1} \right] \leq L_{t-1} \rightsquigarrow \text{Super-Martingale} \end{aligned}$$

- Now we proceed as follows:

$$P_{\theta^*}(\exists t \in \mathbb{N} : M_t > 1/\alpha) \leq P_{\theta^*}(\exists t \in \mathbb{N} : L_t > 1/\alpha)$$

$$\stackrel{(*)}{\leq} \mathbb{E}_{\theta^*}[L_0] \cdot \alpha = \alpha$$

Theorem [Ville (1939)]

For any nonnegative supermartingale L_t and any $x > 1$, we have

$$\mathbb{P}[\exists t : L_t \geq x] \leq \frac{\mathbb{E}[L_0]}{x}$$

Proof.

The idea is to consider the following stopping time

$$N = \inf\{t \geq 1 : L_t \geq x\},$$

and use the optional stopping time theorem. □

Conclusion

- ▶ Inference based on the split likelihood ratio statistic (and variants) leads to simple tests and confidence sets with finite-sample guarantees.

- ▶ Inference based on the split likelihood ratio statistic (and variants) leads to simple tests and confidence sets with finite-sample guarantees.
 - ▶ These methods are most useful in problems where standard asymptotic methods are difficult/impossible to apply.
-
- ▶ **Going forward:** Optimality? Power of the Test?
How does the choice of $\hat{\theta}_1$ affect the power of the test?

Thank You!