

# The Well-Calibrated Bayesian

---

Ryan Brill

February 17, 2022

# 1. Intro

- Subjective probability forecasting is well established among meteorologists

# 1. Intro

- Subjective probability forecasting is well established among meteorologists
- Weather forecasts make predictions such as “there is a 30% chance of rain in Philly tonight”

# 1. Intro

- Subjective probability forecasting is well established among meteorologists
- Weather forecasts make predictions such as “there is a 30% chance of rain in Philly tonight”
- *Subjectivists* (also known as *Bayesians*) believe that a probability forecast is a measure of a forecaster’s “degree of belief” given their information at the time of the forecast

# 1. Intro

- Subjective probability forecasting is well established among meteorologists
- Weather forecasts make predictions such as “there is a 30% chance of rain in Philly tonight”
- *Subjectivists* (also known as *Bayesians*) believe that a probability forecast is a measure of a forecaster’s “degree of belief” given their information at the time of the forecast
- Even if a forecaster bases their predictions on a frequentist statistical model, a Subjectivist still believes that the forecaster *believes* in their own model

# 1. Intro

- Subjective probability forecasting is well established among meteorologists
- Weather forecasts make predictions such as “there is a 30% chance of rain in Philly tonight”
- *Subjectivists* (also known as *Bayesians*) believe that a probability forecast is a measure of a forecaster’s “degree of belief” given their information at the time of the forecast
- Even if a forecaster bases their predictions on a frequentist statistical model, a Subjectivist still believes that the forecaster *believes* in their own model
- A Bayesian has a joint probability distribution over all conceivably observable quantities.

# 1. Intro

- Subjective probability forecasting is well established among meteorologists
- Weather forecasts make predictions such as “there is a 30% chance of rain in Philly tonight”
- *Subjectivists* (also known as *Bayesians*) believe that a probability forecast is a measure of a forecaster’s “degree of belief” given their information at the time of the forecast
- Even if a forecaster bases their predictions on a frequentist statistical model, a Subjectivist still believes that the forecaster *believes* in their own model
- A Bayesian has a joint probability distribution over all conceivably observable quantities.
- Forecasting is then a matter of summarizing the conditional distribution of quantities still unobserved, given current information

# 1. Intro

- A weather forecaster is well calibrated if, for example, of those events to which he assigns a probability 30 percent (say, the event that it will rain tomorrow), the long-run proportion that actually occurs turns out to be 30 percent



# 1. Intro

- A weather forecaster is well calibrated if, for example, of those events to which he assigns a probability 30 percent (say, the event that it will rain tomorrow), the long-run proportion that actually occurs turns out to be 30 percent
- This is not about the “true” probability of rain on a given day, or about any underlying “objective” probability; this refers only to observed long run proportions

# 1. Intro

- A weather forecaster is well calibrated if, for example, of those events to which he assigns a probability 30 percent (say, the event that it will rain tomorrow), the long-run proportion that actually occurs turns out to be 30 percent
- This is not about the “true” probability of rain on a given day, or about any underlying “objective” probability; this refers only to observed long run proportions
- A weather forecaster is coherent if their set of beliefs is internally consistent, or obeys the laws of probability theory

# 1. Intro

- Today we investigate a forecaster's view of their own calibration

# 1. Intro

- Today we investigate a forecaster's view of their own calibration
- Today we discuss the relationship between coherence and calibration

# 1. Intro

- Today we investigate a forecaster's view of their own calibration
- Today we discuss the relationship between coherence and calibration
- Paper: *The Well-Calibrated Bayesian* by A. P. Dawid

## The Well-Calibrated Bayesian

A. P. DAWID\*

---

Suppose that a forecaster sequentially assigns probabilities to events. He is *well calibrated* if, for example, of those events to which he assigns a probability 30 percent, the long-run proportion that actually occurs turns out to be 30 percent. We prove a theorem to the effect that a coherent Bayesian expects to be well calibrated, and consider its destructive implications for the theory of coherence.

**KEY WORDS:** Calibration; Coherence; Martingale; Probability forecasting; Subjectivism; Weather forecasting.

## 2a. Patrick the Forecaster

- Meet Patrick

## 2a. Patrick the Forecaster

- Meet Patrick
- Patrick likes to predict the future



## 2a. Patrick the Forecaster

- Meet Patrick
- Patrick likes to predict the future
- Patrick uses his subjective probability distribution  $\Pi$  to predict the future

## 2a. Patrick the Forecaster



## 2a. Patrick the Forecaster



## 2a. Patrick the Forecaster

- Consider the assessment of “seemingly unrelated” uncertain events: I will cook pasta tonight, Shiba Inu Coin will go down tomorrow, the Rams will win the Super Bowl again next year, etc.

## 2a. Patrick the Forecaster

- Consider the assessment of “seemingly unrelated” uncertain events: I will cook pasta tonight, Shiba Inu Coin will go down tomorrow, the Rams will win the Super Bowl again next year, etc.
- It's natural for Patrick to regard these quantities as probabilistically independent

## 2a. Patrick the Forecaster

- Consider the assessment of “seemingly unrelated” uncertain events: I will cook pasta tonight, Shiba Inu Coin will go down tomorrow, the Rams will win the Super Bowl again next year, etc.
- It's natural for Patrick to regard these quantities as probabilistically independent
- Patrick predicts a probability distribution over the outcomes, and then computes the median

## 2a. Patrick the Forecaster

- Consider Patrick's dilemma when he learns that a large number  $N$  of his forecasts, say 80%, exceed the empirical proportion

## 2a. Patrick the Forecaster

- Consider Patrick's dilemma when he learns that a large number  $N$  of his forecasts, say 80%, exceed the empirical proportion
- Under  $\Pi$ , this event has subjective probability  $\text{Binomial}(N, \frac{1}{2})$ , and so is vanishingly small



## 2a. Patrick the Forecaster

- Consider Patrick's dilemma when he learns that a large number  $N$  of his forecasts, say 80%, exceed the empirical proportion
- Under  $\Pi$ , this event has subjective probability  $\text{Binomial}(N, \frac{1}{2})$ , and so is vanishingly small
- Yet it occurred!

## 2a. Patrick the Forecaster

- Possibility 1:  $\Pi$  is not coherent ( $\Pi$  does not obey the laws of probability theory)

## 2a. Patrick the Forecaster

- Possibility 1:  $\Pi$  is not coherent ( $\Pi$  does not obey the laws of probability theory)
- Possibility 2:  $\Pi$  is coherent, but these “seemingly unrelated” events, which were regarded as subjectively independent by  $\Pi$ , are actually related, because they are all being made by Patrick

## 2a. Patrick the Forecaster

- Possibility 1:  $\Pi$  is not coherent ( $\Pi$  does not obey the laws of probability theory)
- Possibility 2:  $\Pi$  is coherent, but these “seemingly unrelated” events, which were regarded as subjectively independent by  $\Pi$ , are actually related, because they are all being made by Patrick
  - Patrick is a potentially miscalibrated individual - he is not sure whether his future subjective probability assignments will agree with observed frequency

## 2b. Patrick the Weather Forecaster

- Now, Patrick is a *weather* forecaster

## 2b. Patrick the Weather Forecaster

- Now, Patrick is a *weather* forecaster
- A weather forecaster's task is sequential

## 2b. Patrick the Weather Forecaster

- Now, Patrick is a *weather* forecaster
- A weather forecaster's task is sequential
- each day, he forecasts for tomorrow, drawing on his accumulated experience all that has passed up to today

## 2b. Patrick the Weather Forecaster





## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- Forecasts made sequentially on days  $0, 1, 2, \dots$ , each refer to events that will become known the next day

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- Forecasts made sequentially on days  $0, 1, 2, \dots$ , each refer to events that will become known the next day
- Denote  $\mathcal{B}_i$  the totality of events known to Patrick on day  $i$ , so  $\mathcal{B}_0 \subseteq \mathcal{B}_1 \subseteq \dots$

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- Forecasts made sequentially on days  $0, 1, 2, \dots$ , each refer to events that will become known the next day
- Denote  $\mathcal{B}_i$  the totality of events known to Patrick on day  $i$ , so  $\mathcal{B}_0 \subseteq \mathcal{B}_1 \subseteq \dots$
- Each  $\mathcal{B}_i$  is sigma algebra, or a collection of measurable sets

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- Forecasts made sequentially on days  $0, 1, 2, \dots$ , each refer to events that will become known the next day
- Denote  $\mathcal{B}_i$  the totality of events known to Patrick on day  $i$ , so  $\mathcal{B}_0 \subseteq \mathcal{B}_1 \subseteq \dots$
- Each  $\mathcal{B}_i$  is sigma algebra, or a collection of measurable sets
- Patrick's arbitrary subjective probability distribution  $\Pi$  defined over  $\cup_{i=0}^{\infty} \mathcal{B}_i$

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- Forecasts made sequentially on days  $0, 1, 2, \dots$ , each refer to events that will become known the next day
- Denote  $\mathcal{B}_i$  the totality of events known to Patrick on day  $i$ , so  $\mathcal{B}_0 \subseteq \mathcal{B}_1 \subseteq \dots$
- Each  $\mathcal{B}_i$  is sigma algebra, or a collection of measurable sets
- Patrick's arbitrary subjective probability distribution  $\Pi$  defined over  $\cup_{i=0}^{\infty} \mathcal{B}_i$
- Patrick's forecasts made on day  $i$  are for events in  $\mathcal{B}_{i+1}$ , and are calculated from his current subjective conditional distribution  $\Pi(\cdot | \mathcal{B}_i)$

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- $X_i \in \mathcal{B}_i$  is the maximum temperature in Philly on day  $i$

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- $X_i \in \mathcal{B}_i$  is the maximum temperature in Philly on day  $i$
- $m_i$  is the median of Patrick's distribution for  $X_i$  as assessed on day  $i - 1$

## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- $X_i \in \mathcal{B}_i$  is the maximum temperature in Philly on day  $i$
- $m_i$  is the median of Patrick's distribution for  $X_i$  as assessed on day  $i - 1$
- $S_i = \{X_i > m_i\}$



## 2b. Patrick, the Weather Forecaster, Loves Measure Theory

- $X_i \in \mathcal{B}_i$  is the maximum temperature in Philly on day  $i$
- $m_i$  is the median of Patrick's distribution for  $X_i$  as assessed on day  $i - 1$
- $S_i = \{X_i > m_i\}$
- By def.,  $\Pi(S_i | \mathcal{B}_{i-1}) = \frac{1}{2}$

## 2b. Patrick the Weather Forecaster

- Again, we appear to have a conflict if over many days, say 80%, the  $(X_i)$  exceed their assessed medians

## 2b. Patrick the Weather Forecaster

- Again, we appear to have a conflict if over many days, say 80%, the  $(X_i)$  exceed their assessed medians
- Possibility 1: Patrick's subjective prob. dist.  $\Pi$  is not coherent

## 2b. Patrick the Weather Forecaster

- Again, we appear to have a conflict if over many days, say 80%, the  $(X_i)$  exceed their assessed medians
- Possibility 1: Patrick's subjective prob. dist.  $\Pi$  is not coherent
- Possibility 2: If  $\Pi$  is coherent, then Patrick is miscalibrated

### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$

### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$
- Each day  $i$  is associated with some event  $S_i \in \mathcal{B}_i$ , for example, the max temp in Philly on day  $i$

### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$
- Each day  $i$  is associated with some event  $S_i \in \mathcal{B}_i$ , for example, the max temp in Philly on day  $i$
- $Y_i = \mathbb{1}\{S_i\}$

### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$
- Each day  $i$  is associated with some event  $S_i \in \mathcal{B}_i$ , for example, the max temp in Philly on day  $i$
- $Y_i = \mathbb{1}\{S_i\}$
- $\tilde{Y}_i = \Pi(S_i|\mathcal{B}_{i-1}) = \mathbb{E}(Y_i|\mathcal{B}_{i-1})$



### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$
- Each day  $i$  is associated with some event  $S_i \in \mathcal{B}_i$ , for example, the max temp in Philly on day  $i$
- $Y_i = \mathbb{1}\{S_i\}$
- $\tilde{Y}_i = \Pi(S_i|\mathcal{B}_{i-1}) = \mathbb{E}(Y_i|\mathcal{B}_{i-1})$
- $\xi_i = \mathbb{1}\{\text{day } i \text{ is included in the test set}\}$

### 3. General Calibration Theorem

- Setting: Patrick is a weather forecaster who makes sequential forecasts according to  $\Pi$
- Each day  $i$  is associated with some event  $S_i \in \mathcal{B}_i$ , for example, the max temp in Philly on day  $i$
- $Y_i = \mathbb{1}\{S_i\}$
- $\tilde{Y}_i = \Pi(S_i|\mathcal{B}_{i-1}) = \mathbb{E}(Y_i|\mathcal{B}_{i-1})$
- $\xi_i = \mathbb{1}\{\text{day } i \text{ is included in the test set}\}$
- *admissible*: determine the test set of days  $(\xi_i)$  sequentially, so  $\xi_i \in \mathcal{B}_{i-1}$

### 3. General Calibration Theorem

- Number of test days

$$\nu_k = \sum_{i=1}^k \xi_i$$

### 3. General Calibration Theorem

- Number of test days

$$\nu_k = \sum_{i=1}^k \xi_i$$

- Proportion of test days in which the event  $S_i$  occurs

$$p_k = \frac{1}{\nu_k} \sum_{i=1}^k \xi_i Y_i$$

### 3. General Calibration Theorem

- Number of test days

$$\nu_k = \sum_{i=1}^k \xi_i$$

- Proportion of test days in which the event  $S_i$  occurs

$$p_k = \frac{1}{\nu_k} \sum_{i=1}^k \xi_i Y_i$$

- Proportion of test days in which Patrick predicts  $S_i$  to occur, or the average forecast probability

$$\pi_k = \frac{1}{\nu_k} \sum_{i=1}^k \xi_i \tilde{Y}_i$$

### 3. General Calibration Theorem

- Theorem. Assume  $\Pi$  is coherent and  $(\xi_i)$  is admissible. Then

$\Pi$ -almost-surely, if  $\nu_k \rightarrow \infty$ , then  $p_k - \pi_k \rightarrow 0$ .

### 3. General Calibration Theorem

- Theorem. Assume  $\Pi$  is coherent and  $(\xi_i)$  is admissible. Then

$\Pi$ -almost-surely, if  $\nu_k \rightarrow \infty$ , then  $p_k - \pi_k \rightarrow 0$ .

- Proof: martingale convergence

### 3. General Calibration Theorem

- Theorem. Assume  $\Pi$  is coherent and  $(\xi_i)$  is admissible. Then

$\Pi$ -almost-surely, if  $\nu_k \rightarrow \infty$ , then  $p_k - \pi_k \rightarrow 0$ .

- Proof: martingale convergence
- *A coherent Bayesian feels almost certain he is well calibrated*



### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$
- Because  $\frac{1}{\nu_i}, \xi_i, \tilde{Y}_i \in \mathcal{B}_{i-1}$ , we have  $\mathbb{E}(X_i | \mathcal{B}_{i-1}) = 0$

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$
- Because  $\frac{1}{\nu_i}, \xi_i, \tilde{Y}_i \in \mathcal{B}_{i-1}$ , we have  $\mathbb{E}(X_i | \mathcal{B}_{i-1}) = 0$
- Then  $U_k = \sum_{i=1}^k X_i \in \mathcal{B}_k$  is a martingale

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$
- Because  $\frac{1}{\nu_i}, \xi_i, \tilde{Y}_i \in \mathcal{B}_{i-1}$ , we have  $\mathbb{E}(X_i | \mathcal{B}_{i-1}) = 0$
- Then  $U_k = \sum_{i=1}^k X_i \in \mathcal{B}_k$  is a martingale
- $(U_k)$  is a bounded martingale because

$$\mathbb{E}(U_k^2) = \sum_{i=1}^k \mathbb{E}(X_i^2) \leq \frac{1}{4} \mathbb{E} \left[ \sum_{i=1}^k \left( \frac{1}{\nu_i} \xi_i \right)^2 \right] \leq \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{24}$$

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$
- Because  $\frac{1}{\nu_i}, \xi_i, \tilde{Y}_i \in \mathcal{B}_{i-1}$ , we have  $\mathbb{E}(X_i | \mathcal{B}_{i-1}) = 0$
- Then  $U_k = \sum_{i=1}^k X_i \in \mathcal{B}_k$  is a martingale
- $(U_k)$  is a bounded martingale because

$$\mathbb{E}(U_k^2) = \sum_{i=1}^k \mathbb{E}(X_i^2) \leq \frac{1}{4} \mathbb{E} \left[ \sum_{i=1}^k \left( \frac{1}{\nu_i} \xi_i \right)^2 \right] \leq \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{24}$$

- So by Martingale Convergence Thm,  $(U_k)$  converges  $\Pi$  a.s.,

$$U_k = \sum_{i=1}^k X_i = \sum_{i=1}^k \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i) \xrightarrow{\Pi \text{ a.s.}} s \in \mathbb{R}$$

### 3. Proof of General Calibration Theorem

- Let  $X_i = \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i)$
- Recall  $\tilde{Y}_i = \mathbb{E}(Y_i | \mathcal{B}_{i-1}) = \Pi(S_i | \mathcal{B}_{i-1})$
- Because  $\frac{1}{\nu_i}, \xi_i, \tilde{Y}_i \in \mathcal{B}_{i-1}$ , we have  $\mathbb{E}(X_i | \mathcal{B}_{i-1}) = 0$
- Then  $U_k = \sum_{i=1}^k X_i \in \mathcal{B}_k$  is a martingale
- $(U_k)$  is a bounded martingale because

$$\mathbb{E}(U_k^2) = \sum_{i=1}^k \mathbb{E}(X_i^2) \leq \frac{1}{4} \mathbb{E} \left[ \sum_{i=1}^k \left( \frac{1}{\nu_i} \xi_i \right)^2 \right] \leq \frac{1}{4} \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{24}$$

- So by Martingale Convergence Thm,  $(U_k)$  converges  $\Pi$  a.s.,

$$U_k = \sum_{i=1}^k X_i = \sum_{i=1}^k \frac{1}{\nu_i} \xi_i (Y_i - \tilde{Y}_i) \xrightarrow{\Pi \text{ a.s.}} s \in \mathbb{R}$$

- By Kronecker's Lemma,  $\Pi$  a.s. we have

$$p_k - \pi_k = \frac{1}{\nu_k} \sum_{i=1}^k \xi_i (Y_i - \tilde{Y}_i) \xrightarrow{\Pi \text{ a.s.}} 0$$

### 3. Kronecker's Lemma

- Kronecker's Lemma If  $x_n$  is a sequence of real numbers with

$$\sum_{i=1}^{\infty} x_i = s \in \mathbb{R},$$

then for any increasing sequence  $(b_n)$  with  $b_n \rightarrow \infty$ ,

$$\lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{i=1}^k b_i x_i = 0.$$



### 3. Kronecker's Lemma

- Kronecker's Lemma If  $x_n$  is a sequence of real numbers with

$$\sum_{i=1}^{\infty} x_i = s \in \mathbb{R},$$

then for any increasing sequence  $(b_n)$  with  $b_n \rightarrow \infty$ ,

$$\lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{i=1}^k b_i x_i = 0.$$

- Idea: look at the partial sums

### 3. Kronecker's Lemma

- Kronecker's Lemma If  $x_n$  is a sequence of real numbers with

$$\sum_{i=1}^{\infty} x_i = s \in \mathbb{R},$$

then for any increasing sequence  $(b_n)$  with  $b_n \rightarrow \infty$ ,

$$\lim_{k \rightarrow \infty} \frac{1}{b_k} \sum_{i=1}^k b_i x_i = 0.$$

- Idea: look at the partial sums
- Here,  $\frac{1}{\nu_k} \leq \frac{1}{\nu_i}$

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*
- For example, look at all days for which the forecast of rain was close to  $\omega$ , find the long run proportion  $p$  of such days in which rain actually occurred, and plot  $p$  against  $\omega$

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*
- For example, look at all days for which the forecast of rain was close to  $\omega$ , find the long run proportion  $p$  of such days in which rain actually occurred, and plot  $p$  against  $\omega$
- The diagonal  $p = \omega$  indicates perfect empirical calibration

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*
- For example, look at all days for which the forecast of rain was close to  $\omega$ , find the long run proportion  $p$  of such days in which rain actually occurred, and plot  $p$  against  $\omega$
- The diagonal  $p = \omega$  indicates perfect empirical calibration
- If the curve indicates poor calibration, try finding a *continuous* transformation  $\gamma$  to transform the curve into the diagonal



## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*
- For example, look at all days for which the forecast of rain was close to  $\omega$ , find the long run proportion  $p$  of such days in which rain actually occurred, and plot  $p$  against  $\omega$
- The diagonal  $p = \omega$  indicates perfect empirical calibration
- If the curve indicates poor calibration, try finding a *continuous* transformation  $\gamma$  to transform the curve into the diagonal
- In other words, recalibrate  $\omega$  to  $p = \gamma(\omega)$

## 5. How to Achieve Calibration? Recalibration?

- How might Patrick achieve the calibration he almost surely expects from the theorem?
- Try keeping a running tally of assessments and outcomes
- Summarize this tally in a *calibration curve*
- For example, look at all days for which the forecast of rain was close to  $\omega$ , find the long run proportion  $p$  of such days in which rain actually occurred, and plot  $p$  against  $\omega$
- The diagonal  $p = \omega$  indicates perfect empirical calibration
- If the curve indicates poor calibration, try finding a *continuous* transformation  $\gamma$  to transform the curve into the diagonal
- In other words, recalibrate  $\omega$  to  $p = \gamma(\omega)$
- Can Patrick learn from his inadequacies as a forecaster, and use this knowledge to improve future assessments?

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*
- Assume Patrick is coherent. Let  $A_1, A_2$  be disjoint events having  $\Pi$  probability  $\omega_1, \omega_2$ . Since  $\gamma$  is a probability,

$$\gamma(A_1 \cup A_2; \omega_1 + \omega_2) = \gamma(A_1; \omega_1) + \gamma(A_2; \omega_2)$$

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*
- Assume Patrick is coherent. Let  $A_1, A_2$  be disjoint events having  $\Pi$  probability  $\omega_1, \omega_2$ . Since  $\gamma$  is a probability,

$$\gamma(A_1 \cup A_2; \omega_1 + \omega_2) = \gamma(A_1; \omega_1) + \gamma(A_2; \omega_2)$$

- Then by continuity of  $\gamma$ ,  $\exists c, \forall 0 \leq \omega \leq 1$ ,

$$\gamma(A; \omega) = c\omega$$

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*
- Assume Patrick is coherent. Let  $A_1, A_2$  be disjoint events having  $\Pi$  probability  $\omega_1, \omega_2$ . Since  $\gamma$  is a probability,

$$\gamma(A_1 \cup A_2; \omega_1 + \omega_2) = \gamma(A_1; \omega_1) + \gamma(A_2; \omega_2)$$

- Then by continuity of  $\gamma$ ,  $\exists c, \forall 0 \leq \omega \leq 1$ ,

$$\gamma(A; \omega) = c\omega$$

- Details? I can see how  $\gamma$  can't be  $c\omega^k$ ,  $k \neq 1$



## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*
- Assume Patrick is coherent. Let  $A_1, A_2$  be disjoint events having  $\Pi$  probability  $\omega_1, \omega_2$ . Since  $\gamma$  is a probability,

$$\gamma(A_1 \cup A_2; \omega_1 + \omega_2) = \gamma(A_1; \omega_1) + \gamma(A_2; \omega_2)$$

- Then by continuity of  $\gamma$ ,  $\exists c, \forall 0 \leq \omega \leq 1$ ,

$$\gamma(A; \omega) = c\omega$$

- Details? I can see how  $\gamma$  can't be  $c\omega^k$ ,  $k \neq 1$
- $c = 1$  because  $\gamma(\Omega; 1) = 1 = c \cdot 1$ , so

$$\gamma(A; \omega) = \omega$$

## 5. How to Achieve Calibration? Recalibration?

- Paradox: How can Patrick simultaneously assign 2 different probabilities  $\omega$  and  $p = \gamma(\omega)$  to one event?
- He can't...
- Theorem *If  $\omega$  and  $\gamma$  are both coherent, then they are identical*
- Assume Patrick is coherent. Let  $A_1, A_2$  be disjoint events having  $\Pi$  probability  $\omega_1, \omega_2$ . Since  $\gamma$  is a probability,

$$\gamma(A_1 \cup A_2; \omega_1 + \omega_2) = \gamma(A_1; \omega_1) + \gamma(A_2; \omega_2)$$

- Then by continuity of  $\gamma$ ,  $\exists c, \forall 0 \leq \omega \leq 1$ ,

$$\gamma(A; \omega) = c\omega$$

- Details? I can see how  $\gamma$  can't be  $c\omega^k$ ,  $k \neq 1$
- $c = 1$  because  $\gamma(\Omega; 1) = 1 = c \cdot 1$ , so

$$\gamma(A; \omega) = \omega$$

- Corollary: *Patrick can't coherently recalibrate his forecasts!*

## 5. How to Achieve Calibration? Recalibration?

- Since a calibration curve represents a  $\gamma$  function, it cannot be used to “correct” your assessments

## 5. How to Achieve Calibration? Recalibration?

- Since a calibration curve represents a  $\gamma$  function, it cannot be used to “correct” your assessments
- Indeed, we know no valid model for how people should change their opinions when assessments they believe are coherent systematically deviate from the observed frequencies of events

## 5. How to Achieve Calibration? Recalibration?

- Since a calibration curve represents a  $\gamma$  function, it cannot be used to “correct” your assessments
- Indeed, we know no valid model for how people should change their opinions when assessments they believe are coherent systematically deviate from the observed frequencies of events
- We speculate that the fault lies in the idea that  $\gamma$  must depend on  $\omega$  alone

## 5. How to Achieve Calibration? Recalibration?

- Since a calibration curve represents a  $\gamma$  function, it cannot be used to “correct” your assessments
- Indeed, we know no valid model for how people should change their opinions when assessments they believe are coherent systematically deviate from the observed frequencies of events
- We speculate that the fault lies in the idea that  $\gamma$  must depend on  $\omega$  alone
- Rather, what people should learn from outcome feedback is something about properties of the world

## 5. How to Achieve Calibration? Recalibration?

- Under this view, the innocent-sounding assumptions of the General Theorem - that you know the outcomes of all previous events before you state your probability of the next event - is a *psychologically strong requirement*

## 5. How to Achieve Calibration? Recalibration?

- Under this view, the innocent-sounding assumptions of the General Theorem - that you know the outcomes of all previous events before you state your probability of the next event - is a *psychologically strong requirement*
- Not only must you have perfect memory for past outcomes, but you must also be able to use outcome information to develop a better understanding of the world