

X857/75/02

Physics Section 1 — Questions

WEDNESDAY, 15 MAY 1:00 PM – 3:30 PM

Instructions for the completion of Section 1 are given on *page 02* of your question and answer booklet X857/75/01.

Record your answers on the answer grid on page 03 of your question and answer booklet.

Reference may be made to the Data sheet on *page 02* of this booklet and to the Relationships sheet X857/75/11.

Before leaving the examination room you must give your question and answer booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





Speed of light in materials

Material	Speed in m s ⁻¹	
Air	3.0×10^8	
Carbon dioxide	3.0×10^8	
Diamond	1.2×10^8	
Glass	$2 \cdot 0 \times 10^8$	
Glycerol	2·1 × 10 ⁸	
Water	$2\cdot3\times10^8$	

Gravitational field strengths

	Gravitational field strength on the surface in N kg ⁻¹
Earth	9.8
Jupiter	23
Mars	3.7
Mercury	3.7
Moon	1.6
Neptune	11
Saturn	9.0
Sun	270
Uranus	8.7
Venus	8.9

Specific latent heat of fusion of materials

Material	Specific latent heat of fusion in Jkg ⁻¹
Alcohol	0.99 × 10 ⁵
Aluminium	3.95×10^5
Carbon Dioxide	1.80×10^{5}
Copper	$2 \cdot 05 \times 10^5$
Iron	$2 \cdot 67 \times 10^5$
Lead	0.25×10^5
Water	$3 \cdot 34 \times 10^5$

Specific latent heat of vaporisation of materials

<u> </u>		
Material	Specific latent heat of vaporisation in J kg ⁻¹	
Alcohol	11·2 × 10 ⁵	
Carbon Dioxide	3.77×10^5	
Glycerol	$8 \cdot 30 \times 10^5$	
Turpentine	$2\cdot 90\times 10^5$	
Water	22.6×10^5	

Speed of sound in materials

Material	Speed in m s ⁻¹
Aluminium	5200
Air	340
Bone	4100
Carbon dioxide	270
Glycerol	1900
Muscle	1600
Steel	5200
Tissue	1500
Water	1500

Specific heat capacity of materials

Material	Specific heat capacity in J kg ⁻¹ °C ⁻¹
Alcohol	2350
Aluminium	902
Copper	386
Glass	500
Ice	2100
Iron	480
Lead	128
Oil	2130
Water	4180

Melting and boiling points of materials

Material	Melting point in °C	Boiling point in °C
Alcohol	-98	65
Aluminium	660	2470
Copper	1077	2567
Lead	328	1737
Iron	1537	2737
Water	_	100

Radiation weighting factors

3 3,		
Type of radiation	Radiation weighting factor	
alpha	20	
beta	1	
fast neutrons	10	
gamma	1	
slow neutrons	3	
X-rays	1	

SECTION 1


Attempt ALL questions

- 1. Which of the following are both vectors?
 - A weight and acceleration
 - B kinetic energy and acceleration
 - C mass and acceleration
 - D force and speed
 - E speed and acceleration
- 2. A car is travelling at $6.0 \,\mathrm{m\,s^{-1}}$ along a straight level road.

The car then accelerates uniformly at $2.0 \, \text{m s}^{-2}$ for $4.0 \, \text{s}$.

The final speed of the car is


- A $8.0 \,\mathrm{m\,s^{-1}}$
- B $14 \,\mathrm{m \, s^{-1}}$
- C $22 \,\mathrm{m \, s^{-1}}$
- D $26 \,\mathrm{m \, s^{-1}}$
- E $48 \,\mathrm{m \, s^{-1}}$.
- **3.** The graph shows how the speed v of a car varies with time t.



During which part of the journey does the car have the greatest acceleration?

- A OP
- B PQ
- C QR
- D RS
- E ST

4. A ball is kicked horizontally off a high river bank as shown.



The ball lands on the lower river bank at X, $2.0 \, \text{s}$ after the ball is kicked.

The river is $3.0 \,\mathrm{m}$ wide.

The effect of air resistance on the ball is negligible.

The distance d between the edge of the lower river bank and ${\sf X}$ is

- A 1.0 m
- B 4⋅0 m
- C 13 m
- D 16 m
- E 19 m.

5. The table gives the distance from Earth, the approximate surface temperature and the age of five stars.

Star	Distance from Earth (light-years)	Approximate surface temperature (K)	Age (years)
Sirius A	8.6	9900	2·4 × 10 ⁸
Polaris	430	6000	$7 \cdot 0 \times 10^7$
Betelgeuse	640	3600	7·9 × 10 ⁶
Rigel	860	11 000	8·0 × 10 ⁶
VY Canis Majoris	3900	3500	1·0 × 10 ⁷

A student makes the following statements based on this information.

- I As the distance from Earth increases, the age of a star decreases.
- II As the age of a star increases, the approximate surface temperature of the star increases.
- III There is no apparent relationship between the distance from Earth and the approximate surface temperature of a star.

Which of these statements is/are correct?

- A I only
- B II only
- C III only
- D I and III only
- E I, II and III

6. A geostationary satellite orbits the Earth.

Which row in the table shows the altitude above the surface of the Earth and orbital period of the geostationary satellite?

	Altitude above the surface of the Earth (km)	Orbital period (hours)
Α	36 000	12
В	36 000	24
С	36 000	48
D	18 000	12
Е	18 000	24

7. The weight of a robot on Earth is 240 N.

The weight of the robot on Mars is


- A 3.7 N
- B 65 N
- C 91 N
- D 240 N
- E 890 N.
- **8.** A hairdryer is connected to a 230 V supply.

The current in the hairdryer is $2.0 \,\mathrm{A}$.

The electrical charge that passes through the hairdryer in 5 minutes is

- A 10 C
- B 460 C
- C 600 C
- D 1150 C
- E 69 000 C.

9. The graph shows how the resistance R of a thermistor varies with temperature T.




The thermistor is connected in a circuit.


At a temperature of 50 °C the current in the thermistor is $0.004\,A$.


At this temperature the voltage across the thermistor is


- A 0.00002V
- B 0.002 V
- C 0.008 V
- D 8 V
- E 500 V.


10. A student sets up the circuits shown. In which circuit will both LEDs be lit?












11. A circuit is set up as shown.




The room temperature is 20 °C.

The lamp is off.

The lamp will light when

- A the light level is decreased below a certain value
- B the light level is increased above a certain value
- C the resistance of R is increased above a certain value
- D the battery voltage is reduced to 5 V
- E the temperature is increased above a certain value.

12. A circuit is set up as shown.



A student makes the following statements about the readings on the voltmeters.


- $I \qquad V_1 = V_2$
- $|| V_2 = V_3$
- $III \qquad V_S = V_1 + V_2$

Which of these statements must always be true?

- A II only
- B I and II only
- C I and III only
- D II and III only
- E I, II and III

13. A solid substance is placed in an insulated container and heated.

The graph shows how the temperature T of the substance varies with time t.



To calculate the specific latent heat of fusion of the substance a student would use the time from section

- A PQ
- B QR
- C RS
- D ST
- E TU.
- **14.** The pressure p due to a liquid at a depth h is given by the relationship

$$p = \rho g h$$

where ρ is the density of the liquid and g is the gravitational field strength.

A liquid has a density of $990 \, \text{kg m}^{-3}$.

When the pressure due to the liquid is 1470 Pa, the depth in the liquid is

- A 0.069 m
- B 0.15 m
- C 0.67 m
- D 1.5 m
- E 6.6 m.

15. A car is parked in the sun for some time. During this time the air pressure inside the tyres increases.

The reason for this increase in pressure is

- A the volume occupied by the air particles in the tyres has increased
- B the force produced by the air particles in the tyres acts over a smaller area
- C the average spacing between the air particles in the tyres has increased
- D the increased temperature has made the air particles in the tyres expand
- E the air particles in the tyres are moving with greater kinetic energy.
- **16.** The temperature of a sample of gas in a container is 20 °C.

The volume of the gas is $0.30 \,\mathrm{m}^3$.


The container is free to expand in order to maintain a constant pressure.

The temperature of the gas is increased to 50 °C.

The volume now occupied by the gas is

- A $0.12 \,\mathrm{m}^3$
- B $0.27 \,\mathrm{m}^3$
- C $0.30 \, \text{m}^3$
- D $0.33 \,\mathrm{m}^3$
- E $0.75 \,\mathrm{m}^3$.

17. The following diagram gives information about a wave.




Which row in the table shows the amplitude and wavelength of the wave?

	Amplitude (m)	Wavelength (m)
Α	3	4
В	3	8
С	6	4
D	6	8
Е	8	3


18. A student is studying waves with a period of $80.0 \, \text{ms}$ and a wavelength of $4.00 \, \text{m}$. The frequency of these waves is

- A 0.0125 Hz
- B 0⋅320 Hz
- C 12.5 Hz
- D 80.0 Hz
- E 320 Hz.

19. Which of the following diagrams shows the diffraction of water waves as they pass between two walls?



20. A ray of red light passes through a glass block as shown.



Which row in the table shows the angle of incidence and the corresponding angle of refraction at point X?

	Angle of incidence	Angle of refraction
Α	35°	60°
В	30°	55°
С	35°	30°
D	55°	30°
E	60°	35°

21. Which row in the table shows the paths taken by alpha particles and gamma radiation as they pass through a uniform electric field between two metal plates?

	Path taken by alpha particles	Path taken by gamma radiation
A	+	†
В		÷ •
С	+ 0	
D	+ 0	+ 0
E	+ 0	

22. For a particular radioactive source, 1800 atoms decay in a time of 3 minutes.

The activity of the source is

- A 10 Bq
- B 600 Bq
- C 1800 Bq
- D 5400 Bq
- E 324 000 Bq.
- 23. The crew on an aircraft during a transatlantic flight are exposed to cosmic radiation at an equivalent dose rate of $5.0 \,\mu\text{Sy} \, h^{-1}$.

The crew complete 6 transatlantic flights each month. The average duration of a flight is 8 hours.

The equivalent dose received by the crew due to cosmic radiation during transatlantic flights in **one year** is

- A $30 \,\mu\text{Sv}$
- B $40 \,\mu\text{Sv}$
- C 60 μSv
- D 240 μSv
- E 2880 μSv.
- **24.** A radioactive tracer is injected into a patient to enable doctors to check the function of a patient's kidneys.

Radiation from the tracer is monitored outside the patient's body by a detector.

Which row in the table shows the most suitable type of radiation emitted and the half-life for the tracer?

	Type of radiation emitted	Half-life of tracer
Α	alpha	6 hours
В	beta	6 hours
С	beta	6 years
D	gamma	6 hours
Е	gamma	6 years

[Turn over for next question

- **25.** The activity of a radioactive source is 56 MBq.
 - The activity of the source 40 hours later is $3.5 \, MBq$.

The half-life of this source is

- A 8 hours
- B 10 hours
- C 16 hours
- D 20 hours
- E 28 hours.

[END OF SECTION 1. NOW ATTEMPT THE QUESTIONS IN SECTION 2 OF YOUR QUESTION AND ANSWER BOOKLET]

[BLANK PAGE]

DO NOT WRITE ON THIS PAGE

[BLANK PAGE]

DO NOT WRITE ON THIS PAGE

National Qualifications 2019

X857/75/01

Physics
Section 1 — Answer grid
and Section 2

WEDNESDAY, 15 MAY 1:00 PM – 3:30 PM



Mark

ull name of cen	tre			Town	
orename(s)		Sur	name		Number of seat
Date of birtl	h Month	Year	Scottish ca	andidate number	

Total marks — 135

SECTION 1 — 25 marks

Attempt ALL questions.

Instructions for completion of Section 1 are given on page 02.

SECTION 2 — 110 marks

Attempt ALL questions.

Reference may be made to the Data sheet on *page 02* of the question paper X857/75/02 and to the Relationships sheet X857/75/11.

Write your answers clearly in the spaces provided in this booklet. Additional space for answers and rough work is provided at the end of this booklet. If you use this space you must clearly identify the question number you are attempting. Any rough work must be written in this booklet. Score through your rough work when you have written your final copy.

Use blue or black ink.

Before leaving the examination room you must give this booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





The questions for Section 1 are contained in the question paper X857/75/02.

Read these and record your answers on the answer grid on page 03 opposite.

Use blue or black ink. Do NOT use gel pens or pencil.

- 1. The answer to each question is **either** A, B, C, D or E. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is **only one correct** answer to each question.
- 3. Any rough work must be written in the additional space for answers and rough work at the end of this booklet.

Sample question

The energy unit measured by the electricity meter in your home is the

- A ampere
- B kilowatt-hour
- C watt
- D coulomb
- E volt.

The correct answer is ${\bf B}$ — kilowatt-hour. The answer ${\bf B}$ bubble has been clearly filled in (see below).

Α	В	С	D	Ε
0		0	0	0

Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.

Α	В	С	D	Ε
0		0		0

If you then decide to change back to an answer you have already scored out, put a tick (\checkmark) to the right of the answer you want, as shown below:





	Α	В	С	D	E
1	0	0	0	0	0
2	0	0	0	0	0
3	0	0	0	0	0
4	0	0	0	0	0
5	0	0	0	0	0
6	\circ	\circ	\circ	\circ	\circ
7	0	0	0	0	0
8	\circ	0	\circ	0	\circ
9	0	0	0	0	0
10	0	0	0	0	0
11	0	0	0	0	0
12	0	0	0	0	0
13	0	0	0	0	0
14	0	0	0	0	0
15	0	0	0	0	0
16	0	0	0	0	0
17	0	0	0	0	0
18	0	0	0	0	0
19	0	0	0	0	0
20	0	0	0	0	0
21	0	0	0	0	0
22	0	0	0	0	0
23	0	0	0	0	0
24	0	0	0	0	0
25	0	0	0	0	0

* Y 8 5 7 7 5 0 1 0 3 *

page 03

[BLANK PAGE]

DO NOT WRITE ON THIS PAGE

page 04

[Turn over for SECTION 2

DO NOT WRITE ON THIS PAGE


page 05

SECTION 2 — 110 marks Attempt ALL questions

1. A quadcopter is a drone with four rotating blades.



(a) In a race, the quadcopter is flown along a route from point A to point E.



1. (a) (continued)

(i) By scale drawing or otherwise, determine the magnitude of the resultant displacement of the quadcopter from point A to point E.
 Space for working and answer

(ii) By scale drawing or otherwise, determine the direction of the resultant displacement of the quadcopter from point A to point E.Space for working and answer



1. (continued)

(b) The quadcopter takes 32·5 s to complete the race.
 Determine the average velocity of the quadcopter over the whole race.
 Space for working and answer

(c) A second quadcopter completes the race at an average speed of $1 \cdot 25 \,\mathrm{m\,s^{-1}}$.

The distance travelled by this quadcopter during the race is $37 \cdot 0 \,\mathrm{m}$.

Determine the **difference** in the times taken by the quadcopters to complete the race.

3

Space for working and answer

MARKS DO NOT WRITE IN THIS MARGIN

1. (continued)

(d) After passing point E, the quadcopter hovers at a constant height.


Describe how the overall lift force provided by the four rotating blades compares to the weight of the quadcopter.



2. A glider is accelerated from rest by a cable attached to a winch.



The graph shows the horizontal velocity v_h of the glider for the first 20 s of its motion.



- (a) The glider is accelerated by a constant unbalanced force of 925 N.
 - (i) Show that the initial acceleration of the glider is 2·5 m s⁻². **2**Space for working and answer

(ii) Calculate the mass of the glider.


Space for working and answer

3

1

2. (a) (continued)

(iii) At 2.0 s the cable pulls the glider with a force of 1200 N.




(A) Determine the size of the frictional forces acting on the glider at this time.

(B) Suggest one design feature of the glider that reduces the frictional forces acting on it.

(b) At 8·0 s the glider reaches its take-off speed and leaves the ground.
 Determine the distance the glider travels along the ground before take-off.
 3
 Space for working and answer



In 1971, the astronaut Alan Shepard hit a golf ball on the surface of the Moon.



Using your knowledge of physics, comment on the similarities and/or differences between this event and hitting an identical ball on the surface of the Earth.

3

page 12


3. (continued)

DO NOT WRITE IN THIS MARGIN

page 13

4. Astronomers studying a distant star analyse the light from the star that reaches Earth.

The line spectrum from the star is shown, along with the line spectra of the elements hydrogen, helium, mercury, calcium, and sodium.



(a) Determine which of these elements are present in the star.

4. (continued)

- (b) The star is 97 light-years from Earth.
 - (i) State what is meant by the term light-year.

1

(ii) Calculate the distance, in metres, from the star to Earth.

Space for working and answer

3

- (c) Astronomers use satellite-based telescopes to collect information about objects in space.
 - (i) Suggest an advantage of using satellite-based telescopes such as the Hubble Space Telescope.

1

(ii) State one other use of satellites.

1




3

- **5.** A student is investigating how the length of a wire affects its resistance. The student connects different lengths of wire to a power supply of fixed voltage and measures the current in each length of wire.
 - (a) The measurements taken by the student are shown in the table.

Length of wire (m)	Current (A)
0-20	0.94
0.40	0.66
0-60	0.47
0.80	0.37
1.00	0.32

(i) Using the graph paper, draw a graph of these measurements. (Additional graph paper, if required, can be found on *page 38*)



5. (a) (continued)

(ii) State whether the resistance of the wire increases, decreases or stays the same, as the length of wire increases.Justify your answer.

2

(iii) Use your graph to predict the current in a $0.50\,\mathrm{m}$ length of wire, when connected to the power supply.

1

(iv) Suggest one way in which the experimental procedure could be improved to give more reliable results.


1

[Turn over

5. (continued)

(b) A length of the wire with a resistance of $5.2\,\Omega$ is then folded into a rectangular shape and the ends are joined together.

An ohmmeter is connected across the wire between point X and point Y as shown.



State whether the reading on the ohmmeter would be less than, equal to or greater than $5 \cdot 2 \Omega$.

You must justify your answer.



page 18


[Turn over for next question

DO NOT WRITE ON THIS PAGE



page 19

- **6.** A student is investigating connecting different combinations of resistors in circuits.
 - (a) The student sets up a circuit as shown.



(i) Calculate the current in the circuit.


Space for working and answer

4

(ii) Calculate the power dissipated in the 120 Ω resistor. Space for working and answer

6. (continued)

(b) The student then sets up a different circuit as shown.



(i) Determine the total resistance of this circuit.

Space for working and answer

4

(ii) State how the power dissipated in the 120 Ω resistor in this circuit compares to the power dissipated in the 120 Ω resistor in the circuit in part (a) (ii).

Justify your answer.



page 21

7. A hot water dispenser is used to heat enough water for one cup at a time.



The rating plate for the hot water dispenser is shown.

Model: 1-KUPPA 3·5 kW 230 V 50 Hz

The hot water dispenser takes 26 s to heat enough water for one cup.

(a) Show that the energy supplied to the hot water dispenser during this time is $91\,000\,\mathrm{J}$.

Space for working and answer

(continued)

- (b) The hot water dispenser heats $0.250 \, \text{kg}$ of water for each cup.
 - (i) Calculate the minimum energy required to heat 0.250 kg of water from an initial temperature of 20.0 °C to its boiling point.

3

Space for working and answer

(ii) As the water is dispensed into the cup, steam is released.

Determine the maximum mass of steam that can be produced while the water for one cup is being heated.


Space for working and answer

(iii) Explain why, in practice, the mass of steam produced is less than calculated in (b)(ii).



2

8. A water rocket consists of a plastic bottle partly filled with water. Air is pumped in through the water. When the pressure is great enough, the tube detaches from the bottle. Water is forced out of the bottle, which causes the bottle to be launched upwards.



At launch, the air in the bottle is at a pressure of $1.74 \times 10^5 \, Pa$.

(a) On the diagram below, show all the forces acting vertically on the bottle as it is launched.

You must name these forces and show their directions.

(An additional diagram, if required, can be found on page 39)



8. (continued)

(b) The area of water in contact with the pressurised air in the bottle is $4.50 \times 10^{-3} \, m^2$.

Calculate the force exerted on the water by the pressurised air at launch. 3

Space for working and answer

(c) At launch, the air in the bottle has a volume of $7.5 \times 10^{-4} \, \text{m}^3$.

At one point in the flight, the volume of air in the bottle has increased by $1\cdot2\times10^{-4}\,\text{m}^3$.

During the flight the temperature of the air in the bottle remains constant.

(i) Calculate the pressure of the air inside the bottle at this point in the flight.

Space for working and answer



page 25

MARKS DO NOT WRITE IN THIS MARGIN

3

8. (c) (continued)

(ii) Using the kinetic model, explain what happens to the pressure of the air inside the bottle as the volume of the air increases.

page 26

[Turn over for next question

DO NOT WRITE ON THIS PAGE



page 27

MARKS DO NOT WRITE IN THIS MARGIN

A lifeboat crew is made up of local volunteers. When there is an emergency they have to get to the lifeboat quickly.

The lifeboat crew members are alerted to an emergency using a pager.

Text messages are sent to the pager using radio waves.



(a) The radio waves have a frequency of 153 MHz. Calculate the wavelength of the radio waves. Space for working and answer

3

(b) When the pager receives a message it beeps loudly and a light on the pager flashes.


A crew member holding the pager observes the beeps and the flashes happening at the same time.

A second crew member, who is 100 m away from the pager, also observes the beeps and the flashes.

Explain why the second crew member does not observe the beeps and the flashes happening at the same time.

(continued)

(c) The lifeboat has a mass of 25 000 kg. When it is launched, it loses 4.5×10^5 J of gravitational potential energy before it enters the water.



(i) Calculate the maximum speed of the lifeboat as it enters the water. 3 Space for working and answer

(ii) Explain why, in practice, the speed of the lifeboat as it enters the water is less than calculated in (c) (i).



page 29


(a) State the name given to this family of waves.

1

(b) State how the frequency of infrared compares to the frequency of gamma rays.

1

(c) Some examples of sources and detectors of waves in this family are shown.



- (i) From the examples shown, identify
 - (A) the detector of infrared

1

(B) the source of gamma rays.


1

(ii) Suggest one application for the waves that are detected using fluorescent ink.

2

1

- A student carries out an experiment to investigate the effect of different shaped glass blocks on the path of a ray of light.
 - (a) The student directs a ray of red light at a triangular glass block as shown.



(i) Complete the diagram above to show the path of the ray of red light through and out of the glass block.

(An additional diagram, if required, can be found on page 39)

(ii) The diagram shows a dashed line PQ. State the name given to this line.

(iii) On the diagram above, label an angle of incidence i. 1

(b) The student replaces the triangular glass block with a rectangular block made of the same material. The path of the ray of red light is as shown.



State whether the wavelength of the red light in this block is less than, the same as, or greater than the wavelength of the red light in the triangular glass block in (a).


Justify your answer.

DO NOT WRITE ON THIS PAGE

page 32

3

12. A technician carries out an experiment, using the apparatus shown, to determine the half-life of a radioactive source.




(a) Describe how the apparatus can be used to determine the half-life of the radioactive source.

[Turn over



12. (continued)

(b) The technician carries out the experiment over a period of 30 minutes, and displays the data obtained in a graph as shown.



Suggest an improvement that the technician could make to the procedure to more easily determine a value for the half-life of this source.

1

(c) In a second experiment, the technician absorbs $1\cdot 2\,\mu J$ of energy throughout their body from a radioactive source.

The mass of the technician is $80.0 \, \text{kg}$.

(i) Calculate the absorbed dose received by the technician.

Space for working and answer

12. (c) (continued)

(ii) During the experiment, the technician receives an equivalent dose of $4\cdot5\times10^{-8}\,\text{Sv.}$


Calculate the radiation weighting factor of this source.

3

Space for working and answer

(d) The technician wears a film badge to monitor exposure to radiation.

The film badge contains a piece of photographic film behind windows of different materials.



Explain how this badge is used to determine the type of radiation the technician has been exposed to.

2

[Turn over for next question



MARKS DO NOT WRITE IN THIS MARGIN

13. A physics teacher makes the following statement.


'Instead of nuclear fission, perhaps one day nuclear fusion will become a practical source of generating energy.'

Using your knowledge of physics, comment on the similarities and/or differences between using nuclear fission and nuclear fusion to generate energy.

3

[END OF QUESTION PAPER]








ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK

Additional graph paper for Q5 (a) (i)




ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK

Additional diagram for Q8 (a)



Additional diagram for Q11 (a) (i)



MARKS DO NOT WRITE IN THIS MARGIN

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



page 40

MARKS DO NOT WRITE IN THIS MARGIN

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



page 41

DO NOT WRITE ON THIS PAGE



page 42

DO NOT WRITE ON THIS PAGE



page 43

DO NOT WRITE ON THIS PAGE

 $Acknowledgement\ of\ copyright$

Question 3 Viktoyriav/shutterstock.com

Question 9 RNLI logo is reproduced by kind permission of RNLI.



page 44



X857/75/11

Physics Relationships sheet

WEDNESDAY, 15 MAY 1:00 PM - 3:30 PM





$$d = vt$$

$$d = \overline{v}t$$

$$s = vt$$

$$s = \overline{v}t$$

$$a = \frac{v - u}{t}$$

$$F = ma$$

$$W = mg$$

$$E_w = Fd$$

$$E_p = mgh$$

$$E_k = \frac{1}{2}mv^2$$

$$Q = It$$

$$V = IR$$

$$V_2 = \left(\frac{R_2}{R_1 + R_2}\right) V_S$$

$$\frac{V_1}{V_2} = \frac{R_1}{R_2}$$

$$R_T = R_1 + R_2 + \dots$$

$$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$$

$$P = \frac{E}{t}$$

$$P = IV$$

$$P = I^2 R$$

$$P = \frac{V^2}{R}$$

$$E_h = cm\Delta T$$

$$E_h = ml$$

$$p = \frac{F}{A}$$

$$p_1V_1 = p_2V_2$$

$$\frac{p_1}{T_1} = \frac{p_2}{T_2}$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\frac{pV}{T}$$
 = constant

$$f = \frac{N}{t}$$

$$v = f\lambda$$

$$T = \frac{1}{f}$$

$$A = \frac{N}{t}$$

$$D = \frac{E}{m}$$

$$H = Dw_r$$

$$\dot{H} = \frac{H}{t}$$

Additional relationships

Circle

circumference = $2\pi r$

$$area = \pi r^2$$

Sphere

area =
$$4\pi r^2$$

volume =
$$\frac{4}{3}\pi r^3$$

Trigonometry

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$\cos\theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$\tan\theta = \frac{\mathsf{opposite}}{\mathsf{adjacent}}$$

$$\sin^2\theta + \cos^2\theta = 1$$

Electron Arrangements of Elements

		87 Fr 2,8,18,32, 18,8,1 Francium	55 Cs 2,8,18,18, 8,1 Caesium	37 Rb 2,8,18,8,1 Rubidium	19 K 2,8,8,1 Potassium	11 Na 2,8,1 Sodium	Lithium	ζ [: ω	1 1 Hydrogen	Group 1 (1)
	Lan	88 Ra 2,8,18,32, 18,8,2 Radium	56 Ba 2,8,18,18, 8,2 Barium	38 Sr 2,8,18,8,2 Strontium	20 Ca 2,8,8,2 Calcium	Mg 2,8,2 Magnesium	Beryllium	Be	(2)	Group 2
Actinides	Lanthanides	89 Ac 2,8,18,32, 18,9,2 Actinium	57 La 2,8,18,18, 9,2 Lanthanum	39 Y 2,8,18,9,2 Yttrium	21 Sc 2,8,9,2 Scandium	(3)				
89 Ac 2,8,18,32, 18,9,2 Actinium	57 La 2,8,18, 18,9,2 Lanthanum	104 Rf 2,8,18,32, 32,10,2 Rutherfordium	72 Hf 2,8,18,32, 10,2 Hafnium	40 Zr 2,8,18, 10,2 Zirconium	22 Ti 2,8,10,2 Titanium	(4)			Key	
90 Th 2,8,18,32, 18,10,2 Thorium	58 Ce 2,8,18, 20,8,2 Cerium	105 Db 2,8,18,32, 32,11,2 Dubnium	73 Ta 2,8,18, 32,11,2 Tantalum	41 Nb 2,8,18, 12,1 Niobium	23 V 2,8,11,2 Vanadium	(5)		ר	Ato	,
91 Pa 2,8,18,32, 20,9,2 Protactinium	59 Pr 2,8,18,21, 8,2 Praseodymium	106 Sg 2,8,18,32, 32,12,2 Seaborgium	74 W 2,8,18,32, 12,2 Tungsten	42 Mo 2,8,18,13, 1 Molybdenum	24 Cr 2,8,13,1 Chromium	(6)		Name	Atomic number Symbol	
92 U 2,8,18,32, 21,9,2 Uranium	60 Nd 2,8,18,22, 8,2 Neodymium	107 Bh 2,8,18,32, 32,13,2 Bohrium	75 Re 2,8,18,32, 13,2 Rhenium	43 Tc 2,8,18,13, 2 Technetium	25 Mn 2,8,13,2 Manganese	Transition Elements			oer oer	Ú
93 Np 2,8,18,32, 22,9,2 Neptunium	61 Pm 2,8,18,23, 8,2 Promethium	108 Hs 2,8,18,32, 32,14,2 Hassium	76 Os 2,8,18,32, 14,2 Osmium	44 Ru 2,8,18,15, 1 Ruthenium	26 Fe 2,8,14,2 Iron	Element (8)				
94 Pu 2,8,18,32, 24,8,2 Plutonium	62 Sm 2,8,18,24, 8,2 Samarium	109 Mt 2,8,18,32, 32,15,2 Meitnerium	77 Ir 2,8,18,32, 15,2 Iridium	45 Rh 2,8,18,16, 1 Rhodium	27 Co 2,8,15,2 Cobalt	s (9)				
95 Am 2,8,18,32, 25,8,2 Americium	63 Eu 2,8,18,25, 8,2 Europium	110 Ds 2,8,18,32, 32,17,1 Darmstadtium	78 Pt 2,8,18,32, 17,1 Platinum	46 Pd 2,8,18, 18,0 Palladium	28 Ni 2,8,16,2 Nickel	(10)				•
96 Cm 2,8,18,32, 25,9,2 Curium	64 Gd 2,8,18,25, 9,2 Gadolinium	110 111 112 Ds Rg Cn 2,8,18,32, 2,8,18,32, 2,8,18,32, 32,17,1 32,18,1 32,18,2 Darmstadtium Roentgenium Copernicium	79 Au 2,8,18, 32,18,1 Gold	47 Ag 2,8,18, 18,1 Silver	29 Cu 2,8,18,1 Copper	(11)				
97 Bk 2,8,18,32, 27,8,2 Berkelium	65 Tb 2,8,18,27, 8,2 Terbium	112 Cn 2,8,18,32, 32,18,2 Copernicium	80 Hg 2,8,18, 32,18,2 Mercury	48 Cd 2,8,18, 18,2 Cadmium	30 Zn 2,8,18,2 Zinc	(12)				
98 Cf 2,8,18,32, 28,8,2 Californium	66 Dy 2,8,18,28, 8,2 Dysprosium		81 T (2,8,18, 32,18,3 Thallium	49 In 2,8,18, 18,3 Indium	31 Ga 2,8,18,3 Gallium	2,8,3 Aluminium	Boron	 .	(13)	Group 3
99 Es 2,8,18,32, 29,8,2 Einsteinium	67 Ho 2,8,18,29, 8,2 Holmium		82 Pb 2,8,18, 32,18,4 Lead	50 Sn 2,8,18, 18,4 Tin	32 Ge 3 2,8,18,4 Germanium	14 Si 2,8,4 m Silicon	Carbon	. n °	(14)	3 Group 4
100 Fm 2,8,18,32, 30,8,2 Fermium	68 Er 2,8,18,30, 8,2 Erbium		83 Bi 2,8,18, 4 32,18,5 8ismuth	51 Sb 2,8,18, 18,5 Antimony	33 As 4 2,8,18,5 am Arsenic	2,8,5 Phosphorus	Z	3 Z 7	(15)	4 Group 5
101 Md 2,8,18,32, 31,8,2 Mendelevium	69 Tm 2,8,18,31, 8,2 Thulium		84 Po 2,8,18, 32,18,6	52 Te 2,8,18, 18,6 y Tellurium	34 Se 5 2,8,18,6 Selenium	2,8,6	9	ડે ૦ ∞	(16)	5 Group 6
102 No 2,8,18,32, 32,8,2 Nobelium	70 Yb 2,8,18,32, 8,2 Ytterbium		85 At 2,8,18, 32,18,7 n Astatine	53 2,8,18, 18,7 n lodine	35 Br 6 2,8,18,7 n Bromine	2,8,7 Chlorine	 	3 ₇ m 9	(17)	6 Group 7
103 Lr 2,8,18,32, 32,9,2 Lawrencium	71 Lu 2,8,18,32, 9,2 Lutetium		86 Rn 2,8,18, 32,18,8 Radon	54 Xe 2,8,18, 18,8 Xenon	36 Kr 7 2,8,18,8 e Krypton	18 Ar 2,8,8 e Argon	7	3 Z 10	2 He 2 Helium	7 Group 0
										J