



JABchem



Not to be shared without the copyright holder's permission

Past Papers Nat 5 Chemistry

2024 Marking Scheme

Grade Awarded	Mark Required		% candidates achieving grade
	(/125)	%	
A	86+	68.8%	41.5%
B	73+	58.4%	18.8%
C	60+	48.0%	16.3%
D	47+	37.6%	12.9%
No award	<47	<37.6%	10.4%

Section:	Multiple Choice	Extended Answer	Assignment
Average Mark:	16.2 /25	44.7 /75	17.4 /25

2024 National 5 Chemistry Marking Scheme

MC Qu	Answer	% Pupils Correct	Reasoning																
1	B	65	$\text{Rate} = \frac{\Delta \text{Quantity}}{\Delta \text{Time}} = \frac{0.6 \text{ bar}}{5 \text{ min}} = 0.12 \text{ bar min}^{-1}$																
2	D	72	Isotopes have same number of protons but different number of neutrons ☒ A W and X have same number of neutrons ☒ B X and Z have different numbers of protons ☒ C W and Y have different numbers of protons ☒ D Y and Z have same number of protons but different numbers of neutrons																
3	D	66	Atomic number of Sn = 50 ∴ no. of protons = 50 Mass number = no. of protons + no. of neutrons = 50 + 74 = 124																
4	A	46	☒ A tetraethylplumbane would be a covalent molecular substance as it is a liquid at room temperature and does not conduct in any state. ☒ B a covalent network would be solid at room temperature with a high melting point ☒ C an ionic lattice would be a solid at room temperature ☒ D a metallic substance would conduct electricity in the liquid state																
5	C	76	☒ A The O atom is not directly bonded to a hydrogen atom ∴ no hydrogen bonding ☒ B The N atom is not directly bonded to a hydrogen atom ∴ no hydrogen bonding ☒ C The O atom is directly bonded to a hydrogen atom ∴ hydrogen bonding present ☒ D The N atom is not directly bonded to a hydrogen atom ∴ no hydrogen bonding																
6	B	82	☒ A aluminium oxide is insoluble in water so there would be no change to the pH ☒ B calcium oxide is soluble in water ∴ pH would increase as non-metal oxides dissolve to form alkaline solutions ☒ C copper(II) oxide is insoluble in water so there would be no change to the pH ☒ D zinc(II) oxide is insoluble in water so there would be no change to the pH																
7	B	63	$ \begin{array}{ccc} 2\text{Mg} + \text{O}_2 & \longrightarrow & 2\text{MgO} \\ \begin{array}{cc} 2\text{mol} & 1\text{mol} \\ 4\text{mol} & 2\text{mol} \end{array} & & \begin{array}{c} 2\text{mol} \\ 4\text{mol} \end{array} \end{array} $																
8	D	25	☒ A Nitrogen N ₂ gfm = 2×14 = 28g ☒ B Oxygen O ₂ gfm = 2×16 = 32g ☒ C Fluorine F ₂ gfm = 2×19 = 38g ☒ D Neon Ne gfm = 1×20 = 20g																
9	C	45	☒ A same number of moles of sodium hydroxide present after 25cm ³ water added ☒ B dilution of alkali would lower pH as alkali is more dilute after 25cm ³ water added ☒ C number of moles of sodium hydroxide present is constant and pH would be lower ☒ D same number of moles of sodium hydroxide present after 25cm ³ water added																
10	A	68	Alkanes are a homologous series of hydrocarbons which are flammable. Combustion of alkanes are exothermic reactions which would be suitable for use as a fuel.																
11	C	79	<table border="1"> <thead> <tr> <th>Name</th><th>cyclooctyne</th><th>cyclononyne</th><th>cyclodecyne</th></tr> </thead> <tbody> <tr> <td>Number of carbons (n)</td><td>n = 8</td><td>n = 9</td><td>n = 10</td></tr> <tr> <td>Hydrogens: 2n-4</td><td>(2×8)-4 = 16-4 = 12</td><td>(2×9)-4 = 18-4 = 14</td><td>(2×10)-4 = 20-4 = 16</td></tr> <tr> <td>Formula</td><td>C₈H₁₂</td><td>C₉H₁₄</td><td>C₁₀H₁₆</td></tr> </tbody> </table>	Name	cyclooctyne	cyclononyne	cyclodecyne	Number of carbons (n)	n = 8	n = 9	n = 10	Hydrogens: 2n-4	(2×8)-4 = 16-4 = 12	(2×9)-4 = 18-4 = 14	(2×10)-4 = 20-4 = 16	Formula	C ₈ H ₁₂	C ₉ H ₁₄	C ₁₀ H ₁₆
Name	cyclooctyne	cyclononyne	cyclodecyne																
Number of carbons (n)	n = 8	n = 9	n = 10																
Hydrogens: 2n-4	(2×8)-4 = 16-4 = 12	(2×9)-4 = 18-4 = 14	(2×10)-4 = 20-4 = 16																
Formula	C ₈ H ₁₂	C ₉ H ₁₄	C ₁₀ H ₁₆																
12	B	59	Pent-2-ene has formula C ₅ H ₁₀ . Isomers have same formula but different structure. ☒ A Pentane has formula C ₅ H ₁₂ ∴ Different formula means not an isomer ☒ B Pent-1-ene C ₅ H ₁₀ has same formula as pent-2-ene but different structure ☒ C Molecule is pent-2-ene ∴ Same formula & same structure means not an isomer ☒ D 2-methylbutane has formula C ₅ H ₁₂ ∴ Different formula means not an isomer																

13	B	90	H H CH₃ H O H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃ H H H CH₃
----	---	----	--




23	C	76	$\begin{array}{ccccc} \text{nitrogen} & + & \text{hydrogen} & \xrightleftharpoons{\text{iron catalyst}} & \text{ammonia} \\ \text{N}_2 & + & 3\text{H}_2 & \xrightleftharpoons{\text{iron catalyst}} & 2\text{NH}_3 \end{array}$
24	D	53	${}^{228}_{88}\text{Ra} \xrightarrow{\beta} {}^{228}_{89}\text{Ac} \xrightarrow{\beta} {}^{228}_{90}\text{Th} \xrightarrow{\alpha} {}^{224}_{88}\text{Ra}$
25	D	71	☒ A copper gives a flame test colour of blue-green ☒ B copper gives a flame test colour of blue-green ☒ C copper(II) Cu^{2+} ions are blue in solution ☒ D copper(II) Cu^{2+} ions are blue in solution and a flame test colour of blue-green



2024 National 5 Chemistry Marking Scheme

Long Qu	Answer	Reasoning												
1a(i)	One answer from:	$2\text{H}_2\text{O}_2 \longrightarrow 2\text{H}_2\text{O} + \text{O}_2$ or $\text{H}_2\text{O}_2 \longrightarrow \text{H}_2\text{O} + \frac{1}{2}\text{O}_2$												
1a(ii)	catalyst	A catalyst is a chemical which speeds up a chemical reaction without being used up in the reaction. The same mass of catalyst is left at the end of the reaction.												
1a(iii)A	Insoluble in water	For a gas to be collected over water by this method, the gas must be insoluble in water so the full volume of gas collects in the inverted measuring cylinder. Examples of soluble & insoluble gases include: <table><tr><td>Soluble Gases</td><td>NO_2</td><td>NH_3</td><td>HCl</td><td>SO_2</td><td>(CO_2)</td></tr><tr><td>Insoluble Gases</td><td>N_2</td><td>O_2</td><td>noble gases</td><td>hydrocarbons</td><td></td></tr></table> CO_2 is slightly soluble in water so some CO_2 will dissolve in water (like fizzy drinks) but CO_2 can be collected by this method in some experiments. The best way to collect soluble gases is in a gas syringe.	Soluble Gases	NO_2	NH_3	HCl	SO_2	(CO_2)	Insoluble Gases	N_2	O_2	noble gases	hydrocarbons	
Soluble Gases	NO_2	NH_3	HCl	SO_2	(CO_2)									
Insoluble Gases	N_2	O_2	noble gases	hydrocarbons										
1a(iii)B	36	$\text{Rate} = \frac{\Delta \text{Quantity}}{\Delta \text{Time}}$ $1.2 \text{ cm}^3 \text{ s}^{-1} = \frac{\Delta \text{Quantity}}{30 \text{ s}}$ $1.2 \text{ cm}^3 \text{ s}^{-1} \times 30 \text{ s} = \Delta \text{Quantity}$ $36 \text{ cm}^3 = \Delta \text{Quantity}$												
1a(iii)C	Less than 1.2	Reaction rates decrease as reactions proceed as the concentration of reactants decreases as the reactants get used up. There are less collisions between the lower concentration of reactants which lowers the reaction rate. The average reaction rate must be lower between 60 and 90 seconds than it was between 0 and 30 seconds.												
1b(i)	limewater turns cloudy/milky	<table><tr><th>Gas</th><th>Test for Gas</th></tr><tr><td>Carbon dioxide</td><td>turns limewater milky</td></tr><tr><td>Hydrogen</td><td>burns with a pop</td></tr><tr><td>Oxygen</td><td>relights a glowing splint</td></tr></table>	Gas	Test for Gas	Carbon dioxide	turns limewater milky	Hydrogen	burns with a pop	Oxygen	relights a glowing splint				
Gas	Test for Gas													
Carbon dioxide	turns limewater milky													
Hydrogen	burns with a pop													
Oxygen	relights a glowing splint													
1b(ii)	endothermic	<table><tr><th>Type of Reaction</th><th>Definition</th></tr><tr><td>Exothermic</td><td>Reaction which releases (heat) energy</td></tr><tr><td>Endothermic</td><td>Reaction which takes in (heat) energy from the surroundings</td></tr></table>	Type of Reaction	Definition	Exothermic	Reaction which releases (heat) energy	Endothermic	Reaction which takes in (heat) energy from the surroundings						
Type of Reaction	Definition													
Exothermic	Reaction which releases (heat) energy													
Endothermic	Reaction which takes in (heat) energy from the surroundings													
2a(i)	trigonal pyramidal	<table><tr><th>HCl</th><th>H_2O</th><th>NH_3</th><th>CH_4</th></tr><tr><td></td><td></td><td></td><td></td></tr><tr><td>Linear</td><td>Angular</td><td>Trigonal Pyramidal</td><td>Tetrahedral</td></tr></table>	HCl	H_2O	NH_3	CH_4					Linear	Angular	Trigonal Pyramidal	Tetrahedral
HCl	H_2O	NH_3	CH_4											
Linear	Angular	Trigonal Pyramidal	Tetrahedral											
2a(ii)	Answer showing:	or												

2b	Nitric Acid	The end product of the Ostwald Process is nitric acid. Ammonia NH ₃ and Oxygen O ₂ are the reactants which react to form nitrogen monoxide NO and water H ₂ O using a platinum catalyst. Nitrogen monoxide NO reacts with air and water to form nitric acid HNO ₃ Heat is given off in this reaction and the heat supply to start the reaction can be turned off once the reaction has started.																													
2c	Graph showing:	1 mark One mark is awarded for a graph which shows points plotted rather than bars.	1 mark The axis/axes of the graph has/have suitable scale(s). - plotted points occupies at least half of the width and half of the height of the graph paper - The axes have suitable scales	1 mark The axes of the graph have suitable labels and units.	1 mark All data points plotted accurately (within a half box tolerance) with either a line of best fit drawn or plots joined by curve. This mark can only be accessed if linear scales for both axes have been provided.																										
2d	Arrow drawn as shown:	Reactors Ammonia and carbon dioxide																													
3a	Essential oils	Problem Solving: Information from a passage																													
3b	<table><tr><td>0.885</td><td>0.905</td></tr><tr><td>204</td><td>183</td></tr></table>	0.885	0.905	204	183	Problem Solving: Transfer of data from passage to table																									
0.885	0.905																														
204	183																														
3c	Lavender Spike Oil	Problem Solving: Information from a passage																													
3d	Unsaturated oils will decolourise bromine solution	Bromine solution is decolourised when unsaturated hydrocarbons with C=C double bonds are present. The bromine molecules in bromine solution add across the C=C double bonds in the essential oils and the bromine solution loses its yellow colour when all the bromine molecules add onto the unsaturated essential oil molecules																													
3e	Stronger intermolecular bonds	Geraniol has a higher boiling point than linalool and nerol because the geraniol. Molecules are closer to each other duen to stronger intermolecular forces/bonds/attractions between molecules.																													
3f	2.643	<table><tr><td>mass (g)</td><td>=</td><td>density (g cm⁻³)</td><td>x</td><td>volume (cm³)</td></tr><tr><td>mass (g)</td><td>=</td><td>0.881 g cm⁻³</td><td>x</td><td>3 cm³</td></tr><tr><td>mass (g)</td><td>=</td><td>2.643 g</td><td></td><td></td></tr></table>						mass (g)	=	density (g cm ⁻³)	x	volume (cm ³)	mass (g)	=	0.881 g cm ⁻³	x	3 cm ³	mass (g)	=	2.643 g											
mass (g)	=	density (g cm ⁻³)	x	volume (cm ³)																											
mass (g)	=	0.881 g cm ⁻³	x	3 cm ³																											
mass (g)	=	2.643 g																													
4a	homologous series	A homologous series is a family of compounds with same/similar chemical properties and a general formula. Examples of homologous series include: <table><tr><td>Homologous Series</td><td>Alkane</td><td>Alkene</td><td>Cycloalkane</td><td>Alcohol</td><td>Carboxylic acid</td></tr><tr><td>General Formula</td><td>C_nH_{2n+2}</td><td>C_nH_{2n}</td><td>C_nH_{2n}</td><td>C_nH_{2n+1}OH</td><td>C_nH_{2n+1}COOH</td></tr></table>						Homologous Series	Alkane	Alkene	Cycloalkane	Alcohol	Carboxylic acid	General Formula	C _n H _{2n+2}	C _n H _{2n}	C _n H _{2n}	C _n H _{2n+1} OH	C _n H _{2n+1} COOH												
Homologous Series	Alkane	Alkene	Cycloalkane	Alcohol	Carboxylic acid																										
General Formula	C _n H _{2n+2}	C _n H _{2n}	C _n H _{2n}	C _n H _{2n+1} OH	C _n H _{2n+1} COOH																										
4b(i)	141	<table><tr><td>Name</td><td>methanol</td><td>ethanol</td><td>propan-1-ol</td><td>butan-1-ol</td><td>pentan-1-ol</td></tr><tr><td>Boiling Point (°C)</td><td>65</td><td>78</td><td>97</td><td>118</td><td>-</td></tr><tr><td>Difference (°C)</td><td></td><td>13</td><td>19</td><td>21</td><td>(23)</td></tr><tr><td>Prediction (°C)</td><td>-</td><td>-</td><td>-</td><td>-</td><td>141</td></tr></table>						Name	methanol	ethanol	propan-1-ol	butan-1-ol	pentan-1-ol	Boiling Point (°C)	65	78	97	118	-	Difference (°C)		13	19	21	(23)	Prediction (°C)	-	-	-	-	141
Name	methanol	ethanol	propan-1-ol	butan-1-ol	pentan-1-ol																										
Boiling Point (°C)	65	78	97	118	-																										
Difference (°C)		13	19	21	(23)																										
Prediction (°C)	-	-	-	-	141																										
4b(ii)	Increasing in carbons increases the boiling point	The longer the carbon chain in the alcohols the higher the boiling point due to stronger intermolecular bonds/attractions/forces.																													
4c(i)	hydroxyl carboxyl	<table><tr><td>—O—H</td><td>$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$</td></tr><tr><td>hydroxyl group</td><td>carboxyl group</td></tr></table>						—O—H	$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$	hydroxyl group	carboxyl group																				
—O—H	$\begin{array}{c} \text{O} \\ \\ \text{—C—OH} \end{array}$																														
hydroxyl group	carboxyl group																														

4c(ii)		 propan-1-ol → propanoic acid butan-1-ol → butanoic acid																																					
4c(iii)	$(K^+)_2Cr_2O_7^{2-}$	Write down valency below each element's symbol $K \quad Cr_2O_7^{2-}$ 1 2	Put in Cross-over Arrows $K \quad Cr_2O_7$ 1 2	Follow arrows to get formula $K_2Cr_2O_7$ add charges $(K^+)_2Cr_2O_7^{2-}$																																			
5	Open Ended Question: <table><tr><th colspan="4">Candidate Marks</th></tr><tr><td>0 marks</td><td>1 mark</td><td>2 marks</td><td>3 marks</td></tr><tr><td>6%</td><td>50%</td><td>34%</td><td>9%</td></tr></table>	Candidate Marks				0 marks	1 mark	2 marks	3 marks	6%	50%	34%	9%	3 mark answer Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	2 mark answer Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	1 mark answer Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.																							
		Candidate Marks																																					
0 marks	1 mark	2 marks	3 marks																																				
6%	50%	34%	9%																																				
6a	Delocalised electrons	Metals have metallic bonding where positive ions (nucleus and inner shells) are surrounded by delocalised electrons (in the outer shell). Those delocalised electrons can jump from outer shell to outer shell so electricity conduction can take place.																																					
6b	Heat with carbon	<table><tr><th>Method</th><th colspan="2">Electrolysis</th><th colspan="2">Heat With Carbon</th><th colspan="2">Heat Alone</th></tr><tr><td rowspan="3">Metals Made This Way</td><td>Potassium</td><td>Sodium</td><td>Zinc</td><td>Iron</td><td>Mercury</td><td>Silver</td></tr><tr><td>Lithium</td><td>Calcium</td><td>Nickel</td><td>Tin</td><td>Gold</td><td>Platinum</td></tr><tr><td>Magnesium</td><td>Aluminium</td><td>Lead</td><td>Copper</td><td></td><td></td></tr><tr><td>Reason</td><td colspan="2">most reactive metals</td><td colspan="2">medium reactive metals</td><td colspan="2">least reactive metals</td></tr></table>	Method	Electrolysis		Heat With Carbon		Heat Alone		Metals Made This Way	Potassium	Sodium	Zinc	Iron	Mercury	Silver	Lithium	Calcium	Nickel	Tin	Gold	Platinum	Magnesium	Aluminium	Lead	Copper			Reason	most reactive metals		medium reactive metals		least reactive metals					
Method	Electrolysis		Heat With Carbon		Heat Alone																																		
Metals Made This Way	Potassium	Sodium	Zinc	Iron	Mercury	Silver																																	
	Lithium	Calcium	Nickel	Tin	Gold	Platinum																																	
	Magnesium	Aluminium	Lead	Copper																																			
Reason	most reactive metals		medium reactive metals		least reactive metals																																		
6c(i)	<table><tr><td>C</td></tr><tr><td>A</td></tr><tr><td>copper</td></tr><tr><td>D</td></tr><tr><td>B</td></tr></table>	C	A	copper	D	B	Direction of electron Flow: Metals A and C must be above copper in electrochemical series as electrons flow from metals A and C to copper. Metals B and D must be below copper in electrochemical series as electrons flow from copper to metals B and D.		Size of Voltage: Metal C gives a bigger voltage with copper than metal A with copper does. Metal C is higher up electrochemical series than metal A Metal B gives a bigger voltage with copper than metal D with copper does. Metal B is lower down electrochemical series than metal D																														
C																																							
A																																							
copper																																							
D																																							
B																																							
6c(ii)	Glucose (is molecular and) contains no ions	Glucose is covalent because there are only non-metals in its formula $C_6H_{12}O_6$. To complete the circuit an ionic solution must be used as an electrolyte but covalent molecular glucose contains no ions to complete the circuit.																																					
6d(i)	Ion/salt bridge	An ion bridge (or salt bridge) is filter paper soaked in an ionic solution.																																					
6d(ii)	Any soluble lead compound:	One from:	Lead bromide	Lead chloride	Lead nitrate																																		
7a	Kills bacteria and viruses	Problem Solving: Information from a passage																																					
7b	sodium, hydrogen, sulfur and oxygen	sodium hydrogen sulfur oxygen sodium hydrogensulfite																																					
7c	$Cl_2 + H_2O$ ↓ $HOCl + HCl$	chlorine	+	water	→	hypochlorous acid	+	hydrochloric acid																															
		Cl_2	+	H_2O	→	$HOCl$	+	HCl																															

7d	Diagram showing: 	Example:  Question: 
8a(i)	Equation showing:	$\text{Ag}^+_{(\text{aq})} + \text{NO}_3^-_{(\text{aq})} + \text{Na}^+_{(\text{aq})} + \text{Cl}^-_{(\text{aq})} \rightarrow \text{Ag}^+\text{Cl}^-_{(\text{s})} + \text{Na}^+_{(\text{aq})} + \text{NO}_3^-_{(\text{aq})}$ Identify Spectator Ions and cancel out Spectator Ions $\text{Ag}^+_{(\text{aq})} + \cancel{\text{NO}_3^-_{(\text{aq})}} + \cancel{\text{Na}^+_{(\text{aq})}} + \text{Cl}^-_{(\text{aq})} \rightarrow \text{Ag}^+\text{Cl}^-_{(\text{s})} + \cancel{\text{Na}^+_{(\text{aq})}} + \cancel{\text{NO}_3^-_{(\text{aq})}}$ Re-write equation without spectator ions $\text{Ag}^+_{(\text{aq})} + \text{Cl}^-_{(\text{aq})} \rightarrow \text{Ag}^+\text{Cl}^-_{(\text{s})}$
8a(ii)	Precipitation	A precipitation reaction occurs when two ions in solution meet when two solutions are mixed. Some ion combinations form insoluble precipitates when their ions meet. - Take the names of the two original solutions - silver nitrate + sodium chloride - Swap the names over between the two solutions - silver chloride + sodium nitrate - Look up the data booklet for an insoluble precipitate that would form - silver chloride is insoluble and sodium nitrate is soluble - Silver chloride is a precipitate which can be removed by filtration.
8b(i)	pipette	Pipettes are the most accurate way to transfer fixed volumes of solutions in a titration. They are more accurate than burettes, measuring cylinder and beakers.
8b(ii)	1 st titre is inaccurate	The first titre, also known as the rough titre, is not designed to be accurate but to give an indication of where the endpoint of the reaction is. The experiment is repeated and the majority of the volume of reactant from the burette is added in one large addition and then the remaining volume is added drop by drop until the indicator changes colour. This repeated into two titres are produced within 0.2cm ³ .
8b(iii)	0.312	no. of mol = volume x concentration = 0.0156litre x 0.1mol l⁻¹ = 0.00156mol $\text{H}_2\text{C}_2\text{O}_4 + 2\text{NaOH} \rightarrow \text{Na}_2\text{C}_2\text{O}_4 + 2\text{H}_2\text{O}$ 1mol 2mol 0.00156mol 0.00312mol $\text{concentration} = \frac{\text{no. of mol}}{\text{volume}} = \frac{0.00312\text{mol}}{0.01\text{litres}} = 0.312 \text{ mol l}^{-1}$
8b(iv)	Pink to colourless	The conical flask contains sodium hydroxide at the beginning and the phenolphthalein indicator is pink in alkaline conditions. As oxalic acid is added from the burette into the conical flask, the sodium hydroxide is neutralised and the pH of the contents decreases until it reaches pH=7 when neutralisation occurs and the indicator changes colour to colourless.
9a(i)	No fizzing when more nickel carbonate is added	Nickel(II) carbonate reacts with sulfuric acid, carbon dioxide gas is produced in the reaction. The reaction mixture will stop fizzing when the nickel(II) carbonate runs out so more nickel(II) carbonate is added and it will start to fizz again. The process is repeated until adding more nickel(II) carbonate doesn't start fizzing in the reaction mixture indicating there is no acid left and neutralisation is complete.
9a(ii)	Water	All neutralisation reactions produce water. acid + metal oxide → salt + water acid + metal hydroxide → salt + water acid + metal carbonate → salt + water + carbon dioxide

9a(iii)	Base	A base is a chemical which neutralises an acid. There are three types of bases																							
		metal oxide	metal hydroxide	metal carbonate																					
9a(iv)	0.025	no. of mol = volume x concentration = 0.025litre x 1mol l ⁻¹ = 0.025mol																							
9b	filtration	Insoluble solids are removed from liquids by filtrations. - the insoluble solid is the residue in the filter paper - the liquid is the filtrate that goes through the filter paper and collects underneath.																							
9c	Heat solution with Bunsen burner	The water can be evaporated quickly by being heated by a Bunsen burner. The crystals are not as well formed but it does happen much quicker.																							
9d	37.9	gfm NiSO ₄ = (1x58.5)+(1x32)+(4x16) = 58.5 + 32 + 64 = 154.5g $\%Ni = \frac{\text{mass of Nickel}}{\text{gfm}} \times 100 = \frac{58.5}{154.5} \times 100 = 37.9\%$																							
10a	Answer to include:	Compound containing the elements carbon and hydrogen only																							
10b(i)	3460	<table><tr><td>Bond</td><td>Bond energy (kJ)</td><td>Number of bonds made</td><td>Total Energy Released (kJ)</td></tr><tr><td>C=O</td><td>804</td><td>2</td><td>2x 804 = 1608</td></tr><tr><td>O-H</td><td>463</td><td>4</td><td>4x 463 = 1852</td></tr><tr><td colspan="3">Total energy to make all the bonds in the products =</td><td>= 3460</td></tr></table>				Bond	Bond energy (kJ)	Number of bonds made	Total Energy Released (kJ)	C=O	804	2	2x 804 = 1608	O-H	463	4	4x 463 = 1852	Total energy to make all the bonds in the products =			= 3460				
Bond	Bond energy (kJ)	Number of bonds made	Total Energy Released (kJ)																						
C=O	804	2	2x 804 = 1608																						
O-H	463	4	4x 463 = 1852																						
Total energy to make all the bonds in the products =			= 3460																						
10b(ii)	- 816	Energy Change = Total energy to break all the bonds in the reactants - Total energy to make all the bonds in the products Energy Change = 2644 - 3460 Energy Change = - 816 kJ																							
10c(i)	2,4-dimethylhexane	 2,4-dimethylhexane Sidegroups on C₂ & C₄ 2x methyl -CH₃ sidegroups 6 carbons on main chain C-C single bonds in main chain																							
10c(ii)		Any C ₆ H ₁₂ structure contain a carbon ring is correct. Correct answers include: - cyclohexane - methylcyclopentane 1,2-dimethylcyclobutane 1,3-dimethylcyclobutane 1,1,2-trimethylcyclopropane 1,2,3-trimethylcyclopropane																							
10d(i)A	C ₆ H ₁₄	octane $\longrightarrow$ ethene + alkane X C ₈ H ₁₈ $\longrightarrow$ C ₂ H ₄ C ₆ H ₁₄ ∴ hexane																							
10d(i)B	hydration	Addition reactions add a molecule across a C=C double bond. - addition of water across C=C double bond is a hydration reaction - addition of hydrogen across C=C double bond is a hydrogenation reaction The following molecules are all able to add across a C=C double bond: <table><tr><td>H₂</td><td>F₂</td><td>Cl₂</td><td>Br₂</td><td>I₂</td><td>H₂O</td><td>HF</td><td>HCl</td><td>HBr</td><td>HI</td></tr><tr><td colspan="5">Addition reaction gives one product</td><td colspan="5">Addition reaction usually gives two product</td></tr></table>				H ₂	F ₂	Cl ₂	Br ₂	I ₂	H ₂ O	HF	HCl	HBr	HI	Addition reaction gives one product					Addition reaction usually gives two product				
H ₂	F ₂	Cl ₂	Br ₂	I ₂	H ₂ O	HF	HCl	HBr	HI																
Addition reaction gives one product					Addition reaction usually gives two product																				
10d(ii)A	poly(ethenol)	<table><tr><td>Monomer</td><td>ethene</td><td>propene</td><td>chloroethene</td><td>phenylethene</td><td>ethenol</td></tr><tr><td>Polymer</td><td>poly(ethene)</td><td>poly(propene)</td><td>poly(chloroethene)</td><td>poly(phenylethene)</td><td>poly(ethenol)</td></tr></table>				Monomer	ethene	propene	chloroethene	phenylethene	ethenol	Polymer	poly(ethene)	poly(propene)	poly(chloroethene)	poly(phenylethene)	poly(ethenol)								
Monomer	ethene	propene	chloroethene	phenylethene	ethenol																				
Polymer	poly(ethene)	poly(propene)	poly(chloroethene)	poly(phenylethene)	poly(ethenol)																				

10d(ii)B	H H —C—C— H OH	H H C=C H OH monomer H H H H H H —C—C—C—C—C—C— H OH H OH H OH polymer H H —C—C— H OH repeating unit														
11	Open Ended Question: Candidate Marks <table><tr><td>0 marks</td><td>1 mark</td><td>2 marks</td><td>3 marks</td></tr><tr><td>21%</td><td>41%</td><td>28%</td><td>11%</td></tr></table>	0 marks	1 mark	2 marks	3 marks	21%	41%	28%	11%	<table><tr><td>3 mark answer</td><td>2 mark answer</td><td>1 mark answer</td></tr><tr><td>Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.</td><td>Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.</td><td>Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.</td></tr></table>	3 mark answer	2 mark answer	1 mark answer	Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
0 marks	1 mark	2 marks	3 marks													
21%	41%	28%	11%													
3 mark answer	2 mark answer	1 mark answer														
Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.														