



JABchem



Not to be shared without the copyright holder's permission

Past Papers Higher Chemistry

2025 Marking Scheme

Grade Awarded	Mark Required		% candidates achieving grade
	(/150)	%	
A	108+	72%	32.0%
B	92+	61%	25.9%
C	77+	51%	18.2%
D	61+	41%	13.3%
No award	<61	<41%	10.6%

Section:	Multiple Choice	Extended Answer	Assignment
Average Mark:	17.6 /25	56.5 /95	19.7 /30

2025 Higher Chemistry Marking Scheme

MC Qu	Answer	% Pupils Correct	Reasoning
1	A	87	☒ A Electronegativity difference of N-Br bond = 3.0 - 3.0 = 0.0 (pure covalent) ☒ B Electronegativity difference of Cl-H bond = 3.2 - 2.2 = 1.0 ☒ C Electronegativity difference of O-H bond = 3.4 - 2.2 = 1.2 ☒ D Electronegativity difference of I-C bond = 2.7 - 2.6 = 0.1
2	C	77	☒ A Reducing agents are oxidised themselves and lose electrons ☒ B Strong reducing agents are metals on top right of Electrochemical Series. Metals are found on left side of the Period Table ☒ C Strong reducing agents are group 1+2 metals with low electronegativity values ☒ D Strong reducing agents are metals which lose electrons to form positive ions
3	B	72	no. of moles = volume x concentration = 0.1 litres x 0.5 mol l ⁻¹ = 0.05mol $\begin{array}{ccccccc} \text{MnO}_4^- & + & 5\text{Fe}^{2+} & + & 8\text{H}^+ & \longrightarrow & \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O} \\ 1\text{mol} & & 5\text{mol} & & & & \\ 0.05\text{mol} & & 0.25\text{mol} & & & & \end{array}$
4	D	53	3,4-dimethylhex-3-ene Side-groups on C₃ and C₄ 2x -CH₃ methyl side groups 6 Carbons in main chain Functional Group on C₃ C=C double bond
5	B	90	☒ A Primary: 1 carbon directly attached to the carbon with the -OH group ☒ B Tertiary: 3 carbons directly attached to the carbon with the -OH group ☒ C Secondary: 2 carbons directly attached to the carbon with the -OH group ☒ D Secondary: 2 carbons directly attached to the carbon with the -OH group
6	A	93	☒ A Structure is 3-ethylhexanoic acid ☒ B Structure is 2-ethylhexanoic acid ☒ C Structure is 3,3-dimethylhexanoic acid ☒ D Structure is 4,4-dimethylhexanoic acid
7	B	69	$\begin{array}{ccccccc} & \text{H} & & \text{O} & & \text{H} & \text{H} & \text{H} \\ & & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & - \text{O} & - & \text{C} & - \text{C} & - & \text{C} & - \text{H} \\ & & & & & & & \\ & \text{H} & & & & \text{H} & \text{H} & \text{H} \end{array}$ Carboxylic Acid ethanoic acid gfm = 60g Alcohol propan-1-ol gfm 60g
8	D	64	$\begin{array}{cccc} & \text{H} & \text{H} & \text{H} \\ & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} \\ & & & \\ & \text{H} & \text{H} & \text{H} \end{array}$ aldehyde butanal C₄H₈O gfm= 72g $\xrightarrow{\text{reduction}}$ $\begin{array}{cccc} & \text{H} & \text{H} & \text{H} & \text{H} \\ & & & & \\ \text{H} & - \text{C} & - \text{C} & - \text{C} & - \text{C} & - \text{OH} \\ & & & & \\ & \text{H} & \text{H} & \text{H} & \text{H} \end{array}$ primary alcohol butan-1-ol C₄H₁₀O gfm = 74g

9	C	74	6-methylhept-5-ene-2-one methyl -CH₃ group on C₆ Seven carbons on main chain C=C double bond between C₅ and C₆ C=O Carbonyl group on C₂ 7 6 5 4 3 2 1 CH₃C(CH₃)CHCH₂CH₂COCH₃ C=C double bond between C₅ and C₆ C=O Carbonyl group on C₂															
10	B	72	☒ A hydration: water added across a C=C double bond ☒ B fat is hydrolysed into glycerol and 3 fatty acids. Fatty acids then neutralised. ☒ C esterification: alcohol and carboxylic acid join together with H₂O removed ☒ D condensation: two molecules join together with small molecule e.g. H₂O removed															
11	D	82	☒ A fatty acid chains are hydrocarbon ∴ hydrophobic and insoluble in water ☒ B fatty acid chains are hydrocarbon ∴ hydrophobic and insoluble in water ☒ C fatty acid chains are hydrocarbon ∴ hydrophobic ☒ D fatty acid chains are hydrocarbon ∴ hydrophobic and soluble in oil droplets															
12	B	84	☒ A terpenes must have a formula with a multiple of five carbons ☒ B Terpene with formula C₁₅H₂₄ is made from three C₅H₈ isoprene joined together ☒ C terpenes must have a formula with a multiple of five carbons ☒ D terpenes must have a formula with a multiple of five carbons															
13	C	79	<table><tr><td>When rate = 0.02 s⁻¹</td><td>Temperature = 44°C</td><td rowspan="2">∴ Change in temperature = 12°C</td></tr><tr><td>When rate = 0.04 s⁻¹</td><td>Temperature = 56°C</td></tr></table>	When rate = 0.02 s ⁻¹	Temperature = 44°C	∴ Change in temperature = 12°C	When rate = 0.04 s ⁻¹	Temperature = 56°C										
When rate = 0.02 s ⁻¹	Temperature = 44°C	∴ Change in temperature = 12°C																
When rate = 0.04 s ⁻¹	Temperature = 56°C																	
14	A	71	☒ A Line T₂ is to the right and not as high as line T₁ ☒ B The area under the curves should be the same for both lines ☒ C Line T₂ should be lower than line T₁ ☒ D Line T₂ should be to right of line T₁															
15	A	64	<table><tr><td>Enthalpy Change</td><td>=</td><td>activation energy for Forward Reaction</td><td>-</td><td>activation energy for Reverse Reaction</td></tr><tr><td>Enthalpy Change</td><td>=</td><td>165</td><td>-</td><td>179</td></tr><tr><td>Enthalpy Change</td><td>=</td><td colspan="3">-14 kJ</td></tr></table>	Enthalpy Change	=	activation energy for Forward Reaction	-	activation energy for Reverse Reaction	Enthalpy Change	=	165	-	179	Enthalpy Change	=	-14 kJ		
Enthalpy Change	=	activation energy for Forward Reaction	-	activation energy for Reverse Reaction														
Enthalpy Change	=	165	-	179														
Enthalpy Change	=	-14 kJ																
16	D	57	☒ A High temperature will favour the endothermic reverse reaction reducing B ☒ B High activation energy decreases number of particles forming activated complex ☒ C High temperature will favour the endothermic reverse reaction reducing B ☒ D Low temperature favours forward reaction & low activation energy favours B formation															
17	D	78	c a b -a b c = -a + b N₂O₄ 2NO + O₂ 2NO + O₂ N₂O₄ 2NO + O₂ N₂O₄ → → → → → → 2NO₂ N₂O₄ 2NO₂ 2NO + O₂ 2NO₂ 2NO₂															
18	B	74	⑤ ① ② ①x2 ②x-1 ⑤ 2CH₄ CH₄ 2CO 2CH₄ 2CO₂ 2CH₄ + + + + + + 3O₂ 2O₂ O₂ 3 4 O₂ 2CO₂ 3O₂ → → → → → → 2CO CO₂ 2CO₂ 2CO₂ 2CO 2CO + + + + + 4H₂O 2H₂O 4H₂O O₂ 4H₂O															






2025 Higher Chemistry Marking Scheme

Long Qu	Answer	Reasoning																																	
1a(i)	$\text{Na(g)} \rightarrow \text{Na}^+(\text{g}) + \text{e}^-$	The first ionisation energy is the energy to remove one mole of electrons from one mole of atoms in the gaseous state.																																	
1a(ii)	1735	① $\text{Ca(g)} \longrightarrow \text{Ca}^+(\text{g}) + \text{e}^- \quad \Delta H = 590\text{kJ}$ ② $\text{Ca}^+(\text{g)} \longrightarrow \text{Ca}^{2+}(\text{g}) + \text{e}^- \quad \Delta H = 1145\text{kJ}$ ①+② $\text{Ca(g)} \longrightarrow \text{Ca}^+(\text{g}) + \text{e}^- \quad \Delta H = 590\text{kJ}$ $\text{Ca}^+(\text{g)} \longrightarrow \text{Ca}^{2+}(\text{g}) + \text{e}^- \quad \Delta H = 1145\text{kJ}$ $\text{Ca(g)} \longrightarrow \text{Ca}^{2+}(\text{g}) + 2\text{e}^- \quad \Delta H = 1735\text{kJ}$																																	
1a(iii)	Answer to include:	<table><tr><td rowspan="2">1st mark: (one from)</td><td>The second ionisation energy removes an electron from an electron shell that is</td><td>inner full (more) stable closer</td><td rowspan="2">to the nucleus</td></tr><tr><td>Second electron removed from a shell that is</td><td>inner full (more) stable closer</td><td>to the nucleus</td></tr><tr><td rowspan="2">2nd mark: (one from)</td><td>The second</td><td>electron electron shell</td><td>is less</td><td>screened shielded</td></tr><tr><td>Second electron is more strongly</td><td colspan="2">attracted to pulled towards</td><td>the nucleus</td></tr></table>	1 st mark: (one from)	The second ionisation energy removes an electron from an electron shell that is	inner full (more) stable closer	to the nucleus	Second electron removed from a shell that is	inner full (more) stable closer	to the nucleus	2 nd mark: (one from)	The second	electron electron shell	is less	screened shielded	Second electron is more strongly	attracted to pulled towards		the nucleus																	
1 st mark: (one from)	The second ionisation energy removes an electron from an electron shell that is	inner full (more) stable closer		to the nucleus																															
	Second electron removed from a shell that is	inner full (more) stable closer	to the nucleus																																
2 nd mark: (one from)	The second	electron electron shell	is less	screened shielded																															
	Second electron is more strongly	attracted to pulled towards		the nucleus																															
1a(iv)	greater nuclear charge or more protons	The larger number of protons in the nucleus of Mg^{2+} have a greater pull on the outer shell than the Na^+ nucleus has on its outer shell making the Mg^{2+} ion smaller than the Na^+																																	
1b(i)	Electrostatic attraction between positive ions and negative ions	Ionic bonding is usually between metals and non-metals where metals form positive ions and non-metals form negative ions. An ionic lattice is a three-dimensional structure of alternating positive and negative ions. Electrostatic attraction between positive and negative ions is strong and the ionic lattice is a solid with a high melting point.																																	
1b(ii)	6	$\text{mass} = \text{no. of moles} \times \text{gfm} = 0.0012 \times 39.1 = 0.0469\text{g} = 46.9\text{mg}$ $1 \text{ tablet} = 46.9\text{mg} \therefore 8 \text{ tablets} = 46.9\text{mg} \times 8 = 375\text{mg}$ $8 \text{ tablets for 1 day} = 375\text{mg} \therefore 8 \text{ tablets for 7 days} = 375\text{mg} \times 8 = 2628\text{mg}$ 450mg K ↔ 1 banana 2628mg K ↔ 1 banana $\times \frac{2628}{450}$ = 5.84 bananas = 6 whole bananas																																	
2a	<table><tr><td>metallic</td><td></td></tr><tr><td></td><td>network</td></tr><tr><td>covalent</td><td></td></tr><tr><td></td><td>molecular</td></tr></table>	metallic			network	covalent			molecular	<table><tr><td>Bonding</td><td>Structure</td><td colspan="3">Elements in 3rd Period</td></tr><tr><td>Metallic</td><td>Lattice</td><td>Sodium Na</td><td>Magnesium Mg</td><td>Aluminium Al</td></tr><tr><td>Covalent</td><td>Molecular</td><td>Phosphorus P₄</td><td>Sulphur S₈</td><td>Chlorine Cl₂</td></tr><tr><td>Covalent</td><td>Network</td><td colspan="3">Silicon Si</td></tr><tr><td></td><td>Monatomic</td><td colspan="3">Argon Ar</td></tr></table>	Bonding	Structure	Elements in 3 rd Period			Metallic	Lattice	Sodium Na	Magnesium Mg	Aluminium Al	Covalent	Molecular	Phosphorus P ₄	Sulphur S ₈	Chlorine Cl ₂	Covalent	Network	Silicon Si				Monatomic	Argon Ar		
metallic																																			
	network																																		
covalent																																			
	molecular																																		
Bonding	Structure	Elements in 3 rd Period																																	
Metallic	Lattice	Sodium Na	Magnesium Mg	Aluminium Al																															
Covalent	Molecular	Phosphorus P ₄	Sulphur S ₈	Chlorine Cl ₂																															
Covalent	Network	Silicon Si																																	
	Monatomic	Argon Ar																																	
2b(i)	decreases	Going across the third period there are a greater number of protons in the nuclei in the elements and the greater nuclear charge pulls the outer shell more which reduces the covalent radius/atomic size.																																	

2b(ii)	Argon does not form (covalent) bonds	Group 0 elements like Argon have a full outer shell so do not need to share electrons and form covalent bonds to attain a full outer shell.														
2c	Answer to include:	1 st mark:	(van der Waals/intermolecular) forces increase going down group													
		2 nd mark:	London dispersion forces are forces broken between atoms													
		3 rd mark:	The more electrons the stronger the forces													
3	Open Question Answer:	3 mark answer		2 mark answer		1 mark answer										
	<table><tr><td>Mark</td><td>0</td><td>1</td><td>2</td><td>3</td></tr><tr><td>%</td><td>26%</td><td>38%</td><td>27%</td><td>9%</td></tr></table>	Mark	0	1	2	3	%	26%	38%	27%	9%	Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.		Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.		Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
Mark	0	1	2	3												
%	26%	38%	27%	9%												
4a(i)	$\text{C}_3\text{H}_7\text{OH} + 4\frac{1}{2}\text{H}_2\text{O} \downarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$	$\text{C}_3\text{H}_7\text{OH} + 4\frac{1}{2}\text{H}_2\text{O} \longrightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$ (or any multiple)														
4a(ii)	-1050	$E_h = c m T = 4.18 \times 0.15 \times 13.9 = 8.72 \text{ kJ}$ $0.498\text{g propan-1-ol} \longleftrightarrow -8.72 \text{ kJ}$ $60\text{g propan-1-ol} \longleftrightarrow -8.72 \text{ kJ} \times \frac{60}{0.498}$ $= -1050 \text{ kJ mol}^{-1}$														
4a(iii)	Answers from:	Any four = 2 marks Any two = 1 mark	copper/metal can measuring cylinder	thermometer spirit burner & cap	clamp stand balance											
4a(iv)	One answer from	heat loss from surroundings		incomplete combustion		Evaporation of alcohol/fuel										
4b(i)	1 : 3	$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{H} \\ \quad \\ \text{OH} \quad \text{OH} \end{array}$		Formula: $\text{C}_2\text{H}_6\text{O}_2$		$\begin{array}{lcl} \text{O} & : & \text{H} \\ 2 & : & 6 \\ 1 & : & 3 \end{array}$										
4b(ii)A	Acidified dichromate or hot copper (II) oxide	Oxidising Agent	Start Colour	Final Colour	Primary Alcohol ↓ Aldehyde	Secondary Alcohol ↓ Ketone	Aldehyde ↓ Carboxylic acid									
		Acidified dichromate	orange	green	✓	✓	✓									
		hot copper (II) oxide	black	brown	✓	✓	✓									
		Fehling's Solution	blue	brick red	×	×	✓									
Tollen's Reagent	(colourless)	silver mirror	×	×	✓											
4b(ii)B	$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{O} \\ \quad \quad \quad \quad \text{OH} \end{array}$	primary alcohol $\longrightarrow$ aldehyde $\longrightarrow$ carboxylic acid butan-1-ol $\longrightarrow$ butanal $\longrightarrow$ butanoic acid $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{OH} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \end{array} \longrightarrow \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}=\text{O} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array} \longrightarrow \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}=\text{O} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{O} \\ \quad \quad \quad \quad \text{OH} \end{array}$ Does <u>not</u> turn universal indicator red Does turn universal indicator red														
4b(ii)C	butanone (accept butan-2-one)	secondary alcohol $\longrightarrow$		ketone												
		butan-2-ol $\begin{array}{c} \text{H} \quad \text{OH} \quad \text{H} \quad \text{H} \\ \quad \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \end{array}$ $\text{C}_4\text{H}_9\text{OH}$		butanone $\begin{array}{c} \text{H} \quad \text{O} \quad \text{H} \quad \text{H} \\ \quad \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{H} \\ \quad \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \end{array}$ $\text{C}_4\text{H}_8\text{O}$												
4c(i)	Volumetric flask	Volumetric flasks (also known as standard flasks) are used to measure exact volumes of liquids with a mark on the narrow neck to fill up to.														
4c(ii)A	Rinse with sodium hydroxide solution	Rinsing with sodium hydroxide solution and emptying the solution will ensure previous chemicals do not contaminate the sodium hydroxide solution and any water in the burette from previous rinsing does not dilute the sodium hydroxide.														

4c(ii)B	20.4	average titre = $\frac{20.5 + 20.3}{2} = 20.4\text{cm}^3$																																			
4c(ii)C	0.083	NaOH no. of moles = volume x concentration = 0.0198 litres x 0.105 mol l ⁻¹ = 0.002079mol NaOH : CH₃COOH 1mol 1mol 0.002079mol 0.002079mol concentration = $\frac{\text{no. of moles}}{\text{volume}} = \frac{0.002079 \text{ mol}}{0.025 \text{ litres}} = 0.083\text{mol l}^{-1}$																																			
4c(iii)A	K ⁺ CH ₃ COO ⁻	Potassium it is in group 1 of Periodic Table ∴ forms a positive ion with a 1+ charge. Ethanoate has the formula CH ₃ COO ⁻ (p9 of data booklet)																																			
4c(iii)B	5.4	no. of moles = volume x concentration = 0.2 litres x 0.45mol l ⁻¹ = 0.09mol mass = no. of moles x gfm = 0.09 x 60 = 5.4g																																			
5a	Answer to contain:	<table><tr><td>1st mark:</td><td>Sucrose is polar due to hydroxyl groups</td><td>or</td><td>Sucrose can form hydrogen bonds due to hydroxyl groups</td></tr><tr><td>2nd mark:</td><td colspan="3">polar substances like sucrose will dissolve in polar solvents like water</td></tr></table>	1 st mark:	Sucrose is polar due to hydroxyl groups	or	Sucrose can form hydrogen bonds due to hydroxyl groups	2 nd mark:	polar substances like sucrose will dissolve in polar solvents like water																													
1 st mark:	Sucrose is polar due to hydroxyl groups	or	Sucrose can form hydrogen bonds due to hydroxyl groups																																		
2 nd mark:	polar substances like sucrose will dissolve in polar solvents like water																																				
5b(i)	<table><tr><td>Iso</td><td>Pro</td><td>Gly</td><td>Pro</td><td>Phe</td><td>Pro</td><td>Tyr</td></tr><tr><td colspan="7">or</td></tr><tr><td>Tyr</td><td>Pro</td><td>Phe</td><td>Pro</td><td>Gly</td><td>Pro</td><td>Iso</td></tr></table>	Iso	Pro	Gly	Pro	Phe	Pro	Tyr	or							Tyr	Pro	Phe	Pro	Gly	Pro	Iso	There are 7 amino acids in the sequence to be worked out but 10 amino acids across the three smaller peptide chains ∴ three amino acids are 'doublers' (remember the chain could be written backwards) <table><tr><th>Look for common amino acids</th><th>Reverse Chain One</th><th>Line Up Common Fragments</th></tr><tr><td>Tyr-Pro-Phe</td><td>Phe-Pro-Tyr</td><td>Phe-Pro-Tyr</td></tr><tr><td>Gly-Pro-Phe</td><td>Gly-Pro-Phe</td><td>Gly-Pro-Phe</td></tr><tr><td>Iso-Pro-Gly-Pro</td><td>Iso-Pro-Gly-Pro</td><td>Iso-Pro-Gly-Pro</td></tr></table> Amino Acid Sequence = Iso - Pro - Gly - Pro - Phe - Pro - Tyr			Look for common amino acids	Reverse Chain One	Line Up Common Fragments	Tyr-Pro-Phe	Phe-Pro-Tyr	Phe-Pro-Tyr	Gly-Pro-Phe	Gly-Pro-Phe	Gly-Pro-Phe	Iso-Pro-Gly-Pro	Iso-Pro-Gly-Pro	Iso-Pro-Gly-Pro
Iso	Pro	Gly	Pro	Phe	Pro	Tyr																															
or																																					
Tyr	Pro	Phe	Pro	Gly	Pro	Iso																															
Look for common amino acids	Reverse Chain One	Line Up Common Fragments																																			
Tyr-Pro-Phe	Phe-Pro-Tyr	Phe-Pro-Tyr																																			
Gly-Pro-Phe	Gly-Pro-Phe	Gly-Pro-Phe																																			
Iso-Pro-Gly-Pro	Iso-Pro-Gly-Pro	Iso-Pro-Gly-Pro																																			
5b(ii)	Essential	Essential amino acids are amino acids which the body cannot make for itself and must be obtained by the body through their diet.																																			
5c(i)	Esters (or triglyceride)	Fats and oils are made from the condensation reaction between glycerol and three fatty acids. This makes fats and oils which are esters.																																			
5c(ii)A	propane-1,2,3-triol or glycerol		propane-1,2,3-triol 3 carbons Single bonds between carbons Functional groups on Carbons 1,2,3 3 hydroxyl -OH groups																																		
5c(ii)B	Palmitic Acid	<table><tr><th>Fatty Acid</th><th>Formula</th><th>Fits General Formula</th><th>Number of C=C Double Bonds</th></tr><tr><td>Palmitic Acid</td><td>C₁₅H₃₁COOH</td><td>C_nH_{2n+1}COOH</td><td>0</td></tr><tr><td>Oleic Acid</td><td>C₁₇H₃₃COOH</td><td>C_nH_{2n-1}COOH</td><td>1</td></tr><tr><td>Linoleic Acid</td><td>C₁₇H₃₁COOH</td><td>C_nH_{2n-3}COOH</td><td>2</td></tr><tr><td>Linolenic Acid</td><td>C₁₇H₂₉COOH</td><td>C_nH_{2n-5}COOH</td><td>3</td></tr></table>			Fatty Acid	Formula	Fits General Formula	Number of C=C Double Bonds	Palmitic Acid	C ₁₅ H ₃₁ COOH	C _n H _{2n+1} COOH	0	Oleic Acid	C ₁₇ H ₃₃ COOH	C _n H _{2n-1} COOH	1	Linoleic Acid	C ₁₇ H ₃₁ COOH	C _n H _{2n-3} COOH	2	Linolenic Acid	C ₁₇ H ₂₉ COOH	C _n H _{2n-5} COOH	3													
Fatty Acid	Formula	Fits General Formula	Number of C=C Double Bonds																																		
Palmitic Acid	C ₁₅ H ₃₁ COOH	C _n H _{2n+1} COOH	0																																		
Oleic Acid	C ₁₇ H ₃₃ COOH	C _n H _{2n-1} COOH	1																																		
Linoleic Acid	C ₁₇ H ₃₁ COOH	C _n H _{2n-3} COOH	2																																		
Linolenic Acid	C ₁₇ H ₂₉ COOH	C _n H _{2n-5} COOH	3																																		
5c(iii)	Addition (accept reduction)	An addition reaction happens when a molecule adds across a C=C double bond. - an addition reaction where hydrogen is added across a C=C double bond is called hydrogenation - an addition reaction where water is added across a C=C double bond is called hydration																																			

8b(i)	filtration	Solid magnesium hydroxide can be removed from calcium chloride solution by filtration			
8b(ii)	One answer from:	Seawater is a free/cheap raw material		Sell calcium chloride as by-product	
9a(i)	Propagation	Step	Reactants (before Arrow)	→	Products (after Arrow)
		Initiation	No free radicals on Reactant Side	→	Free radicals on Product Side
		Propagation	Free Radicals found on both sides of arrow		
		Termination	Free radicals on Reactant Side	→	No free radicals on Product Side
9a(ii)	Unpaired electron	A free radical is a particles containing an unpaired electron.			
9b	Denaturing	Proteins are denatured when the shape of the protein is permanently changed often due to a rise in temperature or a change in pH.			
9c	One answer from:	plastics	food products	cosmetics	or any other reasonable answer
9d(i)	Oxygen/gas is escaping	The gas produced in the chemical reaction is able to escape the test tube as there is no stopper and the balance will record this loss of mass.			
9d(ii)	Two answers from:	concentration of hydrogen peroxide	temperature	mass volume surface area size	} of potato discs
9d(iii)	One answer from:	As pH increases { the rate increases up to pH=10 the rate reaches a maximum at pH=10 the rate reaches an optimum at pH=10 } (and then decreases) or As the pH increases the rate increases and then decreases			
10a(i)	O H -C- -N-	O H -C- -N- Amide link (also called peptide link in proteins only) O -C- -O- Ester link			
10a(ii)	One answer from:	Bromine solution would { quickly rapidly } decolourise with { CAP capsaicin			
		Bromine solution will decolourise with { CAP capsaicin } but not with { DHC dihydrocapsaicin			
		Bromine solution not decolourise with { DHC dihydrocapsaicin } but will with { CAP capsaicin			
10b	One answer from:	Capsaicinoids do not dissolve in water	Capsaicinoids are insoluble in water	Capsaicinoids are hydrophobic	Capsaicinoids are non-polar
10c	135	CAP has retention time of 2.25 minutes = 135seconds			
10d	3	0.025% of 60g tube = $\frac{0.025}{100} \times 60\text{g} = 0.015\text{g CAP}$ 1000g CAP ↔ £1930.32 0.015g CAP ↔ $\text{£}1930.32 \times \frac{0.015\text{g}}{1000}$ = £0.029 = 2.9p = 3p			

10e	$\begin{array}{c} \text{C}_6\text{H}_8\text{O}_6 \\ \downarrow \\ \text{C}_6\text{H}_6\text{O}_6 + 2\text{H}^+ + 2\text{e}^- \end{array}$	<u>Step 1:</u> Write down main species in reaction $\text{C}_6\text{H}_8\text{O}_6 \rightarrow \text{C}_6\text{H}_6\text{O}_6$ <u>Step 2:</u> Balance all atoms other than O or H $\text{C}_6\text{H}_8\text{O}_6 \rightarrow \text{C}_6\text{H}_6\text{O}_6$ <u>Step 3:</u> Balance O atoms by adding H_2O to the other side $\text{C}_6\text{H}_8\text{O}_6 \rightarrow \text{C}_6\text{H}_6\text{O}_6$ <u>Step 4:</u> Balance H atoms by adding H^+ to the other side $\text{C}_6\text{H}_8\text{O}_6 \rightarrow \text{C}_6\text{H}_6\text{O}_6 + 2\text{H}^+$ <u>Step 5:</u> Balance charge by adding electrons to the most positive side $\text{C}_6\text{H}_8\text{O}_6 \rightarrow \text{C}_6\text{H}_6\text{O}_6 + 2\text{H}^+ + 2\text{e}^-$												
11	Open Question Answer: <table border="1"> <tr> <td>Mark</td><td>0</td><td>1</td><td>2</td><td>3</td></tr> <tr> <td>%</td><td>57%</td><td>23%</td><td>14%</td><td>7%</td></tr> </table>	Mark	0	1	2	3	%	57%	23%	14%	7%	3 mark answer Demonstrates a <u>good</u> understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	2 mark answer Demonstrates a <u>reasonable</u> understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	1 mark answer Demonstrates a <u>limited</u> understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
Mark	0	1	2	3										
%	57%	23%	14%	7%										
12a	2,7-dimethyloctane													
12b														
12c														

2025 Higher Chemistry Section 1 Multiple Choice Analysis

Item	Key	A	B	C	D	Missing	Invalid
1	A	87	5	4	4	0	0
2	C	2	10	77	12	0	0
3	B	15	72	8	4	1	0
4	D	4	38	4	53	0	0
5	B	2	90	7	1	0	0
6	A	93	2	4	2	0	0
7	B	6	69	17	8	0	0
8	D	12	13	11	64	1	0
9	C	6	12	74	7	1	0
10	B	5	72	7	15	0	0
11	D	11	2	5	82	0	0
12	B	7	84	7	2	0	0
13	C	3	10	79	8	0	0
14	A	71	3	18	8	0	0
15	A	64	29	4	3	0	0
16	D	18	16	8	57	0	0
17	D	3	16	3	78	0	0
18	B	2	74	4	20	1	0
19	D	5	7	6	82	0	0
20	C	17	28	46	8	0	0
21	D	14	20	22	43	1	0
22	A	62	18	3	17	0	0
23	C	22	19	36	21	1	0
24	C	20	6	62	12	0	0
25	A	69	26	3	2	0	0

2025 Higher Chemistry Section 2 Marks Data Analysis

2025 Item	Max Mark	Average Mark	% Average Mark	Discrimination	% Not Attempted
Q1ai	1	0.8	82.4%	29	0.4%
Q1aii	1	0.8	75.5%	40	1.2%
Q1aiii	2	0.7	36.1%	43	0.9%
Q1aiv	1	0.6	58.2%	45	2.5%
Q1bi	1	0.1	14.3%	29	2.7%
Q1bii	2	1.3	64.1%	47	1.2%
Q2a	2	1.4	70.1%	38	0.6%
Q2bi	1	0.9	89.2%	31	0.3%
Q2bii	1	0.5	48.4%	38	1.5%
Q2c	3	2.2	74.3%	54	1.8%
Q3	3	1.2	39.5%	62	9.3%
Q4ai	1	0.4	42.0%	35	1.3%
Q4aii	3	1.7	56.4%	62	2.2%
Q4aiii	2	1	47.9%	40	1.0%
Q4aiv	1	0.8	75.2%	49	2.6%
Q4bi	1	0.8	80.4%	48	2.0%
Q4biiA	1	0.4	45.0%	42	3.7%
Q4biiB	1	0.7	68.3%	59	8.3%
Q4biiC	1	0.5	52.5%	55	4.5%
Q4ci	1	0.5	45.7%	39	0.5%
Q4ciiA	1	0.7	65.9%	40	1.7%
Q4ciiB	1	0.7	68.3%	35	0.5%
Q4ciiC	3	2.3	78.1%	54	2.3%
Q4ciiiA	1	0.4	43.8%	42	8.0%
Q4ciiiB	1	0.8	79.7%	47	1.8%
Q5a	2	1.1	54.6%	56	1.6%
Q5bi	1	0.3	26.7%	40	8.3%
Q5bii	1	0.9	92.9%	40	2.3%
Q5ci	1	0.6	59.4%	37	3.4%
Q5ciiA	1	0.8	79.2%	43	1.3%
Q5ciiB	1	0.7	71.4%	33	2.1%
Q5ciii	1	0.6	59.5%	44	1.6%
Q5di	1	0.7	67.6%	52	1.8%
Q5dii	1	0.8	83.5%	48	0.8%
Q5diiiA	1	0.8	81.9%	39	1.2%
Q5diiiB	1	0.6	60.8%	38	6.5%
Q5e	1	0.9	85.5%	30	1.7%
Q6a	1	0.5	47.4%	50	7.1%
Q6b	1	0.8	76.9%	37	2.9%
Q6ci	1	1	98.3%	10	0.6%
Q6cii	1	0.3	31.0%	26	0.4%
Q6di	1	0.7	65.3%	34	0.7%
Q6dii	1	0.7	65.5%	42	0.9%
Q6diii	2	1.2	61.5%	43	1.4%
Q7a	2	1.3	63.7%	53	2.9%
Q7b	2	0.7	33.9%	54	4.1%
Q7ci	2	0.8	40.5%	50	8.1%
Q7cii	2	1.8	88.5%	41	1.4%
Q8ai	2	0.9	44.1%	55	6.1%
Q8aii	2	1.5	77.3%	55	3.4%
Q8aiii	2	1.5	76.6%	59	5.8%
Q8bi	1	0.5	54.8%	34	3.7%
Q8bii	1	0.3	25.5%	27	8.5%
Q9ai	1	0.9	92.1%	40	3.2%
Q9aii	1	0.5	50.3%	41	3.1%
Q9b	1	0.9	92.1%	37	2.8%
Q9c	1	0.8	84.0%	31	2.9%
Q9di	1	0.5	48.1%	44	3.1%
Q9dii	2	1.4	70.0%	30	1.0%
Q9diii	1	0.1	7.6%	20	2.8%
Q10ai	1	0.9	86.8%	36	1.4%
Q10aii	1	0.5	46.7%	50	6.0%
Q10b	1	0.5	47.3%	13	4.3%
Q10c	1	0.5	53.6%	32	1.6%
Q10d	2	1	48.2%	45	3.1%
Q10e	1	0.5	52.0%	54	8.1%
Q11	3	0.7	23.4%	55	20.1%
Q12a	1	0.6	61.1%	41	1.9%
Q12b	1	0.7	66.2%	28	2.1%
Q12c	1	0.2	22.9%	21	3.2%
Paper	95	56.5	59.5%		

2025 Higher Chemistry Section 2 Frequency of Marks Analysis

2025 Item	0	1	2	3
Q1ai	18	82		
Q1aii	24	76		
Q1aiii	41	46	13	
Q1aiv	42	58		
Q1bi	86	14		
Q1bii	24	23	52	
Q2a	12	36	52	
Q2bi	11	89		
Q2bii	52	48		
Q2c	9	12	27	53
Q3	26	38	27	9
Q4ai	58	42		
Q4aii	21	21	25	33
Q4aiii	28	49	23	
Q4aiv	25	75		
Q4bi	20	80		
Q4biiA	55	45		
Q4biiB	32	68		
Q4biiC	47	53		
Q4ci	54	46		
Q4ciiA	34	66		
Q4ciiB	32	68		
Q4ciiC	11	11	10	68
Q4ciiiA	56	44		
Q4ciiiB	20	80		
Q5a	27	37	36	
Q5bi	73	27		
Q5bii	7	93		
Q5ci	41	59		
Q5ciiA	21	79		
Q5ciiB	29	71		
Q5ciii	41	59		
Q5di	32	68		
Q5dii	17	83		
Q5diiiA	18	82		
Q5diiiB	39	61		
Q5e	15	85		
Q6a	53	47		
Q6b	23	77		
Q6ci	2	98		
Q6cii	69	31		
Q6di	35	65		
Q6dii	35	65		
Q6diii	20	38	43	
Q7a	22	30	49	
Q7b	36	60	4	
Q7ci	46	27	27	
Q7cii	8	7	85	
Q8ai	35	43	23	
Q8aii	13	19	68	
Q8aiii	18	10	71	
Q8bi	45	55		
Q8bii	74	26		
Q9ai	8	92		
Q9aii	50	50		
Q9b	8	92		
Q9c	16	84		
Q9di	52	48		
Q9dii	12	36	52	
Q9diii	92	8		
Q10ai	13	87		
Q10aii	53	47		
Q10b	53	47		
Q10c	46	54		
Q10d	30	44	26	
Q10e	48	52		
Q11	57	23	14	7
Q12a	39	61		
Q12b	34	66		
Q12c	77	23		

2025 Higher Chemistry Section 3 Assignment Marks Analysis

Item	Max Mark	Average Mark	% Average Mark	Discrimination
1	1	1	98.3%	14
2	3	1.7	55.1%	43
3a	1	0.3	34.3%	30
3b	1	1	95.6%	10
3c	1	0.6	64.1%	31
3d	1	0.6	60.8%	30
3e	1	1	97.1%	19
3f	1	0.6	57.5%	31
4a	1	1	95.9%	23
4b	1	0.9	88.0%	21
4c	1	0.9	87.3%	26
4d	1	0.7	68.4%	29
5	1	0.3	25.5%	35
6	1	0.6	63.3%	21
7	3	1	32.7%	44
8	1	1	97.6%	13
Paper	20	13	64.8%	

Item	Max Mark	0	1	2	3
1	1	2	98		
2	3	3	42	43	12
3a	1	66	34		
3b	1	4	96		
3c	1	36	64		
3d	1	39	61		
3e	1	3	97		
3f	1	42	58		
4a	1	4	96		
4b	1	12	88		
4c	1	13	87		
4d	1	32	68		
5	1	75	25		
6	1	37	63		
7	3	33	41	20	6
8	1	2	98		
Paper	20				

Item	Max Mark	Item Description
1	1	An aim that describes clearly the purpose of the investigation
2	3	Underlying chemistry
3a	1	A brief summary of the approach used to collect experimental data.
3b	1	Sufficient raw data from the candidate's experiment.
3c	1	Data presented in correctly produced table(s) and correct units shown for calculated values
3d	1	Values calculated correctly using a chemical relationship
3e	1	Data relevant to the experiment obtained from an internet/literature source
3f	1	A citation and reference for a source of the internet/literature data
4a	1	An appropriate format from the options of scatter graph, line graph or bar graph
4b	1	The axis/axes of the graph has/have suitable scale(s).
4c	1	The axes of the graph have suitable labels and units
4d	1	Accurately plotted data points and, where appropriate, a line of best fit.
5	1	A valid comparison of the experimental data with data from the internet/literature source
6	1	A valid conclusion that relates to the aim and is supported by all the data in the report.
7	3	A valid conclusion that relates to the aim and is supported by all the data in the report.
8	1	A valid conclusion that relates to the aim and is supported by all the data in the report.