



JABchem



Not to be shared without the copyright holder's permission

Past Papers Advanced Higher Chemistry



2022 Marking Scheme

Grade Awarded	Mark Required		% candidates achieving grade
	(/130)	%	
A	+	%	%
B	+	%	%
C	+	%	%
D	+	%	%
No award	<	<%	%

























Section:	Multiple Choice	Extended Answer	Project
Average Mark:	/25	/70	No Project in 2022

2022 Adv Higher Chemistry Marking Scheme

















MC Qu	Answer	% Pupils Correct	Reasoning															
1	B		<input checked="" type="checkbox"/> A Each element does have a characteristic spectrum <input checked="" type="checkbox"/> B Heat energy is used to promote electrons in emission spectroscopy <input checked="" type="checkbox"/> C Lines are formed by electrons releasing energy as they drop down energy levels <input checked="" type="checkbox"/> D More electrons dropping down levels leads to more intense lines formed															
2	D		Cr atom electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^1$ Cr ⁺ ion electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5$															
3	D		Octahedral complex ion means M ²⁺ ion makes 6 dative covalent bonds with ligands. <ul style="list-style-type: none">• L⁻ is a bidentate ligand making 2 dative covalent bonds with M²⁺• M²⁺ pairs up with 3 L⁻ bidentate ligands• Formula of complex ion = [ML₃]⁻															
4	A		<input checked="" type="checkbox"/> A homogeneous catalysts are in the same state as the reactants <input checked="" type="checkbox"/> B Adsorption of reactive molecules must take place in heterogeneous catalysis <input checked="" type="checkbox"/> C A catalyst will reduce the activation energy for a reaction pathway <input checked="" type="checkbox"/> D catalysts are often transition metals due to unpaired d electrons in mechanism															
5	D		<input checked="" type="checkbox"/> A Cl ⁻ is reduced to Cl ⁻ and I ⁻ is oxidised to I ₂ but are different substance <input checked="" type="checkbox"/> B IO ₃ ⁻ is reduced to I ₂ and I ⁻ is oxidised to I ₂ but are different substances <input checked="" type="checkbox"/> C Cl ₂ is reduced to Cl ⁻ and I ₂ is oxidised to IO ₃ ⁻ but are different substances <input checked="" type="checkbox"/> D Cl ₂ is reduced to Cl ⁻ and at the same time Cl ₂ is oxidised to ClO ⁻ <table><tr><td>Reaction</td><td colspan="2">Reduction reaction</td><td colspan="2">Oxidation Reaction</td></tr><tr><td>Equation</td><td colspan="2">$Cl_2 + 2e^- \rightarrow 2Cl^-$</td><td colspan="2">$Cl_2 + 2H_2O \rightarrow 2ClO^- + 4H^+ + 2e^-$</td></tr><tr><td>Oxidation State</td><td>0</td><td>-1</td><td>0</td><td>+1</td></tr></table>	Reaction	Reduction reaction		Oxidation Reaction		Equation	$Cl_2 + 2e^- \rightarrow 2Cl^-$		$Cl_2 + 2H_2O \rightarrow 2ClO^- + 4H^+ + 2e^-$		Oxidation State	0	-1	0	+1
Reaction	Reduction reaction		Oxidation Reaction															
Equation	$Cl_2 + 2e^- \rightarrow 2Cl^-$		$Cl_2 + 2H_2O \rightarrow 2ClO^- + 4H^+ + 2e^-$															
Oxidation State	0	-1	0	+1														
6	C		<input checked="" type="checkbox"/> A pH = 7.00 is true at 25°C (298K) <input checked="" type="checkbox"/> B The ionic product $K_w=[H_3O^+][OH^-] = 1.01 \times 10^{-14}$ is only true at 25°C (298K) <input checked="" type="checkbox"/> C $[H_3O^+] = [OH^-]$ is true for any neutral solution at any temperature <input checked="" type="checkbox"/> D $[H_3O^+] = 1.00 \times 10^{-7}$ at 25°C (298K)															
7	A		<input checked="" type="checkbox"/> A [SO ₃] will increase with an increase in pressure but value of K is unchanged <input checked="" type="checkbox"/> B Changes to pressure do not change the value of the equilibrium constant K <input checked="" type="checkbox"/> C Increase in pressure will favour pressure-reducing forward reaction ∴ ↑ [SO ₃] <input checked="" type="checkbox"/> D Increase in pressure will favour pressure-reducing forward reaction ∴ ↑ [SO ₃]															
8	D		enthalpy of formation is the energy change for the formation of 1mol of a substance from its elements in their natural state. <input checked="" type="checkbox"/> A Reactants are ions and not elements in their natural state <input checked="" type="checkbox"/> B Reactants are ions and not elements in their natural state <input checked="" type="checkbox"/> C Cl _(g) is not the natural state of chlorine Cl _{2(g)} <input checked="" type="checkbox"/> D Reactants are elements in their natural state and 1 mole of product is formed															
9	B		Rate = $k[A]^2[B]$ ∴ [A] ² means A is 2nd order (caused by having two particles of A in rate determining step) ∴ [B] ¹ means B is 1st order (caused by having one particles of B in rate determining step) <input checked="" type="checkbox"/> A Doubling [A] leads to a quadrupling of reaction rate as A is 2 nd order <input checked="" type="checkbox"/> B Doubling [B] leads to a doubling of reaction rate as B is 1 st order <input checked="" type="checkbox"/> C Doubling [B] leads to a doubling of reaction rate as B is 1 st order <input checked="" type="checkbox"/> D Doubling [A] and [B] would lead to a combined quadrupling of reaction rate.															
10	D		<input checked="" type="checkbox"/> A The order of each reactant is not decided by stoichiometric coefficients <input checked="" type="checkbox"/> B The order of a reaction is decided by the slow rate determining step <input checked="" type="checkbox"/> C The speed of reaction is determined by the slowest step - rate determining step <input checked="" type="checkbox"/> D The individual orders are determined by experimentally altering concentrations and looking to see the effect on the reaction rate.															






















19	B	 <p>A chiral carbon is a carbon with four different groups attached. This leads to non-superimposable mirror images in three dimensions</p> 															
20	C	<div> <div> <input checked="" type="checkbox"/> A Formula C_4H_4 <input checked="" type="checkbox"/> B Formula C_4H_6 <input checked="" type="checkbox"/> C Formula C_4H_8 fits empirical formula CH_2 <input checked="" type="checkbox"/> D Formula C_4H_{10} </div> <table border="1"> <thead> <tr> <th>Element</th><th>C</th><th>H</th></tr> </thead> <tbody> <tr> <td>Mass or %</td><td>0.12g</td><td>0.02g</td></tr> <tr> <td>No. of moles (divide % by gfm)</td><td>$\frac{0.12}{12} = 0.01$</td><td>$\frac{0.02}{1} = 0.02$</td></tr> <tr> <td>Mole ratio (divide through by smallest value)</td><td>$\frac{0.01}{0.01} = 1$</td><td>$\frac{0.02}{0.01} = 2$</td></tr> <tr> <td>Empirical Formula</td><td colspan="2">CH_2</td></tr> </tbody> </table> </div>	Element	C	H	Mass or %	0.12g	0.02g	No. of moles (divide % by gfm)	$\frac{0.12}{12} = 0.01$	$\frac{0.02}{1} = 0.02$	Mole ratio (divide through by smallest value)	$\frac{0.01}{0.01} = 1$	$\frac{0.02}{0.01} = 2$	Empirical Formula	CH_2	
Element	C	H															
Mass or %	0.12g	0.02g															
No. of moles (divide % by gfm)	$\frac{0.12}{12} = 0.01$	$\frac{0.02}{1} = 0.02$															
Mole ratio (divide through by smallest value)	$\frac{0.01}{0.01} = 1$	$\frac{0.02}{0.01} = 2$															
Empirical Formula	CH_2																
21	B	<input checked="" type="checkbox"/> A $C_3H_8O_2$ has gfm = $(3 \times 12) + (8 \times 1) + (2 \times 16) = 36 + 8 + 32 = 76$ <input checked="" type="checkbox"/> B $C_3H_6O_2$ has gfm = $(3 \times 12) + (6 \times 1) + (2 \times 16) = 36 + 6 + 32 = 74$ <input checked="" type="checkbox"/> C C_3H_8O has gfm = $(3 \times 12) + (8 \times 1) + (1 \times 16) = 36 + 8 + 16 = 72$ <input checked="" type="checkbox"/> D C_3H_8O has gfm = $(3 \times 12) + (8 \times 1) + (1 \times 16) = 36 + 8 + 16 = 72$															
22	A	$\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{6.7}{134} = 0.05 \text{ mol}$ <p>sodium carbonate + malic acid \longrightarrow sodium malate + water + carbon dioxide</p> $\text{Na}_2\text{CO}_3 + \text{C}_4\text{H}_6\text{O}_5 \longrightarrow \text{Na}_2\text{C}_4\text{H}_4\text{O}_5 + \text{H}_2\text{O} + \text{CO}_2$ <p>1mol 1mol 0.05mol 0.05mol</p>															
23	B	$\text{gfm CaCl}_2 \cdot 2\text{H}_2\text{O} = (1 \times 40.1) + (2 \times 35.5) + (4 \times 1) + (2 \times 16) = 40.1 + 71 + 4 + 32 = 147.1 \text{ g}$ $\text{gfm CaCl}_2 = (1 \times 40.1) + (2 \times 35.5) = 40.1 + 71 = 111.1 \text{ g}$ $\text{CaCl}_2 \cdot 2\text{H}_2\text{O} \longrightarrow \text{CaCl}_2 + 2\text{H}_2\text{O}$ <p>1mol 1mol 147.1g 111.1g 2.52g $111.1 \text{ g} \times \frac{2.52}{147.1}$ = 1.90g</p>															
24	A	<input checked="" type="checkbox"/> A Distillation will identify the boiling point of compound while purifying compound <input checked="" type="checkbox"/> B Recrystallisation will purify a compound but not help identify the compound <input checked="" type="checkbox"/> C Solvent Extraction will purify a compound but not help identify the compound <input checked="" type="checkbox"/> D Melting Point Determination will identify a compound but not purify compound															
25	B	<input checked="" type="checkbox"/> A Impurities could move same distance as spot <input checked="" type="checkbox"/> B There is no reactant left (R) and only the product spot (S) is present <input checked="" type="checkbox"/> C More than one chemical could be present in the spot at (S) <input checked="" type="checkbox"/> D Some reactions at equilibrium never reach completion															




2022 Adv Higher Chemistry Marking Scheme

Long Qu	Answer	Reasoning																
1a(i)	∞ figure of 8 shape	<table><tr><th>Subshell Type</th><td>s</td><td>p</td><td>d</td></tr><tr><th>Angular Momentum Quantum number (l)</th><td>$l=0$</td><td>$l=1$</td><td>$l=2$</td></tr><tr><th>Shape(s)</th><td></td><td></td><td> or </td></tr></table>	Subshell Type	s	p	d	Angular Momentum Quantum number (l)	$l=0$	$l=1$	$l=2$	Shape(s)			 or 				
Subshell Type	s	p	d															
Angular Momentum Quantum number (l)	$l=0$	$l=1$	$l=2$															
Shape(s)			 or 															
1a(ii)	<table><tr><td>n</td><td>2</td></tr><tr><td>l</td><td>1</td></tr><tr><td>m_l</td><td>-1 or 0 or +1</td></tr><tr><td>m_s</td><td>$+\frac{1}{2}$ or $-\frac{1}{2}$</td></tr></table>	n	2	l	1	m_l	-1 or 0 or +1	m_s	$+\frac{1}{2}$ or $-\frac{1}{2}$	<table><tr><td>n</td><td>2p subshell is in 2nd shell and has a value for Principal quantum number (n) of n=2</td></tr><tr><td>l</td><td>p subshells have an angular momentum quantum number (l) of l =1</td></tr><tr><td>m_l</td><td>p subshells have a magnetic quantum number (m_l) of -l to +l \therefore -1, 0, +1</td></tr><tr><td>m_s</td><td>All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$</td></tr></table>	n	2p subshell is in 2 nd shell and has a value for Principal quantum number (n) of n=2	l	p subshells have an angular momentum quantum number (l) of l =1	m_l	p subshells have a magnetic quantum number (m_l) of -l to +l \therefore -1, 0, +1	m_s	All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$
n	2																	
l	1																	
m_l	-1 or 0 or +1																	
m_s	$+\frac{1}{2}$ or $-\frac{1}{2}$																	
n	2p subshell is in 2 nd shell and has a value for Principal quantum number (n) of n=2																	
l	p subshells have an angular momentum quantum number (l) of l =1																	
m_l	p subshells have a magnetic quantum number (m_l) of -l to +l \therefore -1, 0, +1																	
m_s	All electrons either have a spin magnetic quantum number of either $+\frac{1}{2}$ or $-\frac{1}{2}$																	
1b	Removing electron from oxygen creates a half-filled 2p orbital requires less energy as a half-filled p subshell is more stable. Or Removing an electron from nitrogen breaks a more stable half-filled 2p orbital which required more energy to break	<table><tr><th>Element</th><th>Nitrogen N</th><th>Oxygen O</th></tr><tr><td>Electronic configuration before electron removal</td><td>$1s^2 2s^2 2p^3$ </td><td>$1s^2 2s^2 2p^4$ </td></tr><tr><td>Electronic configuration after electron removal</td><td>$1s^2 2s^2 2p^2$ </td><td>$1s^2 2s^2 2p^3$ </td></tr><tr><td>Reasoning</td><td>Nitrogen atom has a half-filled 2p orbital and this is more stable electron configuration than the N⁺ ion with 2 electrons in the 2p orbital. More energy required to remove an electron from a half-filled subshell.</td><td>Oxygen atom has 4 electrons in 2p orbitals and losing an electron to become O⁺ ion creates the more stable half-filled subshell. Less energy is required to create a half-filled subshell by removing an electron.</td></tr></table>	Element	Nitrogen N	Oxygen O	Electronic configuration before electron removal	$1s^2 2s^2 2p^3$ 	$1s^2 2s^2 2p^4$ 	Electronic configuration after electron removal	$1s^2 2s^2 2p^2$ 	$1s^2 2s^2 2p^3$ 	Reasoning	Nitrogen atom has a half-filled 2p orbital and this is more stable electron configuration than the N ⁺ ion with 2 electrons in the 2p orbital. More energy required to remove an electron from a half-filled subshell.	Oxygen atom has 4 electrons in 2p orbitals and losing an electron to become O ⁺ ion creates the more stable half-filled subshell. Less energy is required to create a half-filled subshell by removing an electron.				
Element	Nitrogen N	Oxygen O																
Electronic configuration before electron removal	$1s^2 2s^2 2p^3$ 	$1s^2 2s^2 2p^4$ 																
Electronic configuration after electron removal	$1s^2 2s^2 2p^2$ 	$1s^2 2s^2 2p^3$ 																
Reasoning	Nitrogen atom has a half-filled 2p orbital and this is more stable electron configuration than the N ⁺ ion with 2 electrons in the 2p orbital. More energy required to remove an electron from a half-filled subshell.	Oxygen atom has 4 electrons in 2p orbitals and losing an electron to become O ⁺ ion creates the more stable half-filled subshell. Less energy is required to create a half-filled subshell by removing an electron.																
1c	1.05×10^{-16}	$E = R \times Z^2 \times \left(1 - \frac{1}{n^2}\right)$ $= 2.18 \times 10^{-18} \times (8)^2 \times \left(1 - \frac{1}{2^2}\right)$ $= 2.18 \times 10^{-18} \times 64 \times (1 - 0.25)$ $= 2.18 \times 10^{-18} \times 64 \times (0.75)$ $= 1.0464 \times 10^{-16} \text{ J}$																
2a	Reaction not feasible at 298K as $\Delta G > 0$	<table><tr><td>$\Delta H^\circ = \Sigma \Delta H_f^\circ(\text{products}) - \Sigma \Delta H_f^\circ(\text{reactants})$ $= (1 \times 90.3) - \left(\frac{1}{2} \times 0\right) + \left(\frac{1}{2} \times 0\right)$ $= 90.3$ $= 90.3 \text{ kJ mol}^{-1}$</td><td>$\Delta S^\circ = \Sigma S^\circ(\text{products}) - \Sigma S^\circ(\text{reactants})$ $= (1 \times 211) - \left(\frac{1}{2} \times 192\right) + \left(\frac{1}{2} \times 205\right)$ $= 211 - (96 + 102.5)$ $= 211 - 198.5$ $= 12.5 \text{ J K}^{-1} \text{ mol}^{-1}$</td></tr><tr><td colspan="2">$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = 90.3 - \left(298 \times \frac{12.5}{1000}\right) = 90.3 - (3.725) = +86.6 \text{ kJ mol}^{-1}$</td></tr></table>	$\Delta H^\circ = \Sigma \Delta H_f^\circ(\text{products}) - \Sigma \Delta H_f^\circ(\text{reactants})$ $= (1 \times 90.3) - \left(\frac{1}{2} \times 0\right) + \left(\frac{1}{2} \times 0\right)$ $= 90.3$ $= 90.3 \text{ kJ mol}^{-1}$	$\Delta S^\circ = \Sigma S^\circ(\text{products}) - \Sigma S^\circ(\text{reactants})$ $= (1 \times 211) - \left(\frac{1}{2} \times 192\right) + \left(\frac{1}{2} \times 205\right)$ $= 211 - (96 + 102.5)$ $= 211 - 198.5$ $= 12.5 \text{ J K}^{-1} \text{ mol}^{-1}$	$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = 90.3 - \left(298 \times \frac{12.5}{1000}\right) = 90.3 - (3.725) = +86.6 \text{ kJ mol}^{-1}$													
$\Delta H^\circ = \Sigma \Delta H_f^\circ(\text{products}) - \Sigma \Delta H_f^\circ(\text{reactants})$ $= (1 \times 90.3) - \left(\frac{1}{2} \times 0\right) + \left(\frac{1}{2} \times 0\right)$ $= 90.3$ $= 90.3 \text{ kJ mol}^{-1}$	$\Delta S^\circ = \Sigma S^\circ(\text{products}) - \Sigma S^\circ(\text{reactants})$ $= (1 \times 211) - \left(\frac{1}{2} \times 192\right) + \left(\frac{1}{2} \times 205\right)$ $= 211 - (96 + 102.5)$ $= 211 - 198.5$ $= 12.5 \text{ J K}^{-1} \text{ mol}^{-1}$																	
$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = 90.3 - \left(298 \times \frac{12.5}{1000}\right) = 90.3 - (3.725) = +86.6 \text{ kJ mol}^{-1}$																		
2b	600	5 litres per minute = 300 litres per hour 2ppm = 2mg per litre 1litre = 2mg 300 litres = $2\text{mg} \times \frac{300}{1}$ = 600 mg																
2c	<table><tr><td>+2</td></tr><tr><td>+4</td></tr><tr><td>+3</td></tr></table>	+2	+4	+3	<table><tr><td>NO</td><td>total oxidation state from O = $1 \times -2 = -2$</td><td>\therefore Oxidation state of N = +2</td></tr><tr><td>NO₂</td><td>total oxidation state from O = $2 \times -2 = -4$</td><td>\therefore Oxidation state of N = +4</td></tr><tr><td>N₂O₃</td><td>total oxidation state from O = $3 \times -2 = -6$</td><td>\therefore Oxidation State of N = $+\frac{6}{2} = +3$</td></tr></table>	NO	total oxidation state from O = $1 \times -2 = -2$	\therefore Oxidation state of N = +2	NO ₂	total oxidation state from O = $2 \times -2 = -4$	\therefore Oxidation state of N = +4	N ₂ O ₃	total oxidation state from O = $3 \times -2 = -6$	\therefore Oxidation State of N = $+\frac{6}{2} = +3$				
+2																		
+4																		
+3																		
NO	total oxidation state from O = $1 \times -2 = -2$	\therefore Oxidation state of N = +2																
NO ₂	total oxidation state from O = $2 \times -2 = -4$	\therefore Oxidation state of N = +4																
N ₂ O ₃	total oxidation state from O = $3 \times -2 = -6$	\therefore Oxidation State of N = $+\frac{6}{2} = +3$																



2d(i)A	Rate = k [NO] ² [H ₂]	<table><tr><td>no. of mol in slow step</td><td>Order of Reactant</td><td>Effect on Reactant</td><td>Rate Equation</td></tr><tr><td>2mol NO</td><td>2nd order</td><td>[NO]²</td><td rowspan="2">Rate = k [NO]² [H₂]</td></tr><tr><td>1mol H₂</td><td>1st order</td><td>[H₂]</td></tr></table>	no. of mol in slow step	Order of Reactant	Effect on Reactant	Rate Equation	2mol NO	2 nd order	[NO] ²	Rate = k [NO] ² [H ₂]	1mol H ₂	1 st order	[H ₂]																
no. of mol in slow step	Order of Reactant	Effect on Reactant	Rate Equation																										
2mol NO	2 nd order	[NO] ²	Rate = k [NO] ² [H ₂]																										
1mol H ₂	1 st order	[H ₂]																											
2d(i)B	9.94x10 ⁻³	$[\text{NO}]^2 = \frac{\text{Rate}}{K[\text{H}_2]} = \frac{0.0040}{2.7 \times 10^{-3} \times 0.015} = 9.87 \times 10^{-5}$ $[\text{NO}] = 9.94 \times 10^{-3} \text{ mol l}^{-1}$																											
3a	acid partially dissociates	Strong acids are acids which fully dissociates into ions Weak acids are acids with partially dissociate into ions.																											
3b(i)	1.298	$\text{pH} = \frac{1}{2} \text{pK}_a - \frac{1}{2} \log_{10} c$ $\text{pH} = \frac{1}{2}(3.17) - \frac{1}{2} \log_{10}(3.75)$ $\text{pH} = 1.585 - \frac{1}{2}(0.574)$ $\text{pH} = 1.585 - 0.287$ $\text{pH} = 1.298$																											
3b(ii)	7.5%	gfm HF = (1x1)+(1x19) = 1+19 = 20g 3.75mol of HF in 1litre of solution ∴ mass = no. of mol x gfm = 3.75 x 20 = 75g 1 litre of HF solution would have approx. mass of 1000g $\frac{75\text{g}}{1000\text{g}} \times 100 = 7.5\%$																											
3b(iii)	tetrahedral	<table><tr><td>2 electron pairs 2 bonding + 0 lone pairs</td><td>3 electron pairs 3 bonding + 0 lone pairs</td><td>4 electron pairs 4 bonding + 0 lone pairs</td><td>4 electron pairs 3 bonding + 1 lone pairs</td><td>4 electron pairs 2 bonding + 2 lone pairs</td><td>5 electron pairs 5 bonding + 0 lone pairs</td><td>5 electron pairs 3 bonding + 2 lone pairs</td><td>6 electron pairs 6 bonding + 0 lone pairs</td><td>6 electron pairs 4 bonding + 2 lone pairs</td></tr><tr><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></tr><tr><td>Linear</td><td>Trigonal Planar</td><td>Tetrahedral</td><td>Trigonal Pyramidal</td><td>Angular</td><td>Trigonal Bipyramidal</td><td>T-Shaped</td><td>Octahedral</td><td>Square Planar</td></tr></table> $\begin{aligned} \text{Electron pairs} &= \frac{\text{No of outer electrons on central atom} + \text{no. of bonds} - \text{charge}}{2} \\ &= \frac{4 + 4 - 0}{2} \\ &= \frac{8}{2} \\ &= 4 \text{ pairs of electrons (4 bonding pairs)} \end{aligned}$	2 electron pairs 2 bonding + 0 lone pairs	3 electron pairs 3 bonding + 0 lone pairs	4 electron pairs 4 bonding + 0 lone pairs	4 electron pairs 3 bonding + 1 lone pairs	4 electron pairs 2 bonding + 2 lone pairs	5 electron pairs 5 bonding + 0 lone pairs	5 electron pairs 3 bonding + 2 lone pairs	6 electron pairs 6 bonding + 0 lone pairs	6 electron pairs 4 bonding + 2 lone pairs										Linear	Trigonal Planar	Tetrahedral	Trigonal Pyramidal	Angular	Trigonal Bipyramidal	T-Shaped	Octahedral	Square Planar
2 electron pairs 2 bonding + 0 lone pairs	3 electron pairs 3 bonding + 0 lone pairs	4 electron pairs 4 bonding + 0 lone pairs	4 electron pairs 3 bonding + 1 lone pairs	4 electron pairs 2 bonding + 2 lone pairs	5 electron pairs 5 bonding + 0 lone pairs	5 electron pairs 3 bonding + 2 lone pairs	6 electron pairs 6 bonding + 0 lone pairs	6 electron pairs 4 bonding + 2 lone pairs																					
Linear	Trigonal Planar	Tetrahedral	Trigonal Pyramidal	Angular	Trigonal Bipyramidal	T-Shaped	Octahedral	Square Planar																					
4a(i)A	Heating under reflux	Heating under reflux allows heating of a chemical reaction mixture over an extended period of time without volatile substances escaping <ul style="list-style-type: none">a round-bottomed flask is used containing anti-bumping granules to reduce bumping of reactants in the flaskflask is fitted with a condenser to allow condensation of reaction mixture vapour and prevents escape of volatile chemicalsflask is heated using an appropriate heat source e.g. heating mantle it there are flammable reactant/products involved.																											
4a(i)B	Anti-bumping granules																												
4a(ii)	Answer to include:	<div><p>few ions </p></div>																											

4a(iii)A	Buchner funnel with vacuum filtration	Vacuum filtration: filtration under reduced pressure <ul style="list-style-type: none">faster means of separating a precipitate from a filtrate Büchner, Hirsch or sintered glass funnel can be used.										
4a(iii)B	To purify the hippuric acid	Recrystallisation: purify an impure solid with: <ul style="list-style-type: none">dissolving an impure solid gently in a minimum volume of a hot solventhot filtration of resulting mixture to remove any insoluble impuritiescooling the filtrate slowly to allow crystals of the pure compound to form, leaving soluble impurities dissolved in the solventfiltering, washing and drying the pure crystals solvent for recrystallisation is chosen so that the compound being purified is completely soluble at high temperatures and only sparingly soluble at lower temperatures.										
4b(i)	Different atoms types in bonds and different strengths of bonds decides which wavelengths/wavenumbers of infrared are absorbed.	When infrared radiation is absorbed by organic compounds, bonds within the molecule vibrate/stretch/bend <ul style="list-style-type: none">wavelengths of infrared radiation absorbed depend on the type of atoms that make up the bond and the strength of the bondinfrared radiation is passed through a sample of the organic compound and then into a detector that measures the intensity of the transmitted radiation at different wavelengths. The absorbance of infrared radiation is measured in wavenumbers, measured in cm^{-1}										
4b(ii)	3500 - 3300	Information found in data booklet.										
4c(i)	4?	<table border="1"><thead><tr><th>^1H Environment</th><th>Grouping</th></tr></thead><tbody><tr><td>①</td><td>ArH (9.0-6.9)</td></tr><tr><td>②</td><td>Amide link (9.4-5.0)</td></tr><tr><td>③</td><td>$-\text{CH}_2-$ (?)</td></tr><tr><td>④</td><td>R-CH_2-COOH (2.7-2.0)</td></tr></tbody></table> 	^1H Environment	Grouping	①	ArH (9.0-6.9)	②	Amide link (9.4-5.0)	③	$-\text{CH}_2-$ (?)	④	R- CH_2 -COOH (2.7-2.0)
^1H Environment	Grouping											
①	ArH (9.0-6.9)											
②	Amide link (9.4-5.0)											
③	$-\text{CH}_2-$ (?)											
④	R- CH_2 -COOH (2.7-2.0)											
4c(ii)	doublet	<table><tr><td> singlet no adjacent hydrogens</td><td> Doublet 1 adjacent hydrogen</td><td> Triplet 2 adjacent hydrogens</td><td> quartet 3 adjacent hydrogens</td><td> quintet 4 adjacent hydrogens</td></tr></table>	 singlet no adjacent hydrogens	 Doublet 1 adjacent hydrogen	 Triplet 2 adjacent hydrogens	 quartet 3 adjacent hydrogens	 quintet 4 adjacent hydrogens					
 singlet no adjacent hydrogens	 Doublet 1 adjacent hydrogen	 Triplet 2 adjacent hydrogens	 quartet 3 adjacent hydrogens	 quintet 4 adjacent hydrogens								
4c(iii)		^1H nuclei behave like tiny magnets and in a strong magnetic field some align with the field (lower energy), whilst the rest align against it (higher energy). <ul style="list-style-type: none">Absorption of radiation in the radio frequency region of the electromagnetic spectrum causes the ^1H nuclei to 'flip' from the lower to the higher energy alignment.As they fall back from the higher to the lower energy alignment the emitted radiation is detected and plotted on a spectrum.In a ^1H NMR spectrum the chemical shift, δ, (peak position) is related to the environment of the ^1H atom and is measured in parts per million (ppm). The area under the peak is related to the number of ^1H atoms in that environment and is often given by an integration curve on a spectrum. The height of an integration curve is proportional to the number of ^1H atoms in that environment, and so a ratio of ^1H atoms in each environment can be determined.										
4d	Open Question to include:	<table><tr><th>3 mark answer</th><th>2 mark answer</th><th>1 mark answer</th></tr><tr><td>Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.</td><td>Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.</td><td>Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.</td></tr></table>	3 mark answer	2 mark answer	1 mark answer	Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.				
3 mark answer	2 mark answer	1 mark answer										
Demonstrates a good understanding of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a reasonable understanding of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a limited understanding of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.										

5a		<p>Ligands may be negative ions or molecules with non-bonding pairs of electrons that they donate to the central metal atom or ion, forming dative covalent bonds. Electrons in approaching ligands cause the splitting of d orbitals into higher and lower energies as the electrons along the axes to be repelled.</p> <ul style="list-style-type: none">ligands that cause a large difference in energy between subsets of d orbitals are strong field ligands.weak field ligands cause a small energy difference.spectrochemical series is a list of ligands placed in order of their ability to split d orbitals																																
5b(i)	Orbitals fill up in order of increasing energy	<p><u>Aufbau Principle:</u> Electrons fill up in order of increasing energy:</p> <p>1s 2s 2p 3s 3p 4s 3d 4p 5s 4d 5p 6s 4f 5d 6p 7s 5f 6d 7p 8s</p> 																																
5b(ii)A	Both conclusions required:	<table><tr><td>Conclusion 1</td><td>Complex Ion</td><td>$[\text{Co}(\text{OH}_2)_6]^{2+}$</td><td>$[\text{CoF}_6]^{3-}$</td><td>$[\text{Co}(\text{OH}_2)_6]^{3+}$</td><td>$[\text{Co}(\text{CN})_6]^{3-}$</td></tr><tr><td>Higher charge of Cobalt in complex gives higher energy required to pair two electrons with opposite spin (P)</td><td>Charge on Co</td><td>Co^{2+}</td><td>Co^{3+}</td><td>Co^{3+}</td><td>Co^{3+}</td></tr><tr><td>Conclusion 2: Higher spin state has lower difference in energy between subsets of d orbitals (Δ)</td><td>P (kJ mol⁻¹)</td><td>250</td><td>283</td><td>283</td><td>283</td></tr></table> <table><tr><td>Δ (kJ mol⁻¹)</td><td>111</td><td>156</td><td>323</td><td>416</td></tr><tr><td>Spin State</td><td>High</td><td>High</td><td>Low</td><td>Low</td></tr></table>	Conclusion 1	Complex Ion	$[\text{Co}(\text{OH}_2)_6]^{2+}$	$[\text{CoF}_6]^{3-}$	$[\text{Co}(\text{OH}_2)_6]^{3+}$	$[\text{Co}(\text{CN})_6]^{3-}$	Higher charge of Cobalt in complex gives higher energy required to pair two electrons with opposite spin (P)	Charge on Co	Co^{2+}	Co^{3+}	Co^{3+}	Co^{3+}	Conclusion 2: Higher spin state has lower difference in energy between subsets of d orbitals (Δ)	P (kJ mol ⁻¹)	250	283	283	283	Δ (kJ mol ⁻¹)	111	156	323	416	Spin State	High	High	Low	Low				
Conclusion 1	Complex Ion	$[\text{Co}(\text{OH}_2)_6]^{2+}$	$[\text{CoF}_6]^{3-}$	$[\text{Co}(\text{OH}_2)_6]^{3+}$	$[\text{Co}(\text{CN})_6]^{3-}$																													
Higher charge of Cobalt in complex gives higher energy required to pair two electrons with opposite spin (P)	Charge on Co	Co^{2+}	Co^{3+}	Co^{3+}	Co^{3+}																													
Conclusion 2: Higher spin state has lower difference in energy between subsets of d orbitals (Δ)	P (kJ mol ⁻¹)	250	283	283	283																													
Δ (kJ mol ⁻¹)	111	156	323	416																														
Spin State	High	High	Low	Low																														
5b(ii)B	Hexafluoridocobaltate (III)	<p>Hexafluoridocobaltate(III) = $[\text{CoF}_6]^{3-}$</p> <table><tr><td>no. of ligands</td><td>F⁻ ligand</td><td>metal name</td><td>negative charge on complex</td><td>metal ion</td></tr></table> <table><tr><td colspan="2">Neutral ligands include:</td><td colspan="2">Negative Ligands include:</td><td>Central Ion:</td><td>Charge:</td></tr><tr><td>Ligand</td><td>Name</td><td>Ligand</td><td>Name</td><td>Positive Complex: metals keep their name</td><td rowspan="4">Charge of central ion is converted into roman numerals and put in brackets</td></tr><tr><td>H₂O</td><td>aqua</td><td>Chloride Cl⁻</td><td>chlorido</td><td>Negative Complex: Metals end in ATE</td></tr><tr><td>NH₃</td><td>ammine</td><td>Cyanide CN⁻</td><td>cyanido</td><td>Metals end in ATE</td></tr><tr><td>CO</td><td>carbonyl</td><td>Nitrite NO₂⁻</td><td>nitrito</td><td>e.g. Cuprate, Ferrate, Cobaltate</td></tr></table>	no. of ligands	F ⁻ ligand	metal name	negative charge on complex	metal ion	Neutral ligands include:		Negative Ligands include:		Central Ion:	Charge:	Ligand	Name	Ligand	Name	Positive Complex: metals keep their name	Charge of central ion is converted into roman numerals and put in brackets	H ₂ O	aqua	Chloride Cl ⁻	chlorido	Negative Complex: Metals end in ATE	NH ₃	ammine	Cyanide CN ⁻	cyanido	Metals end in ATE	CO	carbonyl	Nitrite NO ₂ ⁻	nitrito	e.g. Cuprate, Ferrate, Cobaltate
no. of ligands	F ⁻ ligand	metal name	negative charge on complex	metal ion																														
Neutral ligands include:		Negative Ligands include:		Central Ion:	Charge:																													
Ligand	Name	Ligand	Name	Positive Complex: metals keep their name	Charge of central ion is converted into roman numerals and put in brackets																													
H ₂ O	aqua	Chloride Cl ⁻	chlorido	Negative Complex: Metals end in ATE																														
NH ₃	ammine	Cyanide CN ⁻	cyanido	Metals end in ATE																														
CO	carbonyl	Nitrite NO ₂ ⁻	nitrito	e.g. Cuprate, Ferrate, Cobaltate																														
5b(ii)C		<table><tr><td>Electronic Configuration</td><td>Low Spin state of Co^{3+}</td><td>High Spin State of Co^{3+}</td></tr><tr><td>Element Cobalt Configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁷ 4s²</td><td></td><td></td></tr><tr><td>Cobalt Co^{3+} Ion Configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁶</td><td></td><td></td></tr></table>	Electronic Configuration	Low Spin state of Co^{3+}	High Spin State of Co^{3+}	Element Cobalt Configuration: 1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 3d ⁷ 4s ²			Cobalt Co^{3+} Ion Configuration: 1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 3d ⁶																									
Electronic Configuration	Low Spin state of Co^{3+}	High Spin State of Co^{3+}																																
Element Cobalt Configuration: 1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 3d ⁷ 4s ²																																		
Cobalt Co^{3+} Ion Configuration: 1s ² 2s ² 2p ⁶ 3s ² 3p ⁶ 3d ⁶																																		
6a(i)	Circled region of alternating C=C double bonds and C-C single bond.																																	
6a(ii)	Antibonding molecular orbitals are higher in energy	Electrons fill bonding molecular orbitals, leaving higher energy antibonding molecular orbitals unfilled.																																
6b(i)	Answer to include:	<p>C=C double bonds do not rotate around their axis so geometric isomers exist depending on the position of side groups in relation to C=C double bond.</p> <table><tr><td>-CH₃ methyl groups are on opposite sides of the C=C double bond</td><td>-CH₃ methyl groups are on same side of the C=C double bond</td></tr><tr><td><p>trans-but-2-ene</p></td><td><p>cis-but-2-ene</p></td></tr></table>	-CH ₃ methyl groups are on opposite sides of the C=C double bond	-CH ₃ methyl groups are on same side of the C=C double bond	 <p>trans-but-2-ene</p>	 <p>cis-but-2-ene</p>																												
-CH ₃ methyl groups are on opposite sides of the C=C double bond	-CH ₃ methyl groups are on same side of the C=C double bond																																	
 <p>trans-but-2-ene</p>	 <p>cis-but-2-ene</p>																																	

6b(ii)	Dotted lines on diagram showing:	Dotted lines must be between: H of an N-H or O-H bond and O of a O-H bond or N of N-H bond			
6c(i)	Blue light provides enough energy to break bonds in bilirubin	Blue light has shorter wavelength than red light and has more energy.			
6c(ii)	257.5	$E = \frac{L \times h \times c}{\lambda} = \frac{6.02 \times 10^{23} \text{ mol}^{-1} \times 6.63 \times 10^{-34} \text{ J s} \times 3 \times 10^8 \text{ m s}^{-1}}{465 \times 10^{-9} \text{ m}}$ $= 257501 \text{ J mol}^{-1}$ $= 257.5 \text{ kJ mol}^{-1}$			
7a(i)	Answer to include:	Place empty weighing bottle or weighing boat on balance. Record the mass of the empty item on balance. Add substance to boat/bottle on balance (careful not to spill any onto the top of balance). Record mass and subtract the empty mass to get mass of substance.			
7a(ii)	0.0204	Absorbance = 0.42 $\therefore [\text{MnO}_4^-] = 2.1 \times 10^{-4} \text{ mol l}^{-1}$ no. of mol = volume \times concentration = $0.1 \text{ litres} \times 2.1 \times 10^{-4} \text{ mol l}^{-1} = 0.000021 \text{ mol}$ gfm = 54.9g mass = no. of mol \times gfm = $0.000021 \text{ mol} \times 54.9 \text{ g mol}^{-1} = 0.00115 \text{ g}$ $\% \text{ mass} = \frac{0.00115}{5.66} \times 100 = 0.0204\%$			
7b(i)	one from:	high state of purity	be stable when solid and in solution	be soluble	reasonably high GFM
7b(ii)	0.358	no. of mol $\text{Cr}_2\text{O}_7^{2-} = \text{volume} \times \text{concentration} = 0.0214 \text{ litres} \times 0.005 \text{ mol l}^{-1} = 0.000107 \text{ mol}$ $6\text{Fe}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \longrightarrow 6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$ 6mol 1mol 0.000642mol 0.000107mol 25cm ³ Fe ²⁺ solution = 0.000642mol 250cm ³ Fe ²⁺ solution = 0.00642mol gfm Fe = 55.8g mass = no. of mol \times gfm = $0.00642 \text{ mol} \times 55.8 \text{ g mol}^{-1} = 0.358 \text{ g}$			
7b(iii)	Answer to contain:	1 mark Green wavelengths absorbed turning HOMO into LUMO		1mark Green wavelengths absorbed \therefore red + blue wavelengths transmitted and purple light emitted	
8a	H ₃ PO ₃ (any order of elements)	$3\text{CH}_3\text{COOH} + \text{PCl}_3 \longrightarrow 3\text{CH}_3\text{COCl} + \text{H}_3\text{PO}_3$			
8b(i)	Species attracted to centre of positive charge	Nucleophiles are negatively charged ions or neutral molecules which are electron rich <ul style="list-style-type: none">attracted towards atoms bearing a partial δ^+ or full positive chargecapable of donating an electron pair to form a new covalent bond Examples: Cl ⁻ , Br ⁻ , OH ⁻ , CN ⁻ , NH ₃ and H ₂ O Electrophiles are positively charged ions or neutral molecules that are electron deficient <ul style="list-style-type: none">attracted towards atoms bearing a partial δ^- or full negative chargecapable of accepting an electron pair to form a new covalent bond Examples: H ⁺ , NO ₂ ⁺ and SO ₃			
8b(ii)	Curly arrow from O on alcohol going to C of acid chloride group				
8b(iii)	Diagram showing:				

8b(iv)	HCl	Condensation reactions join two molecules together and remove a small molecule as they join. Water is the most likely small molecule removed but hydrogen chloride HCl is removed in the condensation of alcohols and acid chloride to form esters.		
8b(v)	Great yield of ester	As the equilibrium lies more to the product side in this reaction then more ester is produced at equilibrium		
8c(i)	Secondary	Primary Amine	Secondary Amine	Tertiary Amine
		$\begin{array}{c} \text{H}-\text{N}-\text{C}_3\text{H}_7 \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{H}_3\text{C}-\text{N}-\text{C}_2\text{H}_5 \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{H}_3\text{C}-\text{N}-\text{CH}_3 \\ \\ \text{CH}_3 \end{array}$
		1 Carbon attached to the Nitrogen	2 Carbons attached to the Nitrogen	3 Carbons attached to the Nitrogen
8c(ii)	amide	Amide links have a carbonyl group attached to a nitrogen which can have one or two carbon groups attached to the nitrogen.	$\begin{array}{cc} \text{O} & \text{H} \\ & \\ -\text{C} & -\text{N}- \end{array}$	$\begin{array}{cc} \text{O} & \text{CH}_3 \\ & \\ -\text{C} & -\text{N}- \end{array}$
8d(i)	Condensation	This reaction removes the Cl from benzoyl chloride and one of the H atoms from the benzene and join them together to form Benzophenone. Hydrogen chloride is removed as they join together.		
8d(ii)	65.2%	Benzoyl chloride $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{21.8}{140.5} = 0.155\text{mol}$		Benzophenone $\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{18.4}{182} = 0.101\text{mol}$ <small>(actual)</small>
		$\begin{array}{ccc} \text{benzoyl chloride} + \text{benzene} & \longrightarrow & \text{benzophenone} + \text{hydrogen chloride} \\ 1\text{mol} & & 1\text{mol} \\ 0.155\text{mol} & & 0.155\text{mol} \\ & & \text{(theoretical)} \end{array}$		
		$\% \text{ Yield} = \frac{\text{Actual}}{\text{Theoretical}} \times 100 = \frac{0.101\text{mol}}{0.155\text{mol}} \times 100 = 65.2\%$		
9	Open Question to include:	3 mark answer	2 mark answer	1 mark answer
		Demonstrates a <u>good understanding</u> of the chemistry involved. A good comprehension of the chemistry has provided in a logically correct, including a statement of the principles involved and the application of these to respond to the problem.	Demonstrates a <u>reasonable understanding</u> of the chemistry involved, making some statement(s) which are relevant to the situation, showing that the problem is understood.	Demonstrates a <u>limited understanding</u> of the chemistry involved. The candidate has made some statement(s) which are relevant to the situation, showing that at least a little of the chemistry within the problem is understood.
10a	Mimics the natural compound that binds to the receptor and produced the biological response in the cell	agonist	An agonist mimics the natural compound and binds to the receptor molecules to produce a response similar to the natural active compound.	
		antagonist	An antagonist prevents the natural compound from binding to the receptor, and so blocks the natural response from occurring.	
10b(i)	5cm ³	A one in one hundred dilution requires a 5cm ³ of eucalyptol transferred by pipette into a 500cm ³ standard/volumetric flask and the filled up to the mark with deionised water.		
10b(ii)	0.0598	mass = density x Volume = 0.921g cm ⁻³ x 5cm ³ = 4.605g		
		$\text{no. of mol} = \frac{\text{mass}}{\text{gfm}} = \frac{4.605 \text{ g}}{154 \text{ g mol}^{-1}} = 0.0299\text{mol}$ $\text{concentration} = \frac{\text{no. of mol}}{\text{Volume}} = \frac{0.0299 \text{ mol}}{0.500 \text{ litres}} = 0.0598\text{mol l}^{-1}$		
10b(iii)	6	$\text{no. of mol} = \text{Volume} \times \text{concentration} = 1\text{litre} \times 9.97 \times 10^{-24} \text{mol l}^{-1} = 9.97 \times 10^{-24} \text{mol}$ $1 \text{ mol} = 6.02 \times 10^{23} \text{ molecules}$ $9.97 \times 10^{-24} \text{mol} = 6.02 \times 10^{23} \text{ molecules} \times \frac{9.97 \times 10^{-24}}{1}$ $= 6.00 \text{ molecules}$		

11a(i)	$C_8H_{10}O_2N_4$		
		Skeletal Structure	Full Structural Formula
11a(ii)	Molecules must have similar shape to bind to same receptor protein	<p>Pharmacologically active molecules must have a similar shape to fit the same receptor molecule. This common shape is called the pharmacophore.</p> <p>Agonists fit the receptor and produce the same biological response as the natural substrate. Antagonist fit the receptor but do not cause the biological response inside the cell.</p>	
11b(i)A		<p>The steps of recrystallisation to purify an impure solid include:</p> <ul style="list-style-type: none"> dissolving an impure solid gently in a minimum volume of a hot solvent hot filtration of the resulting mixture to remove any insoluble impurities cooling the filtrate slowly to allow crystals of the pure compound to form, leaving soluble impurities dissolved in the solvent filtering, washing and drying the pure crystals <p>The solvent used for recrystallisation is chosen so that the compound being purified is completely soluble at high temperatures and only sparingly soluble at lower temperatures. The solvent used should be:</p> <ul style="list-style-type: none"> immiscible with the liquid mixture or solution (usually water) one in which the solute is more soluble in than the liquid mixture or solution (usually water) volatile to allow the solute to be obtained by evaporation of the solvent unreactive with the solute 	
11b(i)B	4.61	$K = \frac{[\text{caffeine}]_{\text{dichloromethane}}}{[\text{caffeine}]_{\text{water}}} = \frac{23.5/60}{8.5/100} = \frac{0.392}{0.085} = 4.61$	
11b(ii)	Do extraction in three separate volumes of 20cm ³	The quantity of caffeine extracted is greater if a number of extractions using smaller volumes of solvent are carried out rather than a single extraction using a large volume of solvent.	
11c	Diagram completed as shown:	