

# Chapter 1

## ThermoS26-06

### 1.1 Legendre Transforms and Thermodynamic Potentials

**Definition 1.1.1.** (5.1) Suppose that  $f : [a, b] \rightarrow \mathbb{R}$  is continuous. For every  $p \in [c, d]$ , define

$$f^*(p) := \sup_{x \in [a, b]} \{xp - f(x)\} \quad (5.1)$$

The function  $f^* : [c, d] \rightarrow \mathbb{R}$  is called the Legendre transform of  $f$ .

**Theorem 1.1.2.** (5.2) Suppose that  $f \in C^2([0, \infty); \mathbb{R})$  with

$$f''(x) > 0, \quad \forall x \in [0, \infty).$$

Then, for all  $p \in \text{Range}(f')$ ,

$$f^*(p) = x_p p - f(x_p), \quad (5.2)$$

where  $x_p \in [0, \infty)$  is the unique solution to

$$f'(x_p) = p.$$

*Proof.* let  $p \in \text{Range}(f')$ .

$$f' : [0, \infty) \rightarrow \text{Range}(f')$$

For every  $p \in \text{Range}(f') \exists! x_p \in [0, \infty)$  and that,

$$f'(x_p) = p.$$

Since  $f' : [0, \infty)$  is strictly increasing,  
 Fix  $p \in \text{Range}(f')$ . By Taylor's Theorem,

$$f(x) = f(x_p) + p(x - x_p) + \frac{1}{2}f''(\xi_p)(x - x_p)^2$$

for some  $\xi_p$  between  $x$  and  $x_p$ .

$$\begin{aligned} \sup_{0 \leq x < \infty} \{x \cdot p - f(x)\} &= \sup_{0 \leq x < \infty} \{x_p \cdot p - f(x_p) - \frac{1}{2}f''(\xi_p)(x - x_p)^2\} \\ &= x_p \cdot p - f(x_p). \end{aligned}$$

□

**Theorem 1.1.3.** (5.3) Suppose that  $f \in C^2([0, \infty); \mathbb{R})$  with

$$f''(x) > 0, \quad \forall x \in [0, \infty).$$

Then,

$$[f^*]'(p) = [f']^{-1}(p), \quad [f^*]''(p) > 0,$$

for all  $p \in \text{Range}(f')$ .

*Proof.* Recall

$$f' : [0, \infty) \rightarrow \text{Range}(f')$$

Then

$$f'([f']^{-1}(p)) = p, \quad \forall p \in \text{Range}(f').$$

Also, observe that  $\frac{d}{dp}[f']^{-1} \in C(\text{Range}(f'); \mathbb{R})$ .

Now,

$$\begin{aligned} \frac{d}{dp}[f^*(p)] &= \frac{d}{dp}\{x_p \cdot p - f(x_p)\} = \frac{d}{dp}\{[f']^{-1}(p) \cdot p - f([f']^{-1}(p))\} \\ &= p \cdot \frac{d}{dp}[f']^{-1}(p) + [f']^{-1}(p) - \frac{df}{dx}([f']^{-1}(p)) \cdot \frac{d}{dp}[f']^{-1}(p) \\ &= [f']^{-1}(p). \end{aligned}$$

Now that this is established, we have

$$\frac{d^2}{dp^2}[f^*(p)] = \frac{d}{dp}[f']^{-1}(p)$$

for all  $p \in \text{Range}(f')$ . Since  $f'' \in C([0, \infty))$  and  $f''(x) > 0, \forall x \in [0, \infty)$ ,

$$\frac{d}{dp}[f']^{-1}(p) > 0, \quad \forall p \in \text{Range}(f').$$

□

Finally, we observe the following:

**Theorem 1.1.4.** (5.4) *Suppose that  $f \in C^2([0, \infty); \mathbb{R})$  with*

$$f''(x) > 0, \quad \forall x \in [0, \infty).$$

*Then  $f^* \in C^2(\text{Range}(f'); \mathbb{R})$  with*

$$[f^*]''(p) > 0, \quad \forall p \in \text{Range}(f')$$

*Furthermore*

$$[f^*]^*(x) = f(x), \quad \forall x \in [0, \infty). \quad (5.3)$$

*In other words, the Legendre transform is involutive and in fact an isomorphism.*

*Proof.* It follows that, for all  $x \in [0, \infty)$ ,

$$[f^*]^*(x) = x p_x - f^*(p_x),$$

where  $p_x \in \text{Range}(f')$  is the unique solution to

$$f^{*'}(p_x) = x.$$

Recall

$$f^*(p_x) = p_x \cdot x_{p_x} - f(x_{p_x}),$$

where  $x_{p_x} \in [0, \infty)$  is the unique solution to

$$f'(x_{p_x}) = p_x.$$

Of course, by uniqueness,

$$x_{p_x} = x.$$

So,

$$[f^*]^*(x) = x \cdot p_x - \{p_x \cdot x_{p_x} - f(x_{p_x})\} = f(x).///$$

□

Why is the Legendre transform useful in Thermodynamics?

This transform allows us to introduce new thermodynamics coordinates/-variables.

Recall

$$\tilde{U} = \tilde{U}(S, V, N)$$

Extensive Quantity	Conjugate Variable	Variable
$\tilde{U}$	$T, -p, \mu$	$S, V, N$
$\tilde{S}$	$\frac{1}{T}, \frac{p}{T}, -\frac{\mu}{T}$	$U, V, N$

Table 1.1: Conjugate variables for  $\tilde{U}$  and  $\tilde{S}$ .

or

$$d\tilde{U} = T dS - p dV + \mu dN$$

The latter means

$$T = \frac{\partial \tilde{U}}{\partial S} \quad p = -\frac{\partial \tilde{U}}{\partial V} \quad \mu = \frac{\partial \tilde{U}}{\partial N}$$

in short hand

**Definition 1.1.5.** (5.5) Suppose that all of the Thermodynamic Postulates hold. Assume that the material in question is unary ( $r = 1$ ). Define

$$\tilde{F}(T, V, N) = \tilde{U}(S_T, V, N) - TS_T \quad (5.4)$$

where  $S_T = S_T(V, N)$  is the unique solution of the equation

$$T_U(S_T(V, N), V, N) = T \quad (5.5)$$

for fixed values of  $N$  and  $V$ .

$\tilde{F}$  is called the Helmholtz free energy.

Formally, we have

$$\begin{aligned} d\tilde{F} &= d\tilde{U} - S dT - T dS \\ &= T dS - p dV + \mu dN - S dT - T dS \\ &= -S dT - p dV + \mu dN. \end{aligned}$$

Extensive Quantity	Conjugate Variable	Variable
$\tilde{U}$	$T, -p, \mu$	$S, V, N$
$\tilde{F}$	$-S, -p, \mu$	$T, V, N$

Table 1.2: Conjugate variables for  $\tilde{U}$  and  $\tilde{F}$ .

**Example 1.1.6.** (5.6) Suppose that

$$\tilde{S}(U, V, N) = \left( \frac{NVUR^2}{v_0\theta} \right)^{1/3}.$$

It follows that

$$\tilde{U}(S, V, N) = \frac{S^3 v_0 \theta}{NVR^2}.$$

The temperature function is

$$T_U(S, V, N) = \frac{3S^2 v_0 \theta}{NVR^2}.$$

Or

$$\frac{1}{T_S(U, V, N)} = \frac{1}{3} \left( \frac{NVUR^2}{v_0\theta} \right)^{-2/3} \frac{NVR^2}{v_0\theta} = \frac{1}{3} \left( \frac{NVR^2}{v_0\theta} \right)^{1/3} U^{-2/3}.$$

Thus

$$T_S(U, V, N) = 3 \left( \frac{v_0\theta}{NVR^2} \right)^{1/3} U^{2/3}.$$

Recall that

$$T_U(S, V, \vec{N}) = T_S(\tilde{U}(S, V, \vec{N}), V, \vec{N})$$

$$T_S(U, V, \vec{N}) = T_U(\tilde{S}(U, V, \vec{N}), V, \vec{N}).$$

By definition

$$\tilde{F}(T, V, N) = \tilde{U}(S_T, V, N) - TS_T$$

where

$$T_U(S_T(V, N), V, N) = T.$$

For our example,

$$T_U(S, V, N) = \frac{3S^2 v_0 \theta}{NVR^2}.$$

Hence,

$$S_T = \sqrt{\frac{TNVR^2}{3v_0\theta}}.$$

Thus,

$$\begin{aligned}\tilde{F}(T, V, N) &= \tilde{U}\left(\sqrt{\frac{TNVR^2}{3v_0\theta}}, V, N\right) - T\sqrt{\frac{TNVR^2}{3v_0\theta}} \\ &= \left(\frac{TNVR^2}{3v_0\theta}\right)^{3/2} \frac{v_0\theta}{NVR^2} - T\sqrt{\frac{TNVR^2}{3v_0\theta}}.\end{aligned}$$

But

$$\tilde{U}(S_T, V, N) = \frac{S_T^3 v_0\theta}{NVR^2} = \left(\frac{TNVR^2}{3v_0\theta}\right)^{3/2} \frac{v_0\theta}{NVR^2}$$

so that

$$\tilde{F}(T, N, V) = \frac{1}{3}T^{3/2} \frac{1}{\sqrt{3}} \sqrt{\frac{NVR^2}{v_0\theta}} - \frac{T^{3/2}}{\sqrt{3}} \sqrt{\frac{NVR^2}{v_0\theta}}.$$

Finally,

$$\tilde{F}(T, N, V) = -\frac{2T^{3/2}}{3} \sqrt{\frac{NVR^2}{3v_0\theta}}. \quad (5.6)$$

Now,

$$\frac{\partial \tilde{F}}{\partial T} = -\frac{2}{3} \frac{3}{2} \sqrt{T} \sqrt{\frac{NVR^2}{3v_0\theta}}.$$

But

$$T_U(S, V, N) = \frac{3S^2 v_0\theta}{NVR^2}$$

Thus,

$$\sqrt{T} = \sqrt{\frac{3v_0\theta}{NVR^2}} S_T$$

and

$$\frac{\partial \tilde{F}}{\partial T} = -S_T. \quad (5.7)$$

**Theorem 1.1.7.** (5.7) Suppose that all of the Thermodynamic Postulates hold. Assume that the material in question is unary ( $r = 1$ ). The following procedure is equivalent means for finding the Helmholtz free energy:

$$\tilde{F}(T, V, N) = U_T - T\tilde{S}(U_T, V, N) \quad (5.8)$$

where  $U_T = U_T(V, N)$  is the unique solution of the equation

$$T_S(U_T(V, N), V, N) = T$$

for fixed values of  $N$  and  $V$ .

Proof: Exercise ///

**Example 1.1.8.** (5.8) Let us use the second version to calculate the free energy for the monatomic ideal gas. Recall,

$$\tilde{S}(U, V, N) = Ns_0 + NR \ln \left( \left( \frac{U}{U_0} \right)^{3/2} \left( \frac{V}{V_0} \right) \left( \frac{N}{N_0} \right)^{-5/2} \right),$$

where  $s_0, R, U_0, V_0, N_0$  are positive constants.

Then

$$\begin{aligned} \frac{\partial \tilde{S}}{\partial U} &= \frac{NR}{\left( \frac{U}{U_0} \right)^{3/2}} \cdot \frac{3}{2} \left( \frac{U}{U_0} \right)^{1/2} \frac{1}{U_0} \\ &= \frac{3NR}{2U_0} \cdot \frac{U_0}{U} = \frac{3NR}{2U} \end{aligned}$$

Thus

$$T_S(U, V, N) = \frac{2}{3} \frac{U}{NR}$$

and

$$U_T = \frac{3NRT}{2}$$

Therefore,

$$\begin{aligned} \tilde{F}(T, V, N) &= \frac{3NRT}{2} - TNs_0 \\ &\quad - NRT \ln \left( \left( \frac{3NRT}{2U_0} \right)^{3/2} \left( \frac{V}{V_0} \right) \left( \frac{N}{N_0} \right)^{-5/2} \right) \end{aligned}$$

Define

$$T_0 := \frac{2U_0}{3N_0R}.$$

Then,

$$\tilde{F}(T, V, N) = \frac{3NRT}{2} - TNs_0 - NRT \ln \left( \left( \frac{T}{T_0} \right)^{3/2} \left( \frac{V}{V_0} \right) \left( \frac{N}{N_0} \right)^{-1} \right) \quad (5.9)$$