



puget_oceantype Elwha River orge Adams Fall Fingerl Green River Nisqually Fall Fingerling 0.6 1.0 0.4 0.5 0.3 0.5 0.2 0.0 0.0 0.0 0.0 -0.3-0.2 -0.5-0.5-0.6 -0.4ooksack Spring Fingerlin kagit Summer Fingerlin Samish Fall Fingerling Skagit Spring Fingerling 0.4 0.6 1.0 0.3 0.2 0.5 0.5 0.0 0.0 0.0 0.0 -0.3-0.2 -0.5-0.5-0.6Skykomish Fall Fingerling 1 Puget Sound Fall Finge illaguamish Fall Fingerli sity of Washington Acce 1.0 0.6 0.6 1.0 0.3 0.5 0.4 0.5 0.0 0.2 0.0



puget_streamtype Nooksack Spring Yearling Skagit Spring Yearling South Puget Sound Fall Yearling 1.0 1 0.25 0.5 0 0.00 0.0 -1 -0.25-0.5 1970 1990 2010 Squaxin Pens Fall Yearling White River Spring Yearling 2 0.5 1 0 0.0 -1 -0.5-2 1970 1990 2010 1970 1990 2010 Time



south_oceantype Columbia River Summe umbia River Upriver Br Cowlitz Fall Tule Elk River 0.6 0.50 0.25 0.3 0.25 0.3 0.00 0.00 0.0 0.0 -0.25 -0.25-0.3 -0.3 -0.50-0.50-0.6-0.6Hoko Falls Lewis River Wild Hanford Wild Brights wer River Hatchery Tu 0.4 0.4 0.50 0.2 0.5 0.25 0.0 0.0 0.0 0.00 -0.2-0.5 -0.4-0.25-0.4 -0.50agan-Similkameen Su Lyons Ferry Fingerling Queets Fall Fingerling Robertson Creek Fall 0.8 0.4 0.4 0.2 1 0.2 0.0 0.0 0.0 0 -0.4-0.2-0.2 -1-0.8 -0.42010 1970 1990 Sooes Fall Fingerling Spring Creek Tule Salmon River 0.50 0.6 0.50 0.25 0.3 0.25 0.00 0.00 0.0 -0.25 -0.25 -0.3-0.50-0.501970 1990 2010 1970 1990 2010 1970 1990 2010 Time