


Week 12 Tuesday  
L-21



$$\frac{dM_r(r)}{dr} = 4\pi r^2 \rho(r)$$

$$\frac{dP(r)}{dr} = -\rho(r) \frac{GM_r(r)}{r^2}$$

$$\frac{dT(r)}{dr} = -\frac{3\kappa_R(r)\rho(r)}{64\pi\sigma} \frac{L_r(r)}{r^2 T(r)^3}$$

$$\frac{dL_r(r)}{dr} = 4\pi r^2 \rho(r) \epsilon(r)$$

$$M_r(r) \quad P(r) \quad L_r(r) \quad T(r)$$

$$\rho(r) \quad \mu(r) \quad \epsilon_{\text{nuc}}(r) \quad \kappa_R(r)$$

$$P(r) = \frac{\rho(r)}{\mu(r)} kT(r)$$

$$\mu(r) = f(\text{comp}, T(r), P(r))$$

$$\kappa_R(r) = f(\text{comp}, T(r), P(r))$$

$$\epsilon_{\text{nuc}}(r) = f(\text{comp}, T(r), P(r))$$

Other energy transport?

**Ideal gas always valid?**

Nuclear mechanism?

How to solve?

What changes with time?

# Summary of derivation for degeneracy pressure:

Momentum is quantized:

$$\vec{p}_{x,y,z} = \frac{h}{2L} n_{x,y,z}$$

Relation between the last filled level ( $n_{\text{Fermi}}$ ) and the total number of electrons:

$$n_F = \left( \frac{3N_{\text{tot}}}{\pi} \right)^{1/3}$$

Fermi momentum:

$$p(n_F) = \frac{h n_F}{2L} = \frac{h}{2} \left( \frac{3n_e}{\pi} \right)^{1/3}$$

Using the relation between concentration, density, and mean molecular weight:

$$= \frac{h}{2} \left( \frac{3}{\pi m_H} \right)^{1/3} \left( \frac{\rho}{\mu_e} \right)^{1/3}$$

Pressure is a momentum flux through a surface:

$$P_x = \int_0^\infty p_x v_x \underbrace{n(p)}_{\substack{\text{\# of electron} \\ dV}} 4\pi p^2 dp$$

In 3D, and spherical coordinates:

$$P_e = \frac{8\pi}{3h^3} \int_0^{p_F} v p^3 dp$$

Non-relativistic case:

$$v(p) = \frac{p}{m_e}$$

$$P_{\text{deg}}^{\text{non-relativistic}} = 1.00 \times 10^{13} \left( \frac{\rho}{\mu_e} \right)^{5/3} [\text{dyne/cm}^2]$$



Relativistic case:

$$v \approx c$$

$$P_{\text{deg}}^{\text{relativistic}} = 1.24 \times 10^{15} \left( \frac{\rho}{\mu_e} \right)^{4/3} \text{dyn/cm}^2$$



Remember this graph from the convection notebook:



# At which $\rho$ and $T$ does the degeneracy pressure becomes important?


1. Transition from non-relativistic to relativistic degeneracy pressure

$$\left(\frac{\rho}{\mu_e}\right)_{\text{transition}} \approx 1.9 \times 11$$

2. Transition from ideal gas pressure to degeneracy pressure

$$\left(\frac{\rho}{\mu_e}\right)_{\text{trans non-rel}} = 2.5 \times 10^{-8} T^{3/2} \left[ \frac{\text{g}}{\text{cm}^3} \right]$$

$$\left(\frac{\rho}{\mu_e}\right)_{\text{trans rel}} = 2.95 \times 10^{-22} T^3 \left[ \frac{\text{g}}{\text{cm}^3} \right]$$



$$P_{\text{deg}}^{\text{non-relativistic}} = 1.00 \times 10^{13} \left( \frac{\rho}{\mu_e} \right)^{5/3} [\text{dyne/cm}^2]$$

$$P_{\text{deg}}^{\text{relativistic}} = 1.24 \times 10^{15} \left( \frac{\rho}{\mu_e} \right)^{4/3} \text{dyn/cm}^2$$



$$P_{\text{deg}}^{\text{non-relativistic}} = 1.00 \times 10^{13} \left( \frac{\rho}{\mu_e} \right)^{5/3} [\text{dyne/cm}^2]$$

$$P_{\text{deg}}^{\text{relativistic}} = 1.24 \times 10^{15} \left( \frac{\rho}{\mu_e} \right)^{4/3} \text{dyn/cm}^2$$

## Polytrops


$$P = k \rho^{1 + 1/n}$$

$$P_{\text{deg}}^{\text{non-relativistic}} = 1.00 \times 10^{13} \left( \frac{\rho}{\mu_e} \right)^{5/3} [\text{dyne/cm}^2]$$

$$P_{\text{deg}}^{\text{relativistic}} = 1.24 \times 10^{15} \left( \frac{\rho}{\mu_e} \right)^{4/3} \text{dyn/cm}^2$$

## Polytrops

$$P = k \rho^{1 + 1/n}$$



$n$

if we pick a

$P_c$   $\rho_c$   $R_\star$

$K$  known

we can find the  
 $M_\star$

$$1 \quad R_\star = \alpha \epsilon_1 = \left( \frac{(n+1)P_c}{4\pi G \rho_c^2} \right)^{1/2} \epsilon_1$$

$$2 \quad P_c = K \rho_c^{(n+1)/n}$$

$$3 \quad M_\star = -\frac{1}{(4\pi)^{1/2}} \left( \frac{n+1}{G} \right)^{3/2} \frac{P_c^{3/2}}{\rho_c^2} \epsilon_1^2 \theta'(\epsilon_1)$$

## 2. At home: Use the properties of polytrope to understand non-relativistic white dwarfs

White dwarfs are the remnant of solar mass object, once the fusion stops. WD are generally made of elements heavier than He, e.g. carbon or oxygen depending on their evolution. In the case where no H or He is present in a gas, the electronic mean molecular weight  $\mu_e \sim 2$ .

These objects remain in hydrostatic equilibrium because of non-relativistic degeneracy pressure, of the form

$$P_e = 1.00 \times 10^{13} \left( \frac{\rho}{\mu_e} \right)^{5/3}.$$

The numerical constant is such that the pressure is in dyn/cm<sup>2</sup> for a density given in g/cm<sup>3</sup>.

You probably notice that the pressure does not depend on temperature, just like a polytrop!

This would correspond to a polytrop of index  $n=3/2$ , with  $K = 1.00 \times 10^{13} / \mu_e^{5/3}$ . For such a polytrop,  $\epsilon_1 = 3.654$ , and  $(-\theta(\epsilon_1)') = 0.20330$ , according to Henson Table 7.1.

## 3. At home: Do the same as #2, for a relativistic white dwarf.