

G-ESTIMATION OF STRUCTURAL NESTED MODELS

Joy Shi, Barbra Dickerman, Miguel Hernán
DEPARTMENTS OF EPIDEMIOLOGY AND BIostatISTICS



HARVARD T.H. CHAN
SCHOOL OF PUBLIC HEALTH

Learning objectives

At the end of this lecture you will be able to

- Specify a structural nested model for a continuous outcome
- Conduct g-estimation to estimate the parameters of a structural nested model
- Describe the relation between g-estimation and instrumental variable estimation

□ Key concepts

- G-estimation
- Structural nested mean models

Study population

- ☐ 1629 cigarette smokers
- ☐ Aged 25-74 years when interviewed in 1971-75 (baseline)
- ☐ Interviewed again in 1982
- ☐ Known sex, age, race, weight, height, education, alcohol use, and smoking intensity at both baseline and follow-up visits, and who answered the general medical history questionnaire at baseline

G-estimation

3

Key variables

Treatment A	Quit smoking between baseline and 1982 1: yes, 0: no
Continuous outcome Y	Weight gain, kg Weight in 1982 minus baseline weight Available for 1566 individuals
Baseline (pre-treatment) covariates	Age, sex, race, alcohol use, intensity of smoking, weight...
Censoring C	Missing weight in 1982 1: yes, 0: no

G-estimation

4

Formal version of causal question: First define the counterfactual means

if everybody had quit smoking

- $E[Y^{a=1}]$
- $Y^{a=1}$ is an individual's outcome under $a=1$

if nobody had quit smoking

- $E[Y^{a=0}]$
- $Y^{a=0}$ is an individual's outcome under $a=0$

Then the formal question is:

- What is the average causal effect $E[Y^{a=1}] - E[Y^{a=0}]$?
 - ignore censoring for now

G-estimation

5

Average causal effect can be defined marginally or conditionally

- Marginal: Effect in the entire population
 - $E[Y^{a=1}] - E[Y^{a=0}]$
 - Methods: Standardization, IP weighting of MSMs, Instrumental variable estimation (under homogeneity)
- Conditional: Effects in subsets defined by effect modifiers
 - $E[Y^{a=1}|V=v] - E[Y^{a=0}|V=v]$ for all values l
 - All of the above combined with stratification
- Conditional: Effects in subsets defined by *all* confounders
 - $E[Y^{a=1}|L=l] - E[Y^{a=0}|L=l]$ for all values l
 - Methods: stratification, outcome regression, and g-estimation of structural nested models

G-estimation

6

Footnote: Average causal effects can also be conditional on PS

- Just replace the individual covariates L by the propensity score PS
 - $E[Y^{a=1}|PS=p] - E[Y^{a=0}|PS=p]$ for all values p
 - Methods: stratification and outcome regression using PS rather than L

Some new terms

- Structural nested models:
 - Models with parameter(s) that can be interpreted as the magnitude of the causal effect conditional on covariates
- G-estimation:
 - Method to estimate the parameters of structural nested models

Plan for this lecture

A. Review

- Conditional exchangeability

B. Structural nested models

C. G-estimation

D. Adjustment for selection bias

E. G-estimation and IV estimation

G-estimation

9

Conditional exchangeability



Formal definition


$$Y^a \perp\!\!\!\perp A | L = l \text{ for all } a$$

- Conditional exchangeability is equivalent to randomization within levels of the variables in L
 - e.g., smoking cessation occurred at random within levels of age, sex, race, smoking intensity, etc.
- Also known as the assumption of no unmeasured confounding

G-estimation

10





Clarification: Average causal effect for continuous outcomes

- Only mean exchangeability is required
 - within levels of the covariates L , treated individuals would have had the same mean outcome as the untreated individuals had they being untreated, and vice versa
 - mean counterfactual outcome is the same in the treated and the untreated with the same value of L
- For dichotomous outcomes, exchangeability and mean exchangeability are the same

An equivalent expression of conditional exchangeability

$$Y^a \perp\!\!\!\perp A | L = l \text{ for all } a$$

$$\begin{aligned} \Pr[A = 1 | L = l, Y^{a=1}] &= \Pr[A = 1 | L = l, Y^{a=0}] \\ &= \Pr[A = 1 | L = l] \end{aligned}$$

- That is, conditional on the measured covariates, the probability of treatment does not depend on the value of the counterfactual outcome

G-estimation

15

Dream with me for a minute

- Suppose conditional exchangeability given L holds
- Suppose we know the values of the counterfactual outcomes $Y^{a=1}$ and $Y^{a=0}$ for all subjects
- Suppose that we fit a logistic model for treatment that includes one of the counterfactual outcomes as a covariate
 - $\text{logit } \Pr[A=1 | L, Y^a] = \alpha_0 + \alpha_1 Y^a + \alpha_2 L$
 - For either $a=0$ or $a=1$

G-estimation

16

What would be the value of
the coefficient α_1 for $Y^{a=0}$?

0

$$\text{logit Pr}[A = 1|L, Y^{a=0}] = \alpha_0 + \alpha_1 Y^{a=0} + \alpha_2 L$$

Zero

0%

Greater than zero

0%

Less than zero

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

What would be the value of
the coefficient α_1 for $Y^{a=1}$?

0

$$\text{logit Pr}[A = 1|L, Y^{a=1}] = \alpha_0 + \alpha_1 Y^{a=1} + \alpha_2 L$$

Zero

0%

Greater than zero

0%

Less than zero

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

We can test whether conditional exchangeability holds by fitting a propensity score model with counterfactual outcomes as covariates and then checking the value of α_1 .

0

True

0%

False

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Reminder: Different notation, different concepts

- Conditional exchangeability of treated and untreated is represented by

$$Y^a \perp\!\!\!\perp A | L = l$$

- Conditional independence of treatment and outcome is represented by

$$Y \perp\!\!\!\perp A | L = l$$

Under the sharp null, these two statements are expected to be true:

$$Y^0 \perp\!\!\!\perp A | L = l$$

$$Y^1 \perp\!\!\!\perp A | L = l$$

where L is the vector of measured variables in our study

0

True

0%

False

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Plan

A. Review

- Conditional exchangeability assumption

B. Structural nested models

C. G-estimation

D. Adjustment for selection bias

E. G-estimation and IV estimation

Average causal effect within levels of the confounders

- These conditional effects are represented by
 - $E[Y^{a=1}|L=l] - E[Y^{a=0}|L=l]$ for all values l
- Or, equivalently
 - $E[Y^{a=1} - Y^{a=0}|L=l]$
- Outcome regression
 - models for the conditional mean outcome
 $E[Y^a|L=l] = E[Y|A=a, L=l]$
- Structural nested models
 - models for the conditional average effect $E[Y^{a=1} - Y^{a=0}|L=l]$

G-estimation

23

Outcome regression One dichotomous confounder L

- If effect modification

$$E[Y|A=a, L=l] = \beta_0 + \beta_1 A + \beta_2 AL + \beta_3 L$$

- Effect in stratum $L=0$: β_1
- Effect in stratum $L=1$: $\beta_1 + \beta_2$

- If no effect modification

$$E[Y|A=a, L=l] = \beta_0 + \beta_1 A + \beta_3 L$$

- Effect in all strata : β_1

G-estimation

24

Structural nested mean model

One dichotomous confounder L

☐ If effect modification

$$E[Y^a - Y^{a=0} | A=a, L=l] = \beta_1 a + \beta_2 aL$$

- Effect in stratum $L=0$: β_1
- Effect in stratum $L=1$: $\beta_1 + \beta_2$

☐ If no effect modification

$$E[Y^a - Y^{a=0} | A=a, L=l] = \beta_1 a$$

- Effect in all strata : β_1

G-estimation

25

Structural nested models have fewer parameters than outcome regression models

☐ Because SNMs model the causal effect directly

- That is, the difference in mean outcomes rather than the mean outcomes

☐ No intercept and no parameters for the confounders

☐ Fewer opportunities of having bias due to model misspecification

G-estimation

26

Structural nested models vs. Marginal structural models

- For time-varying treatments, structural nested models allow product terms between treatment and time-varying covariates
 - marginal structural models do not
- For point treatments (like in our data), both classes of models can be used
 - MSMs estimate marginal effect or conditional effects, SNMs estimate conditional effects only

G-estimation

27

Plan

- A. Review
 - Conditional exchangeability assumption
- B. Structural nested models
- C. G-estimation
- D. Adjustment for selection bias
- E. G-estimation and IV estimation

G-estimation

28

G-estimation

- The procedure to estimate the parameters of structural nested models
- Next we describe, in 5 steps, g-estimation to estimate β_1 in a structural nested mean model
 - $E[Y^a - Y^{a=0} | A=a, L=l] = \beta_1 a$
 - under no effect modification by any confounders
 - Aside: in the presence of EM, there are multiple parameters to be identified: $E[Y^a - Y^{a=0} | A = a, L = l] = \beta_1 a + \beta_2 aL$

G-estimation

29

Step 1: Mean model transformed into an individual-level, rank-preserving model

$$Y^a - Y^{a=0} = \psi_1 a$$

- This step is not really necessary for g-estimation but simplifies teaching the method
- This rank-preserving model has stronger assumptions than the mean model
- The estimate of ψ_1 obtained via g-estimation for the rank-preserving model is also valid for the mean model

G-estimation

30

This model assumes constant treatment effect

$$Y^a - Y^{a=0} = \psi_1 a$$

That is, the individual causal effect ψ_1 is the same for all individuals and equal to the average causal effect β_1

0

True

0%

False

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Step 2: Some rearranging

$$Y^{a=0} = Y^a - \psi_1 a$$

- We'll see why later
- Still too many unknowns in the model

Step 3: Model linked to the observed data

$$Y^{a=0} = Y - \psi_1 A$$

- The observed outcome Y equals the counterfactual outcome Y^a when a is equal to the observed treatment A
 - By consistency of counterfactuals
- How can we validly estimate ψ_1 ?

G-estimation

33

Step 4: Suppose that a friend of yours knows the value of ψ_1

- But they only tell you that ψ_1 is one of the following
 - $\psi^\dagger = -1.11$, $\psi^\dagger = 0$, or $\psi^\dagger = 3.46$
- How can you identify the true value ψ_1 among the 3 possible values ψ^\dagger ?
- For each subject, compute $H(\psi^\dagger) = Y - \psi^\dagger A$ for each of the three possible values ψ^\dagger
- The newly created variables $H(-1.11)$, $H(0)$, and $H(3.46)$ are candidate counterfactuals
 - One of them is the counterfactual outcome $Y^{a=0}$
 - $H(\psi^\dagger) = Y^{a=0}$ if $\psi^\dagger = \psi_1$

G-estimation

34

Computing $H(\psi^\dagger)$ from our observed data

$$Y^{a=0} = Y - \psi_1 A$$

$$H(\psi^\dagger) = Y - \psi^\dagger A$$

Recall that ψ_1 is one of the following:

$\psi^\dagger = -1.11$, $\psi^\dagger = 0$, or $\psi^\dagger = 3.46$

ID	A	L	Y
1	0	0	3.5
2	1	1	1.2
3	1	1	2.7
...

G-estimation

35

Computing $H(\psi^\dagger)$ from our observed data

$$Y^{a=0} = Y - \psi_1 A$$

$$H(\psi^\dagger) = Y - \psi^\dagger A$$

Recall that ψ_1 is one of the following:

$\psi^\dagger = -1.11$, $\psi^\dagger = 0$, or $\psi^\dagger = 3.46$

ID	A	L	Y	$H(\psi^\dagger = -1.11)$
1	0	0	3.5	$3.5 - (-1.11)(0) = 3.5$
2	1	1	1.2	$1.2 - (-1.11)(1) = 2.31$
3	1	1	2.7	$2.7 - (-1.11)(1) = 3.81$
...

G-estimation

36

Computing $H(\psi^\dagger)$ from our observed data

$$Y_{a=0} = Y - \psi_1 A$$

$$H(\psi^\dagger) = Y - \psi^\dagger A$$

Recall that ψ_1 is one of the following:

$\psi^\dagger = -1.11$, $\psi^\dagger = 0$, or $\psi^\dagger = 3.46$

ID	A	L	Y	$H(\psi^\dagger=-1.11)$	$H(\psi^\dagger=0)$
1	0	0	3.5	$3.5 - (-1.11)(0) = 3.5$	$3.5 - 0(0) = 3.5$
2	1	1	1.2	$1.2 - (-1.11)(1) = 2.31$	$1.2 - 0(1) = 1.2$
3	1	1	2.7	$2.7 - (-1.11)(1) = 3.81$	$2.7 - 0(1) = 2.7$
...

G-estimation

37

Computing $H(\psi^\dagger)$ from our observed data

$$Y_{a=0} = Y - \psi_1 A$$

$$H(\psi^\dagger) = Y - \psi^\dagger A$$

Recall that ψ_1 is one of the following:

$\psi^\dagger = -1.11$, $\psi^\dagger = 0$, or $\psi^\dagger = 3.46$

ID	A	L	Y	$H(\psi^\dagger=-1.11)$	$H(\psi^\dagger=0)$	$H(\psi^\dagger=3.46)$
1	0	0	3.5	$3.5 - (-1.11)(0) = 3.5$	$3.5 - 0(0) = 3.5$	$3.5 - 3.46(0) = 3.5$
2	1	1	1.2	$1.2 - (-1.11)(1) = 2.31$	$1.2 - 0(1) = 1.2$	$1.2 - 3.46(1) = -2.26$
3	1	1	2.7	$2.7 - (-1.11)(1) = 3.81$	$2.7 - 0(1) = 2.7$	$2.7 - 3.46(1) = -0.76$
...

G-estimation

38

Step 5: Test all the candidates, choose the true value

- To identify which candidate $H(\psi^\dagger)$ is the counterfactual $Y^{a=0}$, fit 3 models

$$\text{logit Pr}[A=1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(-1.11) + \alpha_2 L$$

$$\text{logit Pr}[A=1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(0) + \alpha_2 L$$

$$\text{logit Pr}[A=1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(3.46) + \alpha_2 L$$

- Remember the assumption of conditional exchangeability?

- and its relation to the parameter α_1 in the model

$$\text{logit Pr}[A=1|L, Y^{a=0}] = \alpha_0 + \alpha_1 Y^{a=0} + \alpha_2 L$$

G-estimation

39

Which candidate $H(\psi^\dagger)$ is equal to the counterfactual outcome?

Recall: $\text{logit Pr}[A = 1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(\psi^\dagger) + \alpha_2 L$

0

Come on. This is hopeless.

0%

The candidate $H(\psi^\dagger)$ with $\alpha_1=0$

0%

The candidate $H(\psi^\dagger)$ with $\alpha_1>0$

0%

The candidate $H(\psi^\dagger)$ with $\alpha_1<0$

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Which ψ^\dagger is equal to the causal parameter ψ_1 ?

Recall: $\text{logit Pr}[A = 1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(\psi^\dagger) + \alpha_2 L$

0

(A) The ψ^\dagger with $\alpha_1=0$ when used to compute the candidate $H(\psi^\dagger)$

0%

(B) The ψ^\dagger with $\alpha_1>0$ when used to compute the candidate $H(\psi^\dagger)$

0%

(C) The ψ^\dagger with $\alpha_1<0$ when used to compute the candidate $H(\psi^\dagger)$

0%

Start the presentation to see live content. For screen share software, share the entire screen. Get help at pollev.com/app

Done: That was g-estimation

- ☐ Search over all possible values of ψ^\dagger until you find one that results in $H(\psi^\dagger)$ with $\alpha_1=0$
 - The candidate $H(\psi^\dagger)$ with $\alpha_1=0$ is the counterfactual and the corresponding ψ^\dagger is the true value ψ_1
- ☐ In practice, use more sophisticated search methods
 - Need some additional coding
- ☐ Sometimes the estimator has closed form
 - No need to search or optimize
 - e.g., linear models like the ones considered here

To get the 95% confidence interval for ψ_1

- ☐ Invert the test for $\alpha_1=0$
 - Find the set of values of ψ^\dagger that result in a P -value > 0.05 when testing for $\alpha_1=0$
 - The 95% confidence limits are the limits of that set of values
 - Can use Wald test, Score test...

- ☐ Bootstrapping

G-estimation

43

Back to our data: Point estimate for ψ_1

- ☐ Point estimate: 3.5 *See gestimation.R, lines 19-42*
 - Value of ψ^\dagger that produces the value of α_1 closest to 0
 - Or, equivalently, p-value closest to 1
- ☐ Adjusting for baseline covariates:
 - Sex, age, race, education, smoking intensity, duration of smoking, exercise, active lifestyle, baseline weight

G-estimation

44

95% confidence interval for estimate of ψ_1

- 95% CI: 2.6, 4.4 approx
- By inversion of the test: values of ψ^\dagger for which the test for $\alpha_1=0$ has $P\text{-value} > 0.05$ in model
 - We might not be able to find the value of ψ^\dagger that corresponds to exactly $p\text{-value}=0.05$
 - Select next closest ψ^\dagger with $p\text{-value} < 0.05$ to give us a (conservative) 95% CI

See *gestimation.R*, lines 19-42



ψ_1	p-value
2.5	0.028
2.6	0.049
2.7	0.081
...	...
3.4	0.888
3.5	0.929
3.6	0.750
...	...
4.2	0.091
4.3	0.055
4.4	0.032

G-estimation

45

More efficient g-estimators exist (narrower 95% CIs)


- Doubly-robust estimators are more efficient
 - Require a model for the mean of $H(\psi^\dagger)$ given the confounders L
- G-estimation based on functions of $H(\psi^\dagger)$ may be more efficient
 - e.g., we could use the model

$$\text{logit Pr}[A=1|L, H(\psi)] = \alpha_0 + \alpha_1 [H(\psi^\dagger)]^3 + \alpha_2 L$$

G-estimation

46

Causal questions of interest

1. What is the effect of smoking cessation on weight gain?
 2. What is the effect of smoking cessation on risk of death?
- 
- G-estimation cannot be generally used with logistic models, particularly in the time-varying setting
 - Can use log models under rare disease assumption

G-estimation

47

Plan

- A. Review
 - Conditional exchangeability assumption
- B. Structural nested models
- C. G-estimation
- D. Adjustment for selection bias
- E. G-estimation and IV estimation

G-estimation

48

Key variables

Treatment A	Quit smoking between baseline and 1982 1: yes, 0: no
Continuous outcome Y	Weight gain, kg Weight in 1982 minus baseline weight Available for 1566 individuals
Baseline (pre-treatment) covariates	Age, sex, race, alcohol use, intensity of smoking, weight...
Censoring C	Missing weight in 1982 1: yes, 0: no

G-estimation

49

Causal question

- ☐ Estimate the counterfactual mean weight gain if nobody's outcome had been censored
 - $E[Y^{a=1, c=0}]$
 - $E[Y^{a=0, c=0}]$
- ☐ Estimate the average causal effect
 - $E[Y^{a=1, c=0}] - E[Y^{a=0, c=0}]$
- ☐ Conditional on confounders when using g-estimation

G-estimation

50

G-estimation + IP weighting

- G-estimation to adjust for confounding
- IP weighting to adjust for selection bias

- First, use IP weighting to estimate IP weights W^C
 - The weights create a pseudo-population without selection bias by measured covariates
 - i.e., no arrows from L and A into C

G-estimation + IP weighting

- Second, use g-estimation to adjust for confounding in the pseudo-population
 - Find the value of ψ that produces the value of α_1 closest to 0 in the IP weighted model
$$\text{logit Pr}[A=1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(\psi^\dagger) + \alpha_2 L$$
 - using estimates of the weights W^C

- Need to use robust variance estimator
 - Conservative 95% confidence interval

Back to our data: Point estimate for ψ_1

- Point estimate: 3.4 *See gestimation.R, lines 44-104*
- 95% CI: 2.5, 4.5 approx
- Adjusting for baseline covariates:
 - Sex, age, race, education, smoking intensity, duration of smoking, exercise, active lifestyle, baseline weight

Plan

- A. Review
 - Conditional exchangeability assumption
- B. Structural nested models
- C. G-estimation
- D. Adjustment for selection bias
- E. G-estimation and IV estimation

G-estimation

to consistently estimate the parameter ψ_1

- From the model $Y^{a=0} = Y - \psi_1 A$
 - Compute $H(\psi^\dagger) = Y - \psi^\dagger A$
 - for a range of possible values ψ^\dagger
 - The variables $H(\psi^\dagger)$ are candidate counterfactuals
 - One of them is the counterfactual outcome $Y^{a=0}$
 - $H(\psi^\dagger) = Y^{a=0}$ if $\psi^\dagger = \psi_1$
 - We just need to identify the value ψ^\dagger that makes $H(\psi^\dagger) = Y^{a=0}$
-

G-estimation

55

How to identify the value ψ^\dagger
that makes $H(\psi^\dagger) = Y^{a=0}$?

- We said that **IF** conditional exchangeability holds

$$Y^a \perp\!\!\!\perp A | L = l$$

- Then ψ_1 is the value ψ^\dagger that makes $\alpha_1=0$ in the model
 $\text{logit } \Pr[A=1|L, H(\psi^\dagger)] = \alpha_0 + \alpha_1 H(\psi^\dagger) + \alpha_2 L$
 - But there is another way
 - If we have an instrumental variable Z
-

G-estimation

56

How to identify the value of ψ
that makes $H(\psi^\dagger) = Y^{a=0}$?

- **By definition** of IV, (unconditional) exchangeability is expected to hold

$$Y^a \perp\!\!\!\perp Z$$

- Then the true value of ψ is the one that makes $\alpha_1=0$ in the model $\text{logit Pr}[Z=1|H(\psi)] = \alpha_0 + \alpha_1 H(\psi^\dagger)$

IV estimation as a
particular case of g-estimation

- G-estimation based on the IV is mathematically equivalent to the standard IV estimand
 - Defined in last week's notes
- The standard IV estimator consistently estimates the parameter ψ^* from our structural mean model
 - Under instrumental assumptions (i)-(iii) and assumption of no additive effect modification (iv)
 - See Chapter 16 of *Causal Inference: What If*.

Within the g-estimation framework, IV becomes a more general method that allows

- ☐ Use of baseline covariates
 - To increase efficiency
 - ☐ Especially if covariates are strongly associated with Y and weakly associated with A
 - To address violations of condition (iii)
- ☐ Use of multiplicative models and assumption of no multiplicative effect modification (iv)
- ☐ Extensions to time-varying treatments and time-varying instruments, survival analyses

Conclusion

- ☐ We have studied (essentially) all available methods to adjust for confounding
- ☐ Of those, IP weighting, standardization/g-formula, and g-estimation can be generally used with time-varying treatments

IP weighting, standardization, or g-estimation?

☐ Why do we have to choose?

- If possible, use all methods separately and doubly-robust versions when available
- Similar effect estimates increase our confidence in the results
- Different effect estimates will make us examine our modeling assumptions

☐ See EPI207 for more about these methods

G-estimation

61

Readings

☐ *Causal Inference: What If*. Chapter 14

G-estimation

62

Progress report

1. Introduction to modeling
2. Stratified analysis:
 - Outcome regression
 - Propensity scores
3. Standardization
4. Inverse probability weighting
 - Marginal structural models
5. Instrumental variable estimation
6. G-estimation
7. Causal survival analysis