

**UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE FÍSICA DE SÃO CARLOS**

Camila Raupp da Luz

Thermal Relaxation Asymmetry in Quantum Systems

São Carlos

2026

Camila Raupp da Luz

Thermal Relaxation Asymmetry in Quantum Systems

Dissertation presented to the Graduate Program in Physics at the Instituto de Física de São Carlos da Universidade de São Paulo, to obtain the degree of Master in Science.

Concentration area: Theoretical and Experimental Physics

Advisor: Dr. Ivan Medina

Coadvisor: Prof. Dr. Diogo O. Soares-Pinto

Original version

São Carlos

2026

É possível elaborar a ficha catalográfica em LaTeX ou incluir a fornecida pela Biblioteca. Para tanto observe a programação contida nos arquivos USPSC-modelo.tex e fichacatalografica.tex e/ou gere o arquivo fichacatalografica.pdf.

A biblioteca da sua Unidade lhe fornecerá um arquivo PDF com a ficha catalográfica definitiva, que deverá ser salvo como fichacatalografica.pdf no diretório do seu projeto.

ERRATA

A errata é um elemento opcional, que consiste de uma lista de erros da obra, precedidos pelas folhas e linhas onde eles ocorrem e seguidos pelas correções correspondentes. Deve ser inserida logo após a folha de rosto e conter a referência do trabalho para facilitar sua identificação, conforme a ABNT NBR 14724 (?).

Modelo de Errata:

RAUPP, C. **Thermal Relaxation Asymmetry in Quantum Systems**. 2026. 61 p. Dissertação (Mestrado em Ciências) - Instituto de Física de São Carlos, Universidade de São Paulo, São Carlos, 2026.

ERRATA

Folha	Linha	Onde se lê	Leia-se
1	10	auto-conclavo	autoconclavo

Folha de aprovação em conformidade
com o padrão definido
pela Unidade.

No presente modelo consta como
folhadeaprovacao.pdf

Este trabalho é dedicado aos alunos da USP, como uma contribuição das Bibliotecas do Campus USP de São Carlos para o desenvolvimento e disseminação da pesquisa científica da Universidade.

ACKNOWLEDGEMENTS

Primeira frase do agradecimento

Segunda frase

Outras frases

Última frase

*“O estudo, a busca da verdade e da beleza são domínios
em que nos é consentido sermos crianças por toda a vida.”*

Albert Einstein

ABSTRACT

RAUPP, C. **Thermal Relaxation Asymmetry in Quantum Systems**. 2026. 61 p.
Dissertation (Master in Science) - Instituto de Física de São Carlos, Universidade de São Paulo, São Carlos, 2026.

This is the english abstract.

Keywords: LaTeX. USPSC class. Thesis. Dissertation. Conclusion course paper. Report.

RESUMO

RAUPP, C. **Thermal Relaxation Asymmetry in Quantum Systems**. 2026. 61 p. Dissertação (Mestrado em Ciências) - Instituto de Física de São Carlos, Universidade de São Paulo, São Carlos, 2026.

O resumo deve ressaltar o objetivo, o método, os resultados e as conclusões do documento. A ordem e a extensão destes itens dependem do tipo de resumo (informativo ou indicativo) e do tratamento que cada item recebe no documento original. O resumo deve ser precedido da referência do documento, com exceção do resumo inserido no próprio documento. (...) Salientamos que algumas Unidades exigem o título dos trabalhos acadêmicos em inglês, tornando necessário a inclusão das referências nos resumos e abstracts, o que foi adotado no **Modelo para TCC em L^AT_EX utilizando a classe USPSC** e no **Modelo para teses e dissertações em L^AT_EX utilizando a classe USPSC**. As palavras-chave devem figurar logo abaixo do resumo, antecedidas da expressão Palavras-chave:, separadas entre si por ponto e finalizadas também por ponto (?).

Palavras-chave: LaTeX. Classe USPSC. Tese. Dissertação. Trabalho de conclusão de curso (TCC). Relatório.

LIST OF FIGURES

Figure 1 – Model for two interacting qubits with a global bath.	36
Figure 2 – Degree of completion for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.	39
Figure 3 – Statistical velocity for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.	40
Figure 4 – Statistical velocity for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.	40
Figure 5 – Acentuação (modo texto - L ^A T _E X)	59

LIST OF TABLES

Table 1 – Model’s Parameters	36
Table 2 – Exemplo de tabela centralizada verticalmente e horizontalmente	51
Table 3 – Exemplo de tabelas com grade	53

LIST OF FRAMES

LIST OF ABBREVIATIONS AND ACRONYMS

ABNT	Associação Brasileira de Normas Técnicas
abnTeX	ABsurdas Normas para TeX
IBGE	Instituto Brasileiro de Geografia e Estatística
LaTeX	Lamport TeX
USP	Universidade de São Paulo
USPSC	Campus USP de São Carlos

LIST OF SYMBOLS

Γ	Letra grega Gama
Λ	Lambda
ζ	Letra grega minúscula zeta
$\in$	Pertence

CONTENTS

1	INTRODUÇÃO	29
2	OPEN QUANTUM SYSTEMS THEORY	31
3	QUANTUM THERMODYNAMICS	33
3.1	Quantum Correlations	33
3.2	Thermodynamic variables in classical and quantum regime	33
3.3	Equilibrium and Nonequilibrium States	33
3.4	Relaxation, Entropy Production and Irreversibility	33
3.4.1	Thermal Kinematics	33
4	FROM DISCRETE TO CONTINUOUS QUANTUM SYSTEMS	35
4.1	Two Interacting Qubits	35
4.1.1	Thermal Relaxation of Two Interacting Qubits	36
4.2	Gaussian States	41
4.2.1	Thermal Relaxation of Gaussian States	41
4.2.1.1	Asymmetry for Thermal States	41
4.2.1.2	Asymmetry for Displaced States	41
4.2.1.3	Asymmetry for Squeezed States	41
5	CONCLUSÃO	43
	REFERENCES	45
	APPENDIX	47
	APPENDIX A – APÊNDICE(S)	49
	APPENDIX B – EXEMPLO DE TABELA CENTRALIZADA VER- TICALMENTE E HORIZONTALMENTE	51
	APPENDIX C – EXEMPLO DE TABELA COM GRADE	53
	ANNEX	55
	ANNEX A – EXEMPLO DE ANEXO	57

ANNEX B – ACENTUAÇÃO (MODO TEXTO - $\text{\LaTeX}$)	59
INDEX	61

1 INTRODUÇÃO

2 OPEN QUANTUM SYSTEMS THEORY

3 QUANTUM THERMODYNAMICS

3.1 Quantum Correlations

3.2 Thermodynamic variables in classical and quantum regime

3.3 Equilibrium and Nonequilibrium States

3.4 Relaxation, Entropy Production and Irreversibility

3.4.1 Thermal Kinematics

4 FROM DISCRETE TO CONTINUOUS QUANTUM SYSTEMS

Nature has a diversity of physical phenomena, being some better described by discrete-variable models while others are better described by continuous-variable models. Quantum mechanics was formulated in both discrete and continuous regimes as the Heisenberg and Schrödinger frameworks, which, despite of having fundamentally different mathematical resources, complements each other when explaining experimental results. Finding the same phenomena in different regimes not only brings to light fundamental aspects of nature but also increases the possibility of technological applications. For these reasons, this chapter explores the thermal relaxation asymmetry for discrete and continuous-variable quantum systems.

Given that this work aims to understand how quantum resources affect the heating and cooling asymmetry, we examine specific models that have these resources. Among a variety of possible discrete models, we chose the two interacting qubits in contact with the same bosonic bath. This model allow us to analyse the effects of quantum correlations and non-Markovianity in the thermal relaxation. While for continuous-variables models, the natural choice was Gaussian states for its simplicity and experimental accessibility. For this case, the quantum resources we explore are displacement and squeezing of thermal states.

4.1 Two Interacting Qubits

The interaction between N qubits with the same energy difference ($\hbar$) can be described by a family of Hamiltonians

$$H_S = -\hbar \sum_{j=1}^N \sigma_j^z - J \sum_{j=1}^{N-1} [(1 + \gamma) \sigma_j^x \sigma_{j+1}^x + (1 - \gamma) \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z] \quad (4.1)$$

where σ_k^x , σ_k^y , and σ_k^z are the Pauli matrices for the k -th qubit. J is the coupling parameter between the qubits, while γ and Δ creates anisotropies in the system. Depending on the values of the anisotropic parameters, we can distinguish different models (1) as shown in Table 1.

Table 1 – Model's Parameters

Parameters	Models
$\Delta = 1, \gamma = 0$	XXX
$\Delta \neq 0, \gamma = 0$	XXZ
$\Delta \neq 0, -1 < \gamma < 1$	XYZ
$\Delta = 0, \gamma = 0$	XX
$\Delta = 0, -1 < \gamma < 1$	XY
$\Delta = 0, \gamma = \pm 1$	Transverse Field Ising (TFI)

What matters for us about this models is that the different interaction cases will lead to different correlations between the qubits. These correlations will be measured by the mutual information and entanglement of formation, defined in Section 3.1. Note that not all of these models can create entanglement, which we will discuss later in this section.

4.1.1 Thermal Relaxation of Two Interacting Qubits

Lets start considering two qubits coupled to the same bosonic bath, as shown in Figure 1. In this first analysis, the qubits has an energy gap of $\frac{\omega_0}{2}$ and exchange energy according to the model XX (with $\Delta = 0$ and $\gamma = 0$), meaning it is isotropic. One qubit will heat up from the temperature T_c while the other will cool down from the temperature T_h , being the bath at an intermediate temperature T_w .




Figure 1 – Model for two interacting qubits with a global bath.

Depending on the initial state of the system, the relaxation process will be different. In our case, we consider the same qubit's energy gap (ω_0) and decaying rate (Γ) respecting the weak coupling regime ($\Gamma \ll \omega_0$). Thus, the only variable that characterized the qubit's initial state is the temperature. We could add coherences to the qubits, but this would make the terms 'heating' (or 'cooling') inaccurate due to the inexistence of thermal states. Therefore, we will keep the qubits locally thermal while adding global correlations if desired.

To make a fair comparison between heating and cooling, we need to find statistically equivalent initial states with different temperatures. Because the asymmetry appears only for relaxation of nonequilibrium states, it is natural to distinguish them according to how far from equilibrium it is. Thereby, the relative entropy, which is related to the nonequilibrium free energy, seems the most appropriate choice of distinguishability measure. That said, statistically equivalent initial states for each qubit must have the same initial relative entropy from the local steady state of the dynamics (Equation 4.2). However, to find the appropriate initial temperature, first, we must find the local steady state (ρ_{ss}^{qubit}).

$$K[\rho_c || \rho_{ss}^{qubit}] = K[\rho_h || \rho_{ss}^{qubit}] \quad (4.2)$$

According to its definition, a steady state does not change in time, which mathematically can be found by solving $\mathcal{L}(\rho_{ss}) = 0$, where ρ_{ss} is the global steady state. Despite this being the most traditional approach, there is another simpler way. First, we know that the steady state of a Davies Map is always a Gibbs state, which allows us to assume that

$$\rho_{ss} = \frac{e^{-\beta_w H_S}}{\mathcal{Z}}, \quad (4.3)$$

where β_w is the inverse temperature of the bath and H_S is the total system hamiltonian 4.1. Then, we can trace out one of the qubits* and write the expression in terms of H_S eigenvalues and eigenvectors to obtain

$$\rho_{qubit} = \begin{bmatrix} e^{\beta_w \omega_0} + \cosh(\beta_w J) & 0 \\ 0 & e^{-\beta_w \omega_0} + \cosh(\beta_w J) \end{bmatrix}. \quad (4.4)$$

Because every two-level system in a Gibbs state has the same format, we can generally isolate the qubit's temperature, resulting in

$$\beta_{qubit}^f = \frac{1}{\omega_0} \ln \left(\frac{e^{\beta_w \omega_0} + \cosh(\beta_w J)}{e^{-\beta_w \omega_0} + \cosh(\beta_w J)} \right). \quad (4.5)$$

Note that, even though the global system thermalizes to the bath's temperature, the qubits' local thermal state corresponds to a different temperature. This happens because the interaction between the qubits generates correlations that are erased by the partial trace. Now, we can solve Equation 4.2 and find the initial temperatures.

Because we want to understand how quantum correlations may affect the thermal relaxation asymmetry, the next step is to find out what is the maximum correlation

* It does not matter which one, because both equilibrate to the same temperature

possible for our initial state. Lets assume the initial global state has the form of an X-state,

$$\rho = \rho_c \otimes \rho_h + \chi = \begin{bmatrix} a & 0 & 0 & f \\ 0 & b & e & 0 \\ 0 & e^* & c & 0 \\ f^* & 0 & 0 & d \end{bmatrix}, \quad (4.6)$$

where the diagonal elements are

$$a = \frac{e^{\omega_0(\beta_c + \beta_h)/2}}{\mathcal{Z}_c \mathcal{Z}_h} \quad (4.7)$$

$$b = \frac{e^{\omega_0(\beta_c - \beta_h)/2}}{\mathcal{Z}_c \mathcal{Z}_h} \quad (4.8)$$

$$c = \frac{e^{-\omega_0(\beta_c - \beta_h)/2}}{\mathcal{Z}_c \mathcal{Z}_h} \quad (4.9)$$

$$d = \frac{e^{-\omega_0(\beta_c + \beta_h)/2}}{\mathcal{Z}_c \mathcal{Z}_h} \quad (4.10)$$

where $\mathcal{Z}_c = 2 \cosh(\omega_0 \beta_c / 2)$, $\mathcal{Z}_h = 2 \cosh(\omega_0 \beta_h / 2)$ and the off-diagonal elements are

$$\chi = e |01\rangle \langle 10| + e^* |10\rangle \langle 01| + f |00\rangle \langle 11| + f^* |11\rangle \langle 00|. \quad (4.11)$$

For ρ to be considered a proper density matrix it must satisfy all the properties: (1) hermiticity, (2) unit trace and (3) positive semi-definite. So, we must choose e and f values that preserve property (3) of ρ , which means that all eigenvalues of ρ must be non-negative ($\lambda_i \geq 0$ for all i). When diagonalizing ρ , we get two identical quadratic equations of the form

$$\lambda = \frac{x + y \pm \sqrt{x^2 + y^2 - 2xy + 4z}}{2} \geq 0 \quad (4.12)$$

where the variables can be $x = (b, a)$, $y = (c, d)$, $z = (|e|^2, |f|^2)$. Because the positive sign will always satisfy the inequality, we will analyse the restriction this inequality brings when choosing the negative sign. Manipulating the expression, we find $z \leq xy$ that means

$$|e| \leq \frac{1}{\mathcal{Z}_c \mathcal{Z}_h} \quad (4.13)$$

$$|f| \leq \frac{1}{\mathcal{Z}_c \mathcal{Z}_h}. \quad (4.14)$$

Now that we know the initial temperatures and maximum initial correlations, we can solve the Lindblad master equation to obtain the density matrices for the global system and each individual qubit as a function of time. After computing the Quantum Fisher Information, we solve the thermal kinematic's variables as described in Section 3.4.1.

The comparison between the degree of completion of the heating qubit (red) and the cooling qubit (blue) is shown in Figure 2, where the dashed lines represent the case with no interaction between the qubits and the solid lines represent the interacting case. Note that in our results, heating is always faster than cooling, because the red degree of completion is above the correspondent blue lines. Another result we can extract is that, when qubits can interact between themselves, they relaxate faster. Furthermore, the presence of initial correlations seems to delay the dynamics when we compare the continuous lines of both graphs.




Figure 2 – Degree of completion for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.

This same observations can be obtained from the statistical velocity shown in Figure 3. Additionally to this analysis, note that, when the velocity is zero, plateaus appears in the degree of completion given its oscillatory behaviour. A possible interpretation of the oscillations is that the system almost relax to an intermediate state, but something interfere and restart the relaxation from that intermediate state.




Figure 3 – Statistical velocity for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.

Figure 4 show the entropy production rate for each of the studied cases. The hypothesis that heating is faster than cooling because it is more efficient in entropy production (2) can be verified by observing that the red curves presents higher initial entropy production. Also note that the interaction case have negative entropy production for some time intervals. This is a




Figure 4 – Statistical velocity for XX-model. Each line correspond to cooling (blue) or heating (red) processes with (solid) or without (dashed) interaction between qubits.

4.2 Gaussian States

4.2.1 Thermal Relaxation of Gaussian States

4.2.1.1 Asymmetry for Thermal States

4.2.1.2 Asymmetry for Displaced States

4.2.1.3 Asymmetry for Squeezed States

5 CONCLUSÃO

REFERENCES

- 1 MEDINA, I. *et al.* Variational-quantum-eigensolver–inspired optimization for spin-chain work extraction. **Phys. Rev. A**, American Physical Society, v. 110, p. 012443, Jul 2024. Available at: <https://link.aps.org/doi/10.1103/PhysRevA.110.012443>.
- 2 IBÁÑEZ, M. *et al.* Heating and cooling are fundamentally asymmetric and evolve along distinct pathways. **Nature Physics**, v. 20, p. 135–141, Jan 2024. Available at: <https://doi.org/10.1038/s41567-023-02269-z>.

APPENDIX

APPENDIX A – APÊNDICE(S)

Elemento opcional, que consiste em texto ou documento elaborado pelo autor, a fim de complementar sua argumentação, conforme a ABNT NBR 14724 (?).

Os apêndices devem ser identificados por letras maiúsculas consecutivas, seguidas de hífen e pelos respectivos títulos. Excepcionalmente, utilizam-se letras maiúsculas dobradas na identificação dos apêndices, quando esgotadas as 26 letras do alfabeto. A paginação deve ser contínua, dando seguimento ao texto principal. (?)

**APPENDIX B – EXEMPLO DE TABELA CENTRALIZADA
VERTICALMENTE E HORIZONTALMENTE**

A Tabela 2 exemplifica como proceder para obter uma tabela centralizada verticalmente e horizontalmente.

Table 2 – Exemplo de tabela centralizada verticalmente e horizontalmente

Coluna A	Coluna B
Coluna A, Linha 1	Este é um texto bem maior para exemplificar como é centralizado verticalmente e horizontalmente na tabela. Segundo parágrafo para verificar como fica na tabela
Quando o texto da coluna A, linha 2 é bem maior do que o das demais colunas	Coluna B, linha 2

Fonte: Elaborada pelos autores.

APPENDIX C – EXEMPLO DE TABELA COM GRADE

A Tabela 3 exemplifica a inclusão de traços estruturadores de conteúdo para melhor compreensão do conteúdo da tabela, em conformidade com as normas de apresentação tabular do IBGE.

Table 3 – Exemplo de tabelas com grade

Coluna A	Coluna B
A1	B1
A2	B2
A3	B3
A4	B4

Fonte: Elaborada pelos autores.

ANNEX

ANNEX A – EXEMPLO DE ANEXO

Elemento opcional, que consiste em um texto ou documento não elaborado pelo autor, que serve de fundamentação, comprovação e ilustração, conforme a ABNT NBR 14724. (?).

O **ANEXO B** exemplifica como incluir um anexo em pdf.

ANNEX B – ACENTUAÇÃO (MODO TEXTO - $\text{\LaTeX}$)Figure 5 – Acentuação (modo texto - $\text{\LaTeX}$)

$\text{\textbackslash'a}$ - á
 $\text{\textbackslash'a}$ - à
 $\text{\textbackslash~a}$ - ã
 $\text{\textbackslash^a}$ - â
 $\text{\textbackslash'e}$ - é
 $\text{\textbackslash^e}$ - ê
 $\text{\textbackslash'\{i\}}$ - í
 $\text{\textbackslash'I}$ - Í
 $\text{\textbackslash'o}$ - ó
 $\text{\textbackslash~o}$ - õ
 $\text{\textbackslash^o}$ - ô
 $\text{\textbackslash'u}$ - ú
 $\text{\textbackslash"u}$ - ü
 $\text{\textbackslashc{c}}$ - ç
 $\text{\textbackslashc{C}}$ - Ç

Fonte: Comandos [...] (??)

INDEX

tabelas, 51, 53